

Orbital-selective Superconductivity in the Pressurized Bilayer Nickelate $\text{La}_3\text{Ni}_2\text{O}_7$: An Infinite Projected Entangled- Pair State Study



arXiv:2311.05491(2023)

Xing-Yu Zhang

$\text{La}_3\text{Ni}_2\text{O}_7$ Superconduct under pressure






La₃Ni₂O₇ structure






Ni-28: [Ar] 3d⁸ 4s² or [Ar] 3d⁹ 4s¹

+3 +2.5 -2



La₃Ni₂O₇



La₃Ni₂O₇ orbitals




$d_{x^2-y^2}$ and d_{z^2} which responsible for SC order?



Different parameters for $d_{x^2-y^2}$ or d_{z^2}


$$\begin{aligned} H_{\text{bilayer}} = & -t_{\parallel} \sum_{\langle i,j \rangle, \mu, \sigma} (c_{i,\mu,\sigma}^{\dagger} c_{j,\mu,\sigma} + H.c.) \\ & + J_{\parallel} \sum_{\langle i,j \rangle, \mu} (\mathbf{S}_{i,\mu} \cdot \mathbf{S}_{j,\mu} - \frac{1}{4} n_{i,\mu} n_{j,\mu}) \\ & - t_{\perp} \sum_{i, \sigma} (c_{i,\mu=1,\sigma}^{\dagger} c_{i,\mu=-1,\sigma} + H.c.) \\ & + J_{\perp} \sum_i \mathbf{S}_{i,\mu=1} \cdot \mathbf{S}_{i,\mu=-1}, \end{aligned}$$

$d_{x^2-y^2}$ responsible for SC order





SC order parameter $\Delta_z = \frac{1}{\sqrt{2}} \langle \sum_{\mu=\pm 1} c_{i,\mu,\uparrow}^\dagger c_{i,-\mu,\downarrow}^\dagger \rangle$


Tuning parameters



$R_3Ni_2O_7$





$$d_{x^2-y^2} \quad t_{\parallel} = 3, J_{\parallel} = 1, t_{\perp} = 0, J_{\perp} = 2$$



Problems:

1. VUMPS doesn't converge
2. Up and down environment converges but the energy is not good
3. Local minimum when n approach to 1



Thank you for listening!

Q&A?