



BAYESIAN INFERENCE

Antoine Deleforge



THANKS



**Prof. Dr.-Ing. Walter
Kellermann**
Head of LMS chair
EARS project coordinator



Dr. Roland Maas
PhD at LMS chair
Research and
development at Amazon



OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- Bayesian inference by examples
 - Direct inference
 - The Expectation-Maximization algorithm
 - Variational Bayes methods
- Markov chain Monte Carlo

OUTLINE

- **What is Bayesian inference?**
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- Bayesian inference by examples
 - Direct inference
 - The Expectation-Maximization algorithm
 - Variational Bayes methods
- Markov chain Monte Carlo

What is Bayesian Inference?

Inference

What is Bayesian Inference?

Inference

Ingredients



What is Bayesian Inference?

Inference

Ingredients



Goals

Estimation Quantitative deductions on causes or consequences of the observations, i.e., find underlying model parameters.

Example: *I observed a certain amount of rain drops forming on my window in the last minute. What is the current rainfall in milimeters?*



What is Bayesian Inference?

Inference

Ingredients



Goals

Estimation Quantitative deductions on causes or consequences of the observations, i.e., find underlying model parameters.

Example: *I observed a certain amount of rain drops forming on my window in the last minute. What is the current rainfall in milimeters?*

Prediction From the inferred model, predict what missing or future observations should be.


Example: *How many more raindrops will form on my window in the next hour?*



What is Bayesian Inference?

Inference

Ingredients



Goals

Estimation Quantitative deductions on causes or consequences of the observations, i.e., find underlying model parameters.

Example: *I observed a certain amount of rain drops forming on my window in the last minute. What is the current rainfall in milimeters?*

Prediction From the inferred model, predict what missing or future observations should be.

Example: *How many more raindrops will form on my window in the next hour?*

Decision Take a decision out of a discrete set of choices

Example: *Is it safe to open my window 1 minute to get some fresh air?*

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition		
Features		
Tools		

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition	{ <p>The model parameters which should be estimated are considered as unknown constant.</p>	
Features	{	
Tools	{	

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition	{ The model parameters which should be estimated are considered as unknown constant .	The model parameters are considered as hidden random variables , following a hypothetical probabilistic model .
Features	{	
Tools	{	

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition	The model parameters which should be estimated are considered as unknown constant .	The model parameters are considered as hidden random variables , following a hypothetical probabilistic model .
Features	Inference entirely based on observed data and frequentist arguments . Useful when few prior knowledge exist on the underlying random process.	
Tools		

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition	The model parameters which should be estimated are considered as unknown constant .	The model parameters are considered as hidden random variables , following a hypothetical probabilistic model .
Features	Inference entirely based on observed data and frequentist arguments . Useful when few prior knowledge exist on the underlying random process.	Incorporate prior knowledge on the hidden variables in the form of a generative probabilistic model. Useful when some reasonable probability density functions (PDFs) can be assumed.
Tools		

What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition	The model parameters which should be estimated are considered as unknown constant .	The model parameters are considered as hidden random variables , following a hypothetical probabilistic model .
Features	Inference entirely based on observed data and frequentist arguments . Useful when few prior knowledge exist on the underlying random process.	Incorporate prior knowledge on the hidden variables in the form of a generative probabilistic model. Useful when some reasonable probability density functions (PDFs) can be assumed.
Tools	<ul style="list-style-type: none"> • Linear estimators • First and second order statistics • A lot of: $\mathbb{E}\{\dots\}$ 	


What is Bayesian Inference?

Statistics: Inference from the real world observations of a random phenomenon using probability theory

	Classical/Frequentist Statistics	Bayesian Statistics
Definition	The model parameters which should be estimated are considered as unknown constant .	The model parameters are considered as hidden random variables , following a hypothetical probabilistic model .
Features	Inference entirely based on observed data and frequentist arguments . Useful when few prior knowledge exist on the underlying random process.	Incorporate prior knowledge on the hidden variables in the form of a generative probabilistic model. Useful when some reasonable probability density functions (PDFs) can be assumed.
Tools	<ul style="list-style-type: none"> • Linear estimators • First and second order statistics • A lot of: $\mathbb{E}\{\dots\}$ 	<ul style="list-style-type: none"> • Bayes' Theorem • Explicit PDFs • A lot of: $\mathcal{N}(\dots)$

What is Bayesian Inference?


- Bayes' Theorem



$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$

What is Bayesian Inference?

- Bayes' Theorem




\mathbf{X} : Observed variables

\mathbf{Z} : Hidden variables

$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$

What is Bayesian Inference?

- Bayes' Theorem



\mathbf{X} : Observed variables


\mathbf{Z} : Hidden variables

Posterior

$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$

What is Bayesian Inference?

- Bayes' Theorem



X : Observed variables

Posterior


$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$

Z : Hidden variables

Likelihood

What is Bayesian Inference?

- Bayes' Theorem



\mathbf{X} : Observed variables

Posterior

$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$


\mathbf{Z} : Hidden variables

Likelihood

Prior

What is Bayesian Inference?

- Bayes' Theorem



\mathbf{X} : Observed variables

Posterior

$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$

\mathbf{Z} : Hidden variables


Likelihood

« Observed data » or
« marginal » likelihood

Prior

What is Bayesian Inference?

- Bayes' Theorem



\mathbf{X} : Observed variables

Posterior

$$p(\mathbf{Z} = z | \mathbf{X} = x) = \frac{p(\mathbf{X} = x | \mathbf{Z} = z)p(\mathbf{Z} = z)}{p(\mathbf{X} = x)}$$

\mathbf{Z} : Hidden variables

Likelihood

« Observed data » or
« marginal » likelihood


Prior

Remark 1: *Bayes does not « forbid » model parameters !*

- No formal difference between a parameter and a hidden variable with constant prior
- Priors distributions often have parameters called « hyperparameters »

What is Bayesian Inference?

- Bayes' Theorem




$$\begin{array}{ccc}
 \mathbf{X} : \text{Observed variables} & \mathbf{Z} : \text{Hidden variables} & \\
 \text{Posterior} & \text{Likelihood} & \text{Prior} \\
 | & | & | \\
 p(\mathbf{Z} = z | \mathbf{X} = x; \boldsymbol{\theta}) & = \frac{p(\mathbf{X} = x | \mathbf{Z} = z; \boldsymbol{\theta}) p(\mathbf{Z} = z; \boldsymbol{\theta})}{p(\mathbf{X} = x; \boldsymbol{\theta})} \\
 \text{« Observed data » or} & & \\
 \text{« marginal » likelihood} & &
 \end{array}$$

Remark 1: *Bayes does not « forbid » model parameters !*

- No formal difference between a parameter and a hidden variable with constant prior
- Priors distributions often have parameters called « hyperparameters »

What is Bayesian Inference?

- Bayes' Theorem



$$\begin{array}{ccc}
 \textbf{\textit{X}} : \text{Observed variables} & \textbf{\textit{Z}} : \text{Hidden variables} & \\
 \text{Posterior} & \text{Likelihood} & \text{Prior} \\
 | & | & | \\
 p(\textit{Z} = z | \textit{X} = x; \boldsymbol{\theta}) & = \frac{p(\textit{X} = x | \textit{Z} = z; \boldsymbol{\theta}) p(\textit{Z} = z; \boldsymbol{\theta})}{p(\textit{X} = x; \boldsymbol{\theta})} \\
 \text{« Observed data » or} & & \\
 \text{« marginal » likelihood} & &
 \end{array}$$

Remark 1: *Bayes does not « forbid » model parameters !*

- No formal difference between a parameter and a hidden variable with constant prior
- Priors distributions often have parameters called « hyperparameters »

Remark 2: *Why hidden variables?*


- Formally not needed: $p(\textit{X})$ can be obtained by marginalizing out hidden variables
- A convenient and powerful **view point** which makes inference possible in complex scenarios through a variety of methods

What is Bayesian Inference?

An Example:

Observations

X_D	X_C
I see drops on my window	I see a cat or a dog at my window



What is Bayesian Inference?


Bayes' Theorem & Example



An Example:

Observations

X_D	X_C
I see drops on my window	I see a cat or a dog at my window



Hypothesis

$Z = 0$	$Z = 1$	$Z = 2$
It's not raining	It's raining rain	It's raining cats and dogs



What is Bayesian Inference?


Bayes' Theorem & Example



An Example:

Observations

X_D	X_C
I see drops on my window	I see a cat or a dog at my window



Hypothesis

$Z = 0$	$Z = 1$	$Z = 2$
It's not raining	It's raining rain	It's raining cats and dogs



$$p(X_i|Z = j)$$

	X_D	X_C
$Z = 0$	0.1	0.1
$Z = 1$	0.99	0.1
$Z = 2$	0.1	0.99

What is Bayesian Inference?


Bayes' Theorem & Example



An Example:

Observations

X_D	X_C
I see drops on my window	I see a cat or a dog at my window



Hypothesis

$Z = 0$	$Z = 1$	$Z = 2$
It's not raining	It's raining rain	It's raining cats and dogs



$$p(X_i|Z = j)$$

	X_D	X_C
$Z = 0$	0.1	0.1
$Z = 1$	0.99	0.1
$Z = 2$	0.1	0.99

$$p(X_D, X_C|Z = 1) = p(X_D, X_C|Z = 2) = 0.099 : \text{equal likelihood!}$$

What is Bayesian Inference?


Bayes' Theorem & Example



An Example:

Observations

X_D	X_C
I see drops on my window	I see a cat or a dog at my window



Hypothesis

$Z = 0$	$Z = 1$	$Z = 2$
It's not raining	It's raining rain	It's raining cats and dogs



$$p(X_i|Z = j)$$

	X_D	X_C
$Z = 0$	0.1	0.1
$Z = 1$	0.99	0.1
$Z = 2$	0.1	0.99

$$p(X_D, X_C|Z = 1) = p(X_D, X_C|Z = 2) = 0.099 : \text{equal likelihood!}$$

- **Add priors:** $p(Z = 0) = 49.995\%$ $p(Z = 1) = 49.995\%$ $p(Z = 2) = 0.01\%$

What is Bayesian Inference?


Bayes' Theorem & Example



An Example:

Observations

X_D	X_C
I see drops on my window	I see a cat or a dog at my window



Hypothesis

$Z = 0$	$Z = 1$	$Z = 2$
It's not raining	It's raining rain	It's raining cats and dogs



$$p(X_i|Z = j)$$

	X_D	X_C
$Z = 0$	0.1	0.1
$Z = 1$	0.99	0.1
$Z = 2$	0.1	0.99

$$p(X_D, X_C|Z = 1) = p(X_D, X_C|Z = 2) = 0.099 : \text{equal likelihood!}$$

- **Add priors:** $p(Z = 0) = 49.995\%$ $p(Z = 1) = 49.995\%$ $p(Z = 2) = 0.01\%$
- **Bayes' theorem:**

$$p(Z = 1|X_D, X_C) \approx 99.98\% \quad p(Z = 2|X_D, X_C) \approx 0.02\%$$

What is Bayesian Inference?

General Methodology
.....

General Methodology

What is Bayesian Inference?


General Methodology
.....

General Methodology

Modeling


What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology




What is Bayesian Inference?

General Methodology



What is Bayesian Inference?

General Methodology



OUTLINE

- **What is Bayesian inference?**
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- Bayesian inference by examples
 - Direct inference
 - The Expectation-Maximization algorithm
 - Variational Bayes methods
- Markov chain Monte Carlo

OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - Direct inference
 - The Expectation-Maximization algorithm
 - Variational Bayes methods
- Markov chain Monte Carlo

OUTLINE


- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - **Direct inference**
 - **The Expectation-Maximization algorithm**
 - **Variational Bayes methods**
- Markov chain Monte Carlo

Bayesian Inference: Examples


Direct Inference
•••



Bayesian Inference: Examples




Bayesian Inference: Examples



Bayesian Inference: Examples



Bayesian Inference: Examples




Bayesian Inference: Examples

Direct Inference
•••




Bayesian Inference: Examples

Direct Inference
•••



Bayesian Inference: Examples

Direct Inference
•••




Bayesian Inference: Examples

Modeling

Bayesian Inference: Examples

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Bayesian Inference: Examples

Modeling

Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Hidden variable: $Z \in \{1, 2, 3\}$




Bayesian Inference: Examples

Direct Inference
○○○

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Hidden variable: $Z \in \{1, 2, 3\}$



Graphical Model:




Bayesian Inference: Examples

Direct Inference
○○○

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Hidden variable: $Z \in \{1, 2, 3\}$



Graphical Model:



Conditionals


$$\begin{cases} p(\mathbf{X}_n = \mathbf{x}_n | Z = 1) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_1, \mathbf{I}) \\ p(\mathbf{X}_n = \mathbf{x}_n | Z = 2) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_2, \mathbf{I}) \\ p(\mathbf{X}_n = \mathbf{x}_n | Z = 3) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_3, \mathbf{I}) \end{cases}$$

Bayesian Inference: Examples

Direct Inference
○○○

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Hidden variable: $Z \in \{1, 2, 3\}$



Graphical Model:



Conditionals


$$\begin{cases} p(\mathbf{X}_n = \mathbf{x}_n | Z = 1) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_1, \mathbf{I}) \\ p(\mathbf{X}_n = \mathbf{x}_n | Z = 2) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_2, \mathbf{I}) \\ p(\mathbf{X}_n = \mathbf{x}_n | Z = 3) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_3, \mathbf{I}) \end{cases}$$

Bayesian Inference: Examples

Direct Inference
○○○

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Hidden variable: $Z \in \{1, 2, 3\}$



Graphical Model:



Conditionals

$$\begin{cases} p(\mathbf{X}_n = \mathbf{x}_n | Z = 1) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_1, \mathbf{I}) \\ p(\mathbf{X}_n = \mathbf{x}_n | Z = 2) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_2, \mathbf{I}) \\ p(\mathbf{X}_n = \mathbf{x}_n | Z = 3) = \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_3, \mathbf{I}) \end{cases}$$

Priors

$$\begin{cases} p(Z = 1) = 0.1 \text{ (Grandma Jane)} \\ p(Z = 2) = 0.3 \text{ (Student house)} \\ p(Z = 3) = 0.6 \text{ (Family with kids)} \end{cases}$$

Bayesian Inference: Examples

Inference

Bayes' Theorem:

$$p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) = \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_N | Z = i)p(Z = i)}{p(\mathbf{x}_1, \dots, \mathbf{x}_N)}$$

Bayesian Inference: Examples

Inference

Bayes' Theorem:

$$\begin{aligned} p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_N | Z = i)p(Z = i)}{p(\mathbf{x}_1, \dots, \mathbf{x}_N)} \\ &= \frac{\prod_{n=1}^N p(\mathbf{x}_n | Z = i)p(Z = i)}{\sum_{k=1}^3 \prod_{n=1}^N p(\mathbf{x}_n | Z = k)p(Z = k)} \end{aligned}$$

Bayesian Inference: Examples

Inference

Bayes' Theorem:

$$\begin{aligned} p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_N | Z = i)p(Z = i)}{p(\mathbf{x}_1, \dots, \mathbf{x}_N)} \\ &= \frac{\prod_{n=1}^N p(\mathbf{x}_n | Z = i)p(Z = i)}{\sum_{k=1}^3 \prod_{n=1}^N p(\mathbf{x}_n | Z = k)p(Z = k)} \end{aligned}$$

↗ Direct computation

Bayesian Inference: Examples

Inference

Bayes' Theorem:

$$\begin{aligned} p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_N | Z = i)p(Z = i)}{p(\mathbf{x}_1, \dots, \mathbf{x}_N)} \\ &= \frac{\prod_{n=1}^N p(\mathbf{x}_n | Z = i)p(Z = i)}{\sum_{k=1}^3 \prod_{n=1}^N p(\mathbf{x}_n | Z = k)p(Z = k)} \end{aligned} \quad \rightarrow \text{Direct computation}$$

Estimation: Maximum a Posteriori (MAP)

$$\hat{z} = \operatorname{argmax}_i \left[p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) \right]$$

Bayesian Inference: Examples

Inference

Bayes' Theorem:

$$\begin{aligned} p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_N | Z = i)p(Z = i)}{p(\mathbf{x}_1, \dots, \mathbf{x}_N)} \\ &= \frac{\prod_{n=1}^N p(\mathbf{x}_n | Z = i)p(Z = i)}{\sum_{k=1}^3 \prod_{n=1}^N p(\mathbf{x}_n | Z = k)p(Z = k)} \end{aligned} \quad \Rightarrow \text{Direct computation}$$

Estimation: Maximum a Posteriori (MAP)

$$\hat{z} = \operatorname{argmax}_i \left[p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) \right] \Rightarrow \hat{z} = 2, \text{the student house}$$

Bayesian Inference: Examples

Inference

Bayes' Theorem:

$$\begin{aligned} p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_N | Z = i)p(Z = i)}{p(\mathbf{x}_1, \dots, \mathbf{x}_N)} \\ &= \frac{\prod_{n=1}^N p(\mathbf{x}_n | Z = i)p(Z = i)}{\sum_{k=1}^3 \prod_{n=1}^N p(\mathbf{x}_n | Z = k)p(Z = k)} \end{aligned} \quad \rightarrow \text{Direct computation}$$

Estimation: Maximum a Posteriori (MAP)

$$\hat{z} = \operatorname{argmax}_i \left[p(Z = i | \mathbf{x}_1, \dots, \mathbf{x}_N) \right] \Rightarrow \hat{z} = 2, \text{the student house}$$

Decision:



These pranksters will hear from me at the Uni!


OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - **Direct inference**
 - **The Expectation-Maximization algorithm**
 - **Variational Bayes methods**
- Markov chain Monte Carlo


OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - Direct inference
 - **The Expectation-Maximization algorithm**
 - **Variational Bayes methods**
- Markov chain Monte Carlo


Bayesian Inference: Examples



Bayesian Inference: Examples




Bayesian Inference: Examples



Bayesian Inference: Examples

EM algorithm
•••




Bayesian Inference: Examples

Modeling

Bayesian Inference: Examples

Modeling

Observed variables: $\{\boldsymbol{x}_n\}_{n=1}^N \subset \mathbb{R}^2$




Bayesian Inference: Examples


EM algorithm
○○○

Modeling

Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$




Hidden Variables: $\{Z_n\}_{n=1}^N \in \{1, 2, 3\}^N$




Bayesian Inference: Examples

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Graphical Model:



Hidden Variables: $\{Z_n\}_{n=1}^N \in \{1, 2, 3\}^N$




Bayesian Inference: Examples


EM algorithm
•••

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Graphical Model:



Hidden Variables: $\{Z_n\}_{n=1}^N \in \{1, 2, 3\}^N$



Conditional:


$$p(X_n = x_n | Z_n = k; \theta) = \mathcal{N}(x_n; \mu_k, \Sigma_k)$$

Bayesian Inference: Examples


EM algorithm
•••

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Graphical Model:



Hidden Variables: $\{Z_n\}_{n=1}^N \in \{1, 2, 3\}^N$



Conditional:

$$p(X_n = x_n | Z_n = k; \theta) = \mathcal{N}(x_n; \mu_k, \Sigma_k)$$


$$\text{Priors: } p(Z_n = k; \theta) = \pi_k, \quad \sum_{k=1}^K \pi_k = 1$$

Bayesian Inference: Examples


EM algorithm
•••

Modeling


Observed variables: $\{x_n\}_{n=1}^N \subset \mathbb{R}^2$



Graphical Model:



Hidden Variables: $\{Z_n\}_{n=1}^N \in \{1, 2, 3\}^N$



Conditional:

$$p(X_n = x_n | Z_n = k; \theta) = \mathcal{N}(x_n; \mu_k, \Sigma_k)$$

$$\text{Priors: } p(Z_n = k; \theta) = \pi_k, \quad \sum_{k=1}^K \pi_k = 1$$

$$\text{Parameters: } \theta = \{\mu_k, \Sigma_k, \pi_k\}_{k=1}^K$$

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

where $p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta})p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})}$

$$\propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

where $p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta})p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})}$

$$\propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

Simple form, but $\boldsymbol{\theta}$ is unknown

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

where $p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta})p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})}$
 $\propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$

Simple form, but $\boldsymbol{\theta}$ is unknown \Rightarrow Maximum likelihood? $\hat{\boldsymbol{\theta}} = \underset{\boldsymbol{\theta}}{\operatorname{argmax}} \mathcal{L}(\boldsymbol{\theta}; \mathbf{X})$

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

$$\text{where } p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}) p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})} \\ \propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

Simple form, but $\boldsymbol{\theta}$ is unknown \Rightarrow Maximum likelihood? $\hat{\boldsymbol{\theta}} = \underset{\boldsymbol{\theta}}{\operatorname{argmax}} \mathcal{L}(\boldsymbol{\theta}; \mathbf{X})$

$$\mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = p(\mathbf{X}_1 = \mathbf{x}_1, \dots, \mathbf{X}_N = \mathbf{x}_N; \boldsymbol{\theta})$$

$$= \log \left(\sum_{z_1, \dots, z_N=1}^K \prod_{n=1}^N \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right)$$

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

$$\text{where } p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}) p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})} \\ \propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

Simple form, but $\boldsymbol{\theta}$ is unknown \Rightarrow Maximum likelihood? $\hat{\boldsymbol{\theta}} = \underset{\boldsymbol{\theta}}{\operatorname{argmax}} \mathcal{L}(\boldsymbol{\theta}; \mathbf{X})$

$$\mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = p(\mathbf{X}_1 = \mathbf{x}_1, \dots, \mathbf{X}_N = \mathbf{x}_N; \boldsymbol{\theta}) \\ = \log \left(\sum_{z_1, \dots, z_N=1}^K \prod_{n=1}^N \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right) \left\{ \begin{array}{l} \bullet \text{Non-convex} \\ \bullet \text{Combinatorial} \\ \bullet \text{Intractable} \end{array} \right.$$

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

$$\text{where } p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}) p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})} \\ \propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

Simple form, but $\boldsymbol{\theta}$ is unknown \Rightarrow Maximum likelihood? $\hat{\boldsymbol{\theta}} = \underset{\boldsymbol{\theta}}{\operatorname{argmax}} \mathcal{L}(\boldsymbol{\theta}; \mathbf{X})$

$$\mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = p(\mathbf{X}_1 = \mathbf{x}_1, \dots, \mathbf{X}_N = \mathbf{x}_N; \boldsymbol{\theta}) \\ = \log \left(\sum_{z_1, \dots, z_N=1}^K \prod_{n=1}^N \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right) \left\{ \begin{array}{l} \bullet \text{Non-convex} \\ \bullet \text{Combinatorial} \\ \bullet \text{Intractable} \end{array} \right.$$

- The joint probability $p(\mathbf{X}, \mathbf{Z}; \boldsymbol{\theta})$ has a much simpler form than the marginal $p(\mathbf{X}; \boldsymbol{\theta})$
- \mathbf{Z} is a hidden variable, and cannot be estimated without knowing $\boldsymbol{\theta}$

Bayesian Inference: Examples

Inference

Bayes' Theorem: $p(Z_1 = z_1, \dots, Z_N = z_N | \mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}) = \prod_{n=1}^N p(Z_n = z_n | \mathbf{x}_n; \boldsymbol{\theta})$

$$\text{where } p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}) p(Z_n = k; \boldsymbol{\theta})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta})} \\ \propto \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$$

Simple form, but $\boldsymbol{\theta}$ is unknown \Rightarrow Maximum likelihood? $\hat{\boldsymbol{\theta}} = \underset{\boldsymbol{\theta}}{\operatorname{argmax}} \mathcal{L}(\boldsymbol{\theta}; \mathbf{X})$


$$\mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = p(\mathbf{X}_1 = \mathbf{x}_1, \dots, \mathbf{X}_N = \mathbf{x}_N; \boldsymbol{\theta})$$

$$= \log \left(\sum_{z_1, \dots, z_N=1}^K \prod_{n=1}^N \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right) \left\{ \begin{array}{l} \bullet \text{Non-convex} \\ \bullet \text{Combinatorial} \\ \bullet \text{Intractable} \end{array} \right.$$


- The joint probability $p(\mathbf{X}, \mathbf{Z}; \boldsymbol{\theta})$ has a much simpler form than the marginal $p(\mathbf{X}; \boldsymbol{\theta})$
- \mathbf{Z} is a hidden variable, and cannot be estimated without knowing $\boldsymbol{\theta}$

⇒ **Expectation-Maximization (EM) algorithm**


Bayesian Inference: Examples



Bayesian Inference: Examples



Bayesian Inference: Examples




Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

Bayesian Inference: Examples




Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

Bayesian Inference: Examples



Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$


$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$$

where $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$ is the conditional cross entropy of \mathbf{Z} given $\mathbf{X}, \boldsymbol{\theta}$ for the distribution $p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)})$.

Bayesian Inference: Examples



Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$


$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$$

where $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$ is the conditional cross entropy of \mathbf{Z} given $\mathbf{X}, \boldsymbol{\theta}$ for the distribution $p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)})$.

$$\text{In particular: } \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

Bayesian Inference: Examples



Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$


$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$$

where $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$ is the conditional cross entropy of \mathbf{Z} given $\mathbf{X}, \boldsymbol{\theta}$ for the distribution $p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)})$.

$$\text{In particular: } \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

$$\text{Therefore: } \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) - \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

Bayesian Inference: Examples



Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$$


where $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$ is the conditional cross entropy of \mathbf{Z} given $\mathbf{X}, \boldsymbol{\theta}$ for the distribution $p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)})$.

$$\text{In particular: } \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

$$\text{Therefore: } \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) - \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

According to Gibb's inequality, we have $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) \geq H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)})$.

Bayesian Inference: Examples



Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$$

where $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$ is the conditional cross entropy of \mathbf{Z} given $\mathbf{X}, \boldsymbol{\theta}$ for the distribution $p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)})$.


$$\text{In particular: } \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

$$\text{Therefore: } \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) - \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

According to Gibb's inequality, we have $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) \geq H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)})$.

$$\mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) - \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) \geq Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)})$$

Bayesian Inference: Examples



Proof of correctness:

$$p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}) = \frac{p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta})}{p(\mathbf{X}|\boldsymbol{\theta})} \quad (\text{Baye's theorem})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\mathbb{E}_{\mathbf{Z}|\mathbf{X},\boldsymbol{\theta}^{(i)}}[\log p(\mathbf{X}|\boldsymbol{\theta})] = \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{X}, \mathbf{Z}|\boldsymbol{\theta}) - \sum_{\mathbf{Z}} p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)}) \log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta})$$

$$\log p(\mathbf{X}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$$

where $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)})$ is the conditional cross entropy of \mathbf{Z} given $\mathbf{X}, \boldsymbol{\theta}$ for the distribution $p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\theta}^{(i)})$.

$$\text{In particular: } \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

$$\text{Therefore: } \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) - \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) = Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}) + H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)}).$$

According to Gibb's inequality, we have $H(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) \geq H(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)})$.

$$\mathcal{L}(\boldsymbol{\theta}; \mathbf{X}) - \mathcal{L}(\boldsymbol{\theta}^{(i)}; \mathbf{X}) \geq Q(\boldsymbol{\theta}|\boldsymbol{\theta}^{(i)}) - Q(\boldsymbol{\theta}^{(i)}|\boldsymbol{\theta}^{(i)})$$

The likelihood can only increase at each step!

Bayesian Inference: Examples

Derivations for the Gaussian mixture model

Bayesian Inference: Examples

Derivations for the Gaussian mixture model

- **E-step:** computing the current posterior probabilities

$$r_{n,k}^{(i)} = p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}^{(i)}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}^{(i)}) p(Z_n = k; \boldsymbol{\theta}^{(i)})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}^{(i)})} \propto \pi_k^{(i)} \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k^{(i)}, \boldsymbol{\Sigma}_k^{(i)})$$

Bayesian Inference: Examples

Derivations for the Gaussian mixture model

- **E-step:** computing the current posterior probabilities

$$r_{n,k}^{(i)} = p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}^{(i)}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}^{(i)}) p(Z_n = k; \boldsymbol{\theta}^{(i)})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}^{(i)})} \propto \pi_k^{(i)} \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k^{(i)}, \boldsymbol{\Sigma}_k^{(i)})$$

We deduce $Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)})$:

$$\begin{aligned} Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)}) &= \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \{ \log \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}, \mathbf{Z}) \} \\ &= \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \left\{ \log \prod_{n=1}^N \sum_{k=1}^K \mathbb{I}\{Z_n = k\} \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right\} \\ &= \sum_{k,n=1}^{K,N} \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \{ \mathbb{I}\{Z_n = k\} \} \log (\pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)) \\ &= \sum_{k,n=1}^{K,N} r_{n,k}^{(i)} \left(\log \pi_k - \frac{1}{2} \log |\boldsymbol{\Sigma}_k| - \frac{1}{2} (\mathbf{x}_n - \boldsymbol{\mu}_k)^\top \boldsymbol{\Sigma}_k^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_k) \right) + \text{const.} \end{aligned}$$

Bayesian Inference: Examples

Derivations for the Gaussian mixture model

- **E-step:** computing the current posterior probabilities

$$r_{n,k}^{(i)} = p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}^{(i)}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}^{(i)}) p(Z_n = k; \boldsymbol{\theta}^{(i)})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}^{(i)})} \propto \pi_k^{(i)} \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k^{(i)}, \boldsymbol{\Sigma}_k^{(i)})$$

We deduce $Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)})$:

$$\begin{aligned} Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)}) &= \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \{ \log \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}, \mathbf{Z}) \} \\ &= \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \left\{ \log \prod_{n=1}^N \sum_{k=1}^K \mathbb{I}\{Z_n = k\} \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right\} \\ &= \sum_{k,n=1}^{K,N} \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \{ \mathbb{I}\{Z_n = k\} \} \log (\pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)) \\ &= \sum_{k,n=1}^{K,N} r_{n,k}^{(i)} \left(\log \pi_k - \frac{1}{2} \log |\boldsymbol{\Sigma}_k| - \frac{1}{2} (\mathbf{x}_n - \boldsymbol{\mu}_k)^\top \boldsymbol{\Sigma}_k^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_k) \right) + \text{const.} \end{aligned}$$

- **M-step:** maximizing $Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)})$ by finding the zeros of the derivative

Bayesian Inference: Examples

Derivations for the Gaussian mixture model

- **E-step:** computing the current posterior probabilities

$$r_{n,k}^{(i)} = p(Z_n = k | \mathbf{x}_n; \boldsymbol{\theta}^{(i)}) = \frac{p(\mathbf{x}_n | Z_n = k; \boldsymbol{\theta}^{(i)}) p(Z_n = k; \boldsymbol{\theta}^{(i)})}{p(\mathbf{x}_1, \dots, \mathbf{x}_N; \boldsymbol{\theta}^{(i)})} \propto \pi_k^{(i)} \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k^{(i)}, \boldsymbol{\Sigma}_k^{(i)})$$

We deduce $Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)})$:

$$\begin{aligned} Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)}) &= \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \{ \log \mathcal{L}(\boldsymbol{\theta}; \mathbf{X}, \mathbf{Z}) \} \\ &= \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \left\{ \log \prod_{n=1}^N \sum_{k=1}^K \mathbb{I}\{Z_n = k\} \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right\} \\ &= \sum_{k,n=1}^{K,N} \mathbb{E}_{\mathbf{Z} | \mathbf{X}, \boldsymbol{\theta}^{(i)}} \{ \mathbb{I}\{Z_n = k\} \} \log (\pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)) \\ &= \sum_{k,n=1}^{K,N} r_{n,k}^{(i)} \left(\log \pi_k - \frac{1}{2} \log |\boldsymbol{\Sigma}_k| - \frac{1}{2} (\mathbf{x}_n - \boldsymbol{\mu}_k)^\top \boldsymbol{\Sigma}_k^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_k) \right) + \text{const.} \end{aligned}$$

- **M-step:** maximizing $Q(\boldsymbol{\theta} | \boldsymbol{\theta}^{(i)})$ by finding the zeros of the derivative

$$\pi_k^{(i+1)} = \frac{1}{N} \sum_{n=1}^N r_{n,k}^{(i)},$$

$$\boldsymbol{\mu}_k^{(i+1)} = \frac{1}{N} \sum_{n=1}^N r_{n,k}^{(i)} \mathbf{x}_n,$$


$$\boldsymbol{\Sigma}_k^{(i+1)} = \frac{1}{N} \sum_{n=1}^N r_{n,k}^{(i)} (\mathbf{x}_n - \boldsymbol{\mu}_k^{(i+1)}) (\mathbf{x}_n - \boldsymbol{\mu}_k^{(i+1)})^\top$$

Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{Z|X,\theta^{(i)}}[\log \mathcal{L}(\theta; X, Z)]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence




Bayesian Inference: Examples

Inference

EM algorithm

- **Initialization:** Random « guess » for θ
- **E-step:** $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- **M-step:** $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- **Convergence**




Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{Z|X,\theta^{(i)}}[\log \mathcal{L}(\theta; X, Z)]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{Z|X,\theta^{(i)}}[\log \mathcal{L}(\theta; X, Z)]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{Z|X,\theta^{(i)}}[\log \mathcal{L}(\theta; X, Z)]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{Z|X,\theta^{(i)}}[\log \mathcal{L}(\theta; X, Z)]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- M-step: $\theta^{(i+1)} = \underset{\theta}{\operatorname{argmax}} Q(\theta|\theta^{(i)})$
- Convergence





Bayesian Inference: Examples

Inference

EM algorithm

- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{\mathbf{Z}|\mathbf{X},\theta^{(i)}}[\log \mathcal{L}(\theta; \mathbf{X}, \mathbf{Z})]$
- M-step: $\theta^{(i+1)} = \operatorname{argmax}_{\theta} Q(\theta|\theta^{(i)})$
- Convergence




Bayesian Inference: Examples

Inference

EM algorithm


- Initialization: Random « guess » for θ
- E-step: $Q(\theta|\theta^{(i)}) = \mathbb{E}_{Z|X,\theta^{(i)}}[\log \mathcal{L}(\theta; X, Z)]$
- M-step: $\theta^{(i+1)} = \operatorname{argmax}_{\theta} Q(\theta|\theta^{(i)})$
- Convergence



Decision:



Minus 10 points for Mr. Green, minus 5 points for the others!




OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - Direct inference
 - **The Expectation-Maximization algorithm**
 - **Variational Bayes methods**
- Markov chain Monte Carlo

OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - Direct inference
 - The Expectation-Maximization algorithm
 - **Variational Bayes methods**
- Markov chain Monte Carlo

Bayesian Inference: Examples



The next day...

Bayesian Inference: Examples



The next day...

Bayesian Inference: Examples



The next day...

Bayesian Inference: Examples




The next day...

Bayesian Inference: Examples




The next day...

Bayesian Inference: Examples




The next day...

Bayesian Inference: Examples



Bayesian Inference: Examples



Bayesian Inference: Examples

Modeling A « fully Bayesian » model

$$\left. \begin{aligned} p(\boldsymbol{x}_n | Z_n = k, \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) &= \mathcal{N}(\boldsymbol{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) \\ p(Z_n = k | \boldsymbol{\pi}) &= \pi_k \end{aligned} \right\} \text{GMM (same as before)}$$

Bayesian Inference: Examples

Modeling A « fully Bayesian » model

$$\left. \begin{aligned} p(\mathbf{x}_n | Z_n = k, \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) &= \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) \\ p(Z_n = k | \boldsymbol{\pi}) &= \pi_k \end{aligned} \right\} \text{GMM (same as before)}$$

$$\left. \begin{aligned} p(\boldsymbol{\Lambda}_k) &= \mathcal{W}(\boldsymbol{\Lambda}_k; \mathbf{W}_0, \nu_0) \\ p(\boldsymbol{\mu}_k | \boldsymbol{\Lambda}_k) &= \mathcal{N}(\boldsymbol{\mu}_k; \mathbf{m}_0, \boldsymbol{\Lambda}_k^{-1}) \\ p(\boldsymbol{\pi}) &= \text{SymDir}(\boldsymbol{\pi} | \alpha_0) \end{aligned} \right\}$$

Priors on all parameters

Note: These are the *conjugate priors* for the normal and the multinomial distributions, i.e., they are such that $p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k | \mathbf{x}_n, Z_n = k) \cong p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k)$ and $p(\boldsymbol{\pi} | z_n) \cong p(\boldsymbol{\pi})$

Bayesian Inference: Examples

Modeling A « fully Bayesian » model

$$\left. \begin{aligned} p(\mathbf{x}_n | Z_n = k, \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) &= \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) \\ p(Z_n = k | \boldsymbol{\pi}) &= \pi_k \end{aligned} \right\} \text{GMM (same as before)}$$

$$p(\boldsymbol{\Lambda}_k) = \mathcal{W}(\boldsymbol{\Lambda}_k; \mathbf{W}_0, \nu_0)$$


$$p(\boldsymbol{\mu}_k | \boldsymbol{\Lambda}_k) = \mathcal{N}(\boldsymbol{\mu}_k; \mathbf{m}_0, \boldsymbol{\Lambda}_k^{-1})$$

$$p(\boldsymbol{\pi}) = \text{SymDir}(\boldsymbol{\pi} | \alpha_0)$$

Priors on all parameters

Note: These are the *conjugate priors* for the normal and the multinomial distributions, i.e., they are such that $p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k | \mathbf{x}_n, Z_n = k) \cong p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k)$ and $p(\boldsymbol{\pi} | z_n) \cong p(\boldsymbol{\pi})$

Graphical model:



Bayesian Inference: Examples

Modeling A « fully Bayesian » model

$$\left. \begin{aligned} p(\mathbf{x}_n | Z_n = k, \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) &= \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) \\ p(Z_n = k | \boldsymbol{\pi}) &= \pi_k \end{aligned} \right\} \text{GMM (same as before)}$$

$$p(\boldsymbol{\Lambda}_k) = \mathcal{W}(\boldsymbol{\Lambda}_k; \mathbf{W}_0, \nu_0)$$


$$p(\boldsymbol{\mu}_k | \boldsymbol{\Lambda}_k) = \mathcal{N}(\boldsymbol{\mu}_k; \mathbf{m}_0, \boldsymbol{\Lambda}_k^{-1})$$

$$p(\boldsymbol{\pi}) = \text{SymDir}(\boldsymbol{\pi} | \alpha_0)$$

Priors on all parameters

Note: These are the *conjugate priors* for the normal and the multinomial distributions, i.e., they are such that $p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k | \mathbf{x}_n, Z_n = k) \cong p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k)$ and $p(\boldsymbol{\pi} | z_n) \cong p(\boldsymbol{\pi})$

Graphical model:



Choice of hyperparameters:

$$\mathbf{W}_0 = \mathbf{I}$$

$$\nu_0 = D = 2$$

$$\mathbf{m}_0 = \text{mean}(\mathbf{X})$$

$\alpha_0 > 0$: low values will allow Gaussian weights to be close to 0

Bayesian Inference: Examples

Inference

- The posterior distribution $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$ is intractable

Bayesian Inference: Examples

Inference

- The posterior distribution $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$ is intractable
- **Technique:** use a *variational approximation* $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X}) \approx q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$, where $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is restricted to a family of distributions having a simpler form than the true $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$

Bayesian Inference: Examples

Inference

- The posterior distribution $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$ is intractable
- **Technique:** use a *variational approximation* $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X}) \approx q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$, where $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is restricted to a family of distributions having a simpler form than the true $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$
- The variational distribution $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is typically assumed to **factorize** over some partition of the latent variables.

Bayesian Inference: Examples

Inference

- The posterior distribution $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$ is intractable
- **Technique:** use a *variational approximation* $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X}) \approx q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$, where $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is restricted to a family of distributions having a simpler form than the true $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$
- The variational distribution $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is typically assumed to **factorize** over some partition of the latent variables.
- Here we use: $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = q_{\mathbf{Z}}(\mathbf{Z})q_{\boldsymbol{\Lambda}\boldsymbol{\mu}\boldsymbol{\pi}}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$. Remarkably, this is the **only** assumption needed to obtain a tractable *EM-like* inference procedure.

Bayesian Inference: Examples

Inference

- The posterior distribution $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$ is intractable
- **Technique:** use a *variational approximation* $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X}) \approx q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$, where $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is restricted to a family of distributions having a simpler form than the true $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$
- The variational distribution $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is typically assumed to **factorize** over some partition of the latent variables.
- Here we use: $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = q_{\mathbf{Z}}(\mathbf{Z})q_{\boldsymbol{\Lambda}\boldsymbol{\mu}\boldsymbol{\pi}}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$. Remarkably, this is the **only** assumption needed to obtain a tractable *EM-like* inference procedure.
- Such procedures are referred to as **Variational Bayesian EM algorithms**.

Bayesian Inference: Examples

Inference

- The posterior distribution $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$ is intractable
- **Technique:** use a *variational approximation* $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X}) \approx q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$, where $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is restricted to a family of distributions having a simpler form than the true $p(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi} | \mathbf{X})$
- The variational distribution $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$ is typically assumed to **factorize** over some partition of the latent variables.
- Here we use: $q(\mathbf{Z}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = q_{\mathbf{Z}}(\mathbf{Z})q_{\boldsymbol{\Lambda}\boldsymbol{\mu}\boldsymbol{\pi}}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})$. Remarkably, this is the **only** assumption needed to obtain a tractable *EM-like* inference procedure.
- Such procedures are referred to as ***Variational Bayesian EM algorithms***.
For $p(\mathbf{Z}, \mathbf{W} | \mathbf{X}) \approx q_{\mathbf{Z}}(\mathbf{Z})q_{\mathbf{W}}(\mathbf{W})$:

VB-EM {

- **E-Z step:** $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_W^{(i-1)}(\mathbf{W})} \{ \log p(\mathbf{Z} | \mathbf{X}, \mathbf{W}) \} \right)$
- **E-W step:** $q_W^{(i)}(\mathbf{W}) \propto \exp \left(\mathbb{E}_{q_Z^{(i)}(\mathbf{Z})} \{ \log p(\mathbf{W} | \mathbf{X}, \mathbf{Z}) \} \right)$

Bayesian Inference: Examples

Inference

Proof of correctness

- Using a similar reasonning as for EM, we can show that the VB-EM iteratively minimizes the Kulback-Leibler divergence between the true posterior $p(\mathbf{Z}, \mathbf{W}|\mathbf{X})$ and its variational approximation $q(\mathbf{Z}, \mathbf{W}) = q_Z(\mathbf{Z})q_W(\mathbf{W})$:

$$(q_Z^{(\infty)}, q_W^{(\infty)}) = \operatorname{argmin}_{q_Z, q_W} \text{KL}(q||p)$$

- VB-EM** {
- **E-Z step:** $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_W^{(i-1)}(\mathbf{W})} \{ \log p(\mathbf{Z}|\mathbf{X}, \mathbf{W}) \} \right)$
 - **E-W step:** $q_W^{(i)}(\mathbf{W}) \propto \exp \left(\mathbb{E}_{q_Z^{(i)}(\mathbf{Z})} \{ \log p(\mathbf{W}|\mathbf{X}, \mathbf{Z}) \} \right)$

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

• **E-Λμπ-Step:**

$$\log q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = \log p(\boldsymbol{\pi}) + \sum_{k=1}^K \log p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) + \sum_{n,k=1}^{N,K} q_{Z_n}^{(i-1)}(k) \log \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) + \text{const.}$$

Using the decomposition $q_Z^{(i-1)}(\mathbf{Z}) = \prod_{n=1}^N q_{Z_n}^{(i-1)}(k)$ (see E-Z-step).

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

- **E-Λμπ-Step:**

$$\log q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = \log p(\boldsymbol{\pi}) + \sum_{k=1}^K \log p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) + \sum_{n,k=1}^{N,K} q_{Z_n}^{(i-1)}(k) \log \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) + \text{const.}$$

Using the decomposition $q_Z^{(i-1)}(\mathbf{Z}) = \prod_{n=1}^N q_{Z_n}^{(i-1)}(k)$ (see E-Z-step).

This leads to the factorization: $q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = q_{\pi}^{(i)}(\boldsymbol{\pi}) \prod_{k=1}^K q_{\Lambda_k \mu_k}^{(i)}(\boldsymbol{\Lambda}_k, \boldsymbol{\mu}_k)$

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

- **E-Λμπ-Step:**

$$\log q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = \log p(\boldsymbol{\pi}) + \sum_{k=1}^K \log p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) + \sum_{n,k=1}^{N,K} q_{Z_n}^{(i-1)}(k) \log \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) + \text{const.}$$

Using the decomposition $q_Z^{(i-1)}(\mathbf{Z}) = \prod_{n=1}^N q_{Z_n}^{(i-1)}(k)$ (see E-Z-step).

This leads to the factorization: $q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = q_{\pi}^{(i)}(\boldsymbol{\pi}) \prod_{k=1}^K q_{\Lambda_k \mu_k}^{(i)}(\boldsymbol{\Lambda}_k, \boldsymbol{\mu}_k)$

with $\begin{cases} q_{\pi}^{(i)}(\boldsymbol{\pi}) = \text{Dir}(\boldsymbol{\pi}; \alpha_0 + N_1^{(i)}, \dots, \alpha_0 + N_K^{(i)}) \\ q_{\Lambda_k \mu_k}^{(i)}(\boldsymbol{\Lambda}_k, \boldsymbol{\mu}_k) = \mathcal{N}\left(\boldsymbol{\mu}_k; \mathbf{m}_k^{(i)}, \frac{\boldsymbol{\Lambda}_k^{-1}}{1 + N_k^{(i)}}\right) \mathcal{W}\left(\boldsymbol{\Lambda}_k; \mathbf{W}_k^{(i)}, \nu_k^{(i)}\right), \end{cases}$

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

- **E-Λμπ-Step:**

$$\log q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = \log p(\boldsymbol{\pi}) + \sum_{k=1}^K \log p(\boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k) + \sum_{n,k=1}^{N,K} q_{Z_n}^{(i-1)}(k) \log \pi_k \mathcal{N}(\mathbf{x}_n; \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1}) + \text{const.}$$

Using the decomposition $q_Z^{(i-1)}(\mathbf{Z}) = \prod_{n=1}^N q_{Z_n}^{(i-1)}(k)$ (see E-Z-step).

This leads to the factorization: $q_{\Lambda \mu \pi}^{(i)}(\boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi}) = q_{\pi}^{(i)}(\boldsymbol{\pi}) \prod_{k=1}^K q_{\Lambda_k \mu_k}^{(i)}(\boldsymbol{\Lambda}_k, \boldsymbol{\mu}_k)$

with $\begin{cases} q_{\pi}^{(i)}(\boldsymbol{\pi}) = \text{Dir}(\boldsymbol{\pi}; \alpha_0 + N_1^{(i)}, \dots, \alpha_0 + N_K^{(i)}) \\ q_{\Lambda_k \mu_k}^{(i)}(\boldsymbol{\Lambda}_k, \boldsymbol{\mu}_k) = \mathcal{N}\left(\boldsymbol{\mu}_k; \mathbf{m}_k^{(i)}, \frac{\boldsymbol{\Lambda}_k^{-1}}{1 + N_k^{(i)}}\right) \mathcal{W}\left(\boldsymbol{\Lambda}_k; \mathbf{W}_k^{(i)}, \nu_k^{(i)}\right), \end{cases}$

where: $\bar{\mathbf{x}}_k^{(i)} = \frac{1}{N_k^{(i)}} \sum_{n=1}^N q_{Z_n}^{(i-1)}(k) \mathbf{x}_n, \quad \mathbf{S}_k^{(i)} = \frac{1}{N_k^{(i)}} \sum_{n=1}^N q_{Z_n}^{(i-1)}(k) (\mathbf{x}_n - \bar{\mathbf{x}}_k^{(i)}) (\mathbf{x}_n - \bar{\mathbf{x}}_k^{(i)})^\top,$

$$N_k^{(i)} = \sum_{n=1}^N q_{Z_n}^{(i-1)}(k), \quad \mathbf{m}_k^{(i)} = \frac{\mathbf{m}_0 + N_k^{(i)} \bar{\mathbf{x}}_k^{(i)}}{1 + N_k^{(i)}},$$

$$\mathbf{W}_k^{(i)-1} = \mathbf{W}_0^{-1} + N_k^{(i)} \mathbf{S}_k^{(i)} + \frac{N_k^{(i)}}{N_k^{(i)} + 1} (\bar{\mathbf{x}}_k^{(i)} - \mathbf{m}_0) (\bar{\mathbf{x}}_k^{(i)} - \mathbf{m}_0)^\top, \quad \nu_k^{(i)} = \nu_0 + N_k^{(i)}$$

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

- E-Z-Step:

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

•E-Z-Step:

$$\log q_Z^{(i)}(\mathbf{Z}) = \mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{\log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})\} + \text{const.}$$

$$= \sum_{n,k=1}^{N,K} \mathbb{I}\{Z_n = k\} \log \rho_{n,k}^{(i)} + \text{const.}$$

where $\rho_{n,k}^{(i)} = \mathbb{E}_{q_{\pi}^{(i)}} \{\log \pi_k\} + \frac{1}{2} \mathbb{E}_{q_{\Lambda_k}^{(i)}} \{\log |\boldsymbol{\Lambda}_k|\} - \frac{D}{2} \log 2\pi - \frac{1}{2} \mathbb{E}_{q_{\Lambda_k \mu_k}^{(i)}} \{(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top \boldsymbol{\Lambda}_k (\mathbf{x}_n - \boldsymbol{\mu}_k)\}$

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

•E-Z-Step:

$$\log q_Z^{(i)}(\mathbf{Z}) = \mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{\log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})\} + \text{const.}$$

$$= \sum_{n,k=1}^{N,K} \mathbb{I}\{Z_n = k\} \log \rho_{n,k}^{(i)} + \text{const.}$$

where $\rho_{n,k}^{(i)} = \mathbb{E}_{q_\pi^{(i)}} \{\log \pi_k\} + \frac{1}{2} \mathbb{E}_{q_{\Lambda_k}^{(i)}} \{\log |\boldsymbol{\Lambda}_k|\} - \frac{D}{2} \log 2\pi - \frac{1}{2} \mathbb{E}_{q_{\Lambda_k \mu_k}^{(i)}} \{(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top \boldsymbol{\Lambda}_k (\mathbf{x}_n - \boldsymbol{\mu}_k)\}$

It follows that $q_Z^{(i)}(\mathbf{Z}) = \prod_{n=1}^N q_{Z_n}^{(i)}(k)$ where $q_{Z_n}^{(i)}(k) = \frac{\rho_{n,k}^{(i)}}{\sum_{j=1}^K \rho_{n,j}^{(i)}} = r_{n,k}^{(i)}$

Bayesian Inference: Examples

Inference

Derivations for the Bayesian mixture of Gaussian

•E-Z-Step:

$$\log q_Z^{(i)}(\mathbf{Z}) = \mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{\log p(\mathbf{Z}|\mathbf{X}, \boldsymbol{\Lambda}, \boldsymbol{\mu}, \boldsymbol{\pi})\} + \text{const.}$$

$$= \sum_{n,k=1}^{N,K} \mathbb{I}\{Z_n = k\} \log \rho_{n,k}^{(i)} + \text{const.}$$

where $\rho_{n,k}^{(i)} = \mathbb{E}_{q_\pi^{(i)}} \{\log \pi_k\} + \frac{1}{2} \mathbb{E}_{q_{\Lambda_k}^{(i)}} \{\log |\boldsymbol{\Lambda}_k|\} - \frac{D}{2} \log 2\pi - \frac{1}{2} \mathbb{E}_{q_{\Lambda_k \mu_k}^{(i)}} \{(\mathbf{x}_n - \boldsymbol{\mu}_k)^\top \boldsymbol{\Lambda}_k (\mathbf{x}_n - \boldsymbol{\mu}_k)\}$

It follows that $q_Z^{(i)}(\mathbf{Z}) = \prod_{n=1}^N q_{Z_n}^{(i)}(k)$ where $q_{Z_n}^{(i)}(k) = \frac{\rho_{n,k}^{(i)}}{\sum_{j=1}^K \rho_{n,j}^{(i)}} = r_{n,k}^{(i)}$

Finally, we can express $r_{n,k}^{(i)}$ as a function of the parameters calculated in previous step:

$$r_{n,k}^{(i)} \propto |\mathbf{W}_k^{(i)}|^{1/2} \exp \left(\psi(\alpha_0 + N_k^{(i)}) + \sum_{i=1}^D \psi \left(\frac{\nu_k^{(i)} + 1 - i}{2} \right) - \frac{D}{2N_k^{(i)} + 2} - \frac{\nu_k^{(i)}}{2} (\mathbf{x}_n - \mathbf{m}_k^{(i)})^\top \mathbf{W}_k^{(i)} (\mathbf{x}_n - \mathbf{m}_k^{(i)}) \right)$$


where $\psi(\cdot)$ denotes the digamma function.

Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- **Initialization:** Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step :** $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step:** $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- **Convergence**




Bayesian Inference: Examples

Inference

VB-EM in action

- **Initialization:** Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step :** $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step:** $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- **Convergence**




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- **E- $\Lambda\mu\pi$ -Step** : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- **E-Z-Step**: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence




Bayesian Inference: Examples

Inference

VB-EM in action

- Initialization: Random means + GMM E-step for $q_Z^{(0)}(\mathbf{Z})$
- E- $\Lambda\mu\pi$ -Step : $q_{\Lambda\mu\pi}^{(i)}(\Lambda, \mu, \pi) \propto \exp \left(\mathbb{E}_{q_Z^{(i-1)}} \{ \log p(\Lambda, \mu, \pi | \mathbf{X}, \mathbf{Z}) \} \right)$
- E-Z-Step: $q_Z^{(i)}(\mathbf{Z}) \propto \exp \left(\mathbb{E}_{q_{\Lambda\mu\pi}^{(i)}} \{ \log p(\mathbf{Z} | \mathbf{X}, \Lambda, \mu, \pi) \} \right)$
- Convergence



Conclusions on GMM VB-EM

- Similar computational time as GMM-EM (though slightly more iterations)
- Priors on Gaussian weights handle automatically degenerate or unused clusters
- Determination of K
- Works even for very small data samples

OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- **Bayesian inference by examples**
 - Direct inference
 - The Expectation-Maximization algorithm
 - **Variational Bayes methods**
- Markov chain Monte Carlo

OUTLINE

- What is Bayesian inference?
 - Overview
 - Classical vs. Bayesian approach
 - Bayes Theorem & Example
 - General Methodology
- Bayesian inference by examples
 - Direct inference
 - The Expectation-Maximization algorithm
 - Variational Bayes methods
- **Markov chain Monte Carlo**