

Partial Differential Equations

Chapter VII - Finite Difference Method

John Cagnol, Pauline Lafitte

The Engineering Program of CentraleSupélec

Lecture 9 – March 3rd 2020

VII.1. Introduction

Consider a well-posed PDE.

Consider a well-posed PDE.

For instance, let $f \in C^0(\Omega)$ and

$$\begin{cases} -\Delta u = f & \text{on } \Omega, \\ u|_{\partial\Omega} = 0. \end{cases}$$

Consider a well-posed PDE.

For instance, let $f \in C^0([0, 1])$ and

$$\begin{cases} -u'' = f & \text{on }]0, 1[, \\ u(0) = 0, \quad u(1) = 0 \end{cases}$$

Consider a well-posed PDE.

For instance, let $f \in C^0([0, 1])$, $c \in C^0([0, 1], \mathbb{R}^+)$ and

$$\begin{cases} -u'' + c u = f & \text{on }]0, 1[, \\ u(0) = 0, \quad u(1) = 0 \end{cases}$$

Consider a well-posed PDE.

For instance, let $f \in C^0([0, 1])$, $c \in C^0([0, 1], \mathbb{R}^+)$ and

$$\begin{cases} -u'' + c u = f & \text{on }]0, 1[, \\ u(0) = 0, \quad u(1) = 0 \end{cases}$$

The core idea underlying the FEM is:

- To mesh the domain and build a finite-dimensional subspace of elementary functions.
- To project u on that subspace of functions.
- To end up with a finite number of unknowns and to solve a linear system.

Consider a well-posed PDE.

For instance, let $f \in C^0([0, 1])$, $c \in C^0([0, 1], \mathbb{R}^+)$ and

$$\begin{cases} -u'' + c u = f & \text{on }]0, 1[, \\ u(0) = 0, \quad u(1) = 0 \end{cases}$$

The core idea underlying the FEM is:

- To mesh the domain and build a finite-dimensional subspace of elementary functions.
- To project u on that subspace of functions.
- To end up with a finite number of unknowns and to solve a linear system.

There is another way...

As you know...

As you know...

A derivative can be approximated by the Newton quotient.

$$f'(x) = \frac{f(x+h) - f(x)}{h} + o(1)$$

(provided f is differentiable at x)

As you know...

A derivative can be approximated by the Newton quotient.

$$f'(x) = \frac{f(x+h) - f(x)}{h} + O(h)$$

(provided f is twice differentiable at x)

As you know...

A derivative can be approximated by the Newton quotient.

$$f'(x) = \frac{f(x+h) - f(x)}{h} + O(h)$$

(provided f is twice differentiable at x)

Higher order derivatives can be approximated by iteration of this process.

As you know...

A derivative can be approximated by the Newton quotient.

$$f'(x) = \frac{f(x+h) - f(x)}{h} + O(h)$$

(provided f is twice differentiable at x)

Higher order derivatives can be approximated by iteration of this process.

The core idea underlying the FDM is:

- To mesh the domain thus obtaining a grid
- To approximate the derivatives of u using the value of u on the grid
- To have a finite numbers of unknowns (the values of u on the grid) and to solve a linear system.

VII.2. FDM in dimension $d = 1$

VII.2.1. Approximating the derivative

Forward, backward and central differences

Let $b > a$, and f be a real-valued function defined on $[a, b]$.

Let $J \in \mathbb{N}^*$ and

$$(x_j)_{j \in \{0, \dots, J+1\}}$$

be a mesh of $[a, b]$ where $x_0 = a$ and $x_{J+1} = b$.

Note $h = (b - a)/(J + 1)$

Forward, backward and central differences

Let $b > a$, and f be a real-valued function defined on $[a, b]$.

Let $J \in \mathbb{N}^*$ and

$$(x_j)_{j \in \{0, \dots, J+1\}}$$

be a mesh of $[a, b]$ where $x_0 = a$ and $x_{J+1} = b$.

Note $h = (b - a)/(J + 1)$

Definition VII.2.1

*The **forward difference** at $x = x_j$ is defined by $f(x + h) - f(x)$*

Forward, backward and central differences

Let $b > a$, and f be a real-valued function defined on $[a, b]$.

Let $J \in \mathbb{N}^*$ and

$$(x_j)_{j \in \{0, \dots, J+1\}}$$

be a mesh of $[a, b]$ where $x_0 = a$ and $x_{J+1} = b$.

Note $h = (b - a)/(J + 1)$

Definition VII.2.1

*The **forward difference** at $x = x_j$ is defined by $f(x_{j+1}) - f(x_j)$*

Forward, backward and central differences

Let $b > a$, and f be a real-valued function defined on $[a, b]$.

Let $J \in \mathbb{N}^*$ and

$$(x_j)_{j \in \{0, \dots, J+1\}}$$

be a mesh of $[a, b]$ where $x_0 = a$ and $x_{J+1} = b$.

Note $h = (b - a)/(J + 1)$

Definition VII.2.1

*The **forward difference** at $x = x_j$ is defined by $f(x_{j+1}) - f(x_j)$*

Definition VII.2.2

*The **backward difference** at $x = x_j$ is defined by $f(x) - f(x - h)$*

Forward, backward and central differences

Let $b > a$, and f be a real-valued function defined on $[a, b]$.

Let $J \in \mathbb{N}^*$ and

$$(x_j)_{j \in \{0, \dots, J+1\}}$$

be a mesh of $[a, b]$ where $x_0 = a$ and $x_{J+1} = b$.

Note $h = (b - a)/(J + 1)$

Definition VII.2.1

*The **forward difference** at $x = x_j$ is defined by $f(x_{j+1}) - f(x_j)$*

Definition VII.2.2

*The **backward difference** at $x = x_j$ is defined by $f(x_j) - f(x_{j-1})$*

Forward, backward and central differences

Let $b > a$, and f be a real-valued function defined on $[a, b]$.

Let $J \in \mathbb{N}^*$ and

$$(x_j)_{j \in \{0, \dots, J+1\}}$$

be a mesh of $[a, b]$ where $x_0 = a$ and $x_{J+1} = b$.

Note $h = (b - a)/(J + 1)$

Definition VII.2.1

*The **forward difference** at $x = x_j$ is defined by $f(x_{j+1}) - f(x_j)$*

Definition VII.2.2

*The **backward difference** at $x = x_j$ is defined by $f(x_j) - f(x_{j-1})$*

Definition VII.2.3

*The **central difference** at $x = x_j$ is defined by $f(x + h/2) - f(x - h/2)$*

Forward, backward and central differences

Remark VII.2.4

The points $x + \frac{h}{2}$ and $x - \frac{h}{2}$ are not the grid.

Forward, backward and central differences

Remark VII.2.4

The points $x + \frac{h}{2}$ and $x - \frac{h}{2}$ are not the grid.

The problem is often remedied by taking the average on the two points nearby on the grid:

- *we replace $f(x + \frac{h}{2})$ by $(f(x + h) + f(x))/2$*
- *we replace $f(x - \frac{h}{2})$ by $(f(x - h) + f(x))/2$*

Forward, backward and central differences

Remark VII.2.4

The points $x + \frac{h}{2}$ and $x - \frac{h}{2}$ are not the grid.

The problem is often remedied by taking the average on the two points nearby on the grid:

- *we replace $f(x + \frac{h}{2})$ by $(f(x_{j+1}) + f(x_j))/2$*
- *we replace $f(x - \frac{h}{2})$ by $(f(x_{j-1}) + f(x_j))/2$*

Forward, backward and central differences

Remark VII.2.4

The points $x + \frac{h}{2}$ and $x - \frac{h}{2}$ are not the grid.

The problem is often remedied by taking the average on the two points nearby on the grid:

- *we replace $f(x + \frac{h}{2})$ by $(f(x_{j+1}) + f(x_j))/2$*
- *we replace $f(x - \frac{h}{2})$ by $(f(x_{j-1}) + f(x_j))/2$*

This leads to

$$f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right) \text{ replaced by } \frac{f(x_{j+1}) + f(x_j)}{2} - \frac{f(x_{j-1}) + f(x_j)}{2}$$

Forward, backward and central differences

Remark VII.2.4

The points $x + \frac{h}{2}$ and $x - \frac{h}{2}$ are not the grid.

The problem is often remedied by taking the average on the two points nearby on the grid:

- *we replace $f(x + \frac{h}{2})$ by $(f(x_{j+1}) + f(x_j))/2$*
- *we replace $f(x - \frac{h}{2})$ by $(f(x_{j-1}) + f(x_j))/2$*

This leads to

$$f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right) \text{ replaced by } \frac{f(x_{j+1}) - f(x_{j-1}))}{2}$$

Forward, backward and central differences

Remark VII.2.4

The points $x + \frac{h}{2}$ and $x - \frac{h}{2}$ are not the grid.

The problem is often remedied by taking the average on the two points nearby on the grid:

- *we replace $f(x + \frac{h}{2})$ by $(f(x_{j+1}) + f(x_j))/2$*
- *we replace $f(x - \frac{h}{2})$ by $(f(x_{j-1}) + f(x_j))/2$*

This leads to

$$f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right) \text{ replaced by } \frac{f(x_{j+1}) - f(x_{j-1}))}{2}$$

*Subsequently, the **central difference** at $x = x_j$ is often defined by*

$$\frac{f(x_{j+1}) - f(x_{j-1}))}{2}$$

Forward, backward and central difference quotient

Definition VII.2.5

The **forward difference quotient** is given by

$$\frac{f(x+h) - f(x)}{h}$$

Forward, backward and central difference quotient

Definition VII.2.5

The **forward difference quotient** is given by

$$\frac{f(x+h) - f(x)}{h} = \frac{f(x_{j+1}) - f(x_j)}{x_{j+1} - x_j}$$

Forward, backward and central difference quotient

Definition VII.2.5

The **forward difference quotient** is given by

$$\frac{f(x+h) - f(x)}{h} = \frac{f(x_{j+1}) - f(x_j)}{x_{j+1} - x_j}$$

Definition VII.2.6

The **backward difference quotient** is given by

$$\frac{f(x) - f(x-h)}{h}$$

Forward, backward and central difference quotient

Definition VII.2.5

The **forward difference quotient** is given by

$$\frac{f(x+h) - f(x)}{h} = \frac{f(x_{j+1}) - f(x_j)}{x_{j+1} - x_j}$$

Definition VII.2.6

The **backward difference quotient** is given by

$$\frac{f(x) - f(x-h)}{h} = \frac{f(x_j) - f(x_{j-1})}{x_j - x_{j-1}}$$

Forward, backward and central difference quotient

Definition VII.2.5

The **forward difference quotient** is given by

$$\frac{f(x+h) - f(x)}{h} = \frac{f(x_{j+1}) - f(x_j)}{x_{j+1} - x_j}$$

Definition VII.2.6

The **backward difference quotient** is given by

$$\frac{f(x) - f(x-h)}{h} = \frac{f(x_j) - f(x_{j-1})}{x_j - x_{j-1}}$$

Definition VII.2.7

The **central difference quotient** is given by

$$\frac{f(x + \frac{h}{2}) - f(x - \frac{h}{2})}{h}$$

Forward, backward and central difference quotient

Definition VII.2.5

The **forward difference quotient** is given by

$$\frac{f(x+h) - f(x)}{h} = \frac{f(x_{j+1}) - f(x_j)}{x_{j+1} - x_j}$$

Definition VII.2.6

The **backward difference quotient** is given by

$$\frac{f(x) - f(x-h)}{h} = \frac{f(x_j) - f(x_{j-1})}{x_j - x_{j-1}}$$

Definition VII.2.7

The **central difference quotient** is given by

$$\frac{f(x + \frac{h}{2}) - f(x - \frac{h}{2})}{h} \quad \text{replaced by} \quad \frac{f(x_{j+1}) - f(x_{j-1}))}{x_{j+1} - x_{j-1}}$$

Relation between difference quotients and derivatives

Provided f is twice differentiable, the Taylor expansion gives

$$f(x + h) = f(x) + h f'(x) + O(h^2)$$

Relation between difference quotients and derivatives

Provided f is twice differentiable, the Taylor expansion gives

$$f(x + h) = f(x) + hf'(x) + O(h^2)$$

Therefore

$$\frac{f(x + h) - f(x)}{h} - f'(x) = O(h)$$

Relation between difference quotients and derivatives

Provided f is twice differentiable, the Taylor expansion gives

$$f(x + h) = f(x) + hf'(x) + O(h^2)$$

Therefore

$$\frac{f(x + h) - f(x)}{h} - f'(x) = O(h) \xrightarrow{h \rightarrow 0} 0$$

Relation between difference quotients and derivatives

Provided f is twice differentiable, the Taylor expansion gives

$$f(x + h) = f(x) + hf'(x) + O(h^2)$$

Therefore

$$\frac{f(x + h) - f(x)}{h} - f'(x) = O(h) \xrightarrow{h \rightarrow 0} 0$$

Definition VII.2.8

*The **local truncation error** or **discretization error** is the difference between the difference quotient and the derivative. If the local truncation error is $O(h^k)$, the method is of **order k**.*

Relation between difference quotients and derivatives

Provided f is twice differentiable, the Taylor expansion gives

$$f(x + h) = f(x) + hf'(x) + O(h^2)$$

Therefore

$$\frac{f(x + h) - f(x)}{h} - f'(x) = O(h) \xrightarrow{h \rightarrow 0} 0$$

Definition VII.2.8

*The **local truncation error** or **discretization error** is the difference between the difference quotient and the derivative. If the local truncation error is $O(h^k)$, the method is of **order k**.*

Proposition VII.2.9

The forward difference quotient approximates the derivative. The method is of order 1.

Relation between difference quotients and derivatives

Provided f is twice differentiable, the Taylor expansion gives

$$f(x - h) = f(x) - hf'(x) + O(h^2)$$

Therefore

$$\frac{f(x) - f(x - h)}{h} - f'(x) = O(h) \xrightarrow{h \rightarrow 0} 0$$

Proposition VII.2.10

*The backward difference quotient approximates the derivative.
The method is of order 1.*

Relation between different quotients and derivatives

Provided f is three times differentiable, the Taylor expansion gives

$$f\left(x + \frac{h}{2}\right) = f(x) + \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

$$f\left(x - \frac{h}{2}\right) = f(x) - \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

Relation between different quotients and derivatives

Provided f is three times differentiable, the Taylor expansion gives

$$f\left(x + \frac{h}{2}\right) = f(x) + \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

$$f\left(x - \frac{h}{2}\right) = f(x) - \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

Therefore

$$f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right) = h f'(x) + O(h^3)$$

Relation between different quotients and derivatives

Provided f is three times differentiable, the Taylor expansion gives

$$f\left(x + \frac{h}{2}\right) = f(x) + \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

$$f\left(x - \frac{h}{2}\right) = f(x) - \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

Therefore

$$\frac{f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right)}{h} - f'(x) = O(h^2)$$

Relation between different quotients and derivatives

Provided f is three times differentiable, the Taylor expansion gives

$$f\left(x + \frac{h}{2}\right) = f(x) + \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

$$f\left(x - \frac{h}{2}\right) = f(x) - \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

Therefore

$$\frac{f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right)}{h} - f'(x) = O(h^2) \xrightarrow{h \rightarrow 0} 0$$

Relation between different quotients and derivatives

Provided f is three times differentiable, the Taylor expansion gives

$$f\left(x + \frac{h}{2}\right) = f(x) + \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

$$f\left(x - \frac{h}{2}\right) = f(x) - \frac{h}{2} f'(x) + \frac{1}{2} \left(\frac{h}{2}\right)^2 f''(x) + O(h^3)$$

Therefore

$$\frac{f\left(x + \frac{h}{2}\right) - f\left(x - \frac{h}{2}\right)}{h} - f'(x) = O(h^2) \xrightarrow{h \rightarrow 0} 0$$

Proposition VII.2.11

The central difference quotient approximates the derivative.

The method is of order 2.

Relation between different quotients and derivatives

Question:

Is there a way to approximate the derivative with a method of order 4?

Relation between different quotients and derivatives

Question:

Is there a way to approximate the derivative with a method of order 4?

Answer:

Will be given in Exercise Q.V.1 during the next lab session.

Stencils

Definition VII.2.12


The **stencil** of a numerical approximation method is the geometric arrangement of the nodes that relate to the point of interest.

In dimension 1, the stencil is “flat”.

We can have more complex arrangements in dimension $d > 1$.

Example

Stencils for the forward, backward and central difference quotients.



Remarks regarding the notations

Remark VII.2.13

Difference quotients *are also called*

- **Newton quotient**
- **Fermat difference quotient**
- **Finite difference approximation of the derivative**

Remarks regarding the notations

Remark VII.2.13

Difference quotients *are also called*

- **Newton quotient**
- **Fermat difference quotient**
- **Finite difference approximation of the derivative**

Remark VII.2.14

*In some books, forward/backward/central **finite difference approximation of the derivative** is shortened by forward/backward/central **difference**.*

This can be confusing.

In case of doubt, make sure you look at the definitions.

Remarks regarding the notations

Remark VII.2.15

In some books,

- *Forward finite differences are denoted Δ*
- *Backward finite differences are denoted ∇*
- *Central finite differences are denoted δ .*

*We will **not** use these notations in this class to avoid confusion with the Laplace and Del operators.*

VII.2.2. Approximating higher-order derivatives

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{f'(x + \frac{h}{2}) - f'(x - \frac{h}{2})}{h} + O(h^2)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{f'(x + \frac{h}{2}) - f'(x - \frac{h}{2})}{h} + O(h^2)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{\frac{f(x + \frac{h}{2} + \frac{h}{2}) - f(x + \frac{h}{2} - \frac{h}{2})}{h} - \frac{f(x - \frac{h}{2} + \frac{h}{2}) - f(x - \frac{h}{2} - \frac{h}{2})}{h}}{h} + O(h^2) + O(h^2)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{\frac{f(x+\frac{h}{2}+\frac{h}{2})-f(x+\frac{h}{2}-\frac{h}{2})}{h} - \frac{f(x-\frac{h}{2}+\frac{h}{2})-f(x-\frac{h}{2}-\frac{h}{2})}{h}}{h} + O(h^2) + O(h^2)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{\frac{f(x+h)-f(x)}{h} - \frac{f(x)-f(x-h)}{h}}{h} + O(h^2)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2} + O(h) + O(h^2)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2} + O(h)$$

(second-order central)

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2} + O(h)$$

(second-order central)

We can do better than $O(h)$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times, the Taylor expansion gives

$$f(x+h) = f(x) + hf'(x) + \frac{1}{2}h^2 f''(x) + \frac{1}{6}h^3 f'''(x) + O(h^4)$$

$$f(x-h) = f(x) - hf'(x) + \frac{1}{2}h^2 f''(x) - \frac{1}{6}h^3 f'''(x) + O(h^4)$$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times, the Taylor expansion gives

$$f(x+h) = f(x) + hf'(x) + \frac{1}{2}h^2 f''(x) + \frac{1}{6}h^3 f'''(x) + O(h^4)$$

$$f(x-h) = f(x) - hf'(x) + \frac{1}{2}h^2 f''(x) - \frac{1}{6}h^3 f'''(x) + O(h^4)$$

$$f(x+h) + f(x-h) = 2f(x) + h^2 f''(x) + O(h^4)$$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f(x+h) + f(x-h) = 2f(x) + h^2 f''(x) + O(h^4)$$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$f(x+h) - 2f(x) + f(x-h) = h^2 f''(x) + O(h^4)$$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$\frac{f(x+h) - 2f(x) + f(x-h)}{h^2} = f''(x) + O(h^2)$$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

$$\frac{f(x+h) - 2f(x) + f(x-h)}{h^2} - f''(x) = O(h^2)$$

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

Proposition VII.2.16

The second-order central difference quotient

$$\frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$

approximates the second derivative. The method is of order 2.

One can obtain finite difference approximations to higher order derivatives by applying the finite difference approximation to f' .

For example, consider f differentiable four times,

Proposition VII.2.16

The second-order central difference quotient

$$\frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$

approximates the second derivative. The method is of order 2.

The stencil is: 

More generally,

Proposition VII.2.17

The n th-order central difference quotient

$$\frac{1}{h^n} \sum_{i=0}^n (-1)^i \binom{n}{i} f(x + (n/2 - i)h)$$

approximates the n -th derivative. The method is of order 2.

Remark VII.2.18

For odd n , the function is not evaluated on the grid (as for $n = 1$). The problem may be remedied taking the average on the two nearest points of the grid.

Similarly,

Proposition VII.2.19

The n th-order forward difference quotient

$$\frac{1}{h^n} \sum_{i=0}^n (-1)^i \binom{n}{i} f(x + (n-i)h) = \frac{1}{h^n} \sum_{i=0}^n (-1)^i \binom{n}{i} f(x_{j+n-i})$$

approximates the n -th derivative. The method is of order 1.

Proposition VII.2.20

The n th-order backward difference quotient

$$\frac{1}{h^n} \sum_{i=0}^n (-1)^i \binom{n}{i} f(x - ih) = \frac{1}{h^n} \sum_{i=0}^n (-1)^i \binom{n}{i} f(x_{j-i})$$

approximates the n -th derivative. The method is of order 1.

$x = x_j$ needs to be chosen so the x_{j+n-i} / x_{j-i} are defined.

VII.2.3. Approximating the solution of a PDE

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$


Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

We proved the existence and uniqueness of $u \in C^2([0, 1])$.
(Exercise E.V.2 Lab 5)

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$:

$(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:

$(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


Define

- $c_j = c(x_j)$,
- $f_j = f(x_j)$.

And let u_j be the unknowns that will approximate $u(x_j)$.

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:


$(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

For $x = x_0$

$$u_0 = u(0) = 0$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider


$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$: $u_0 = 0$
 $(x_j)_{j \in \{0, \dots, J+1\}}, x_0 = 0, x_{J+1} = 0$.

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$: $u_0 = 0$
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

For $x = x_1$

$$-u''(x_1) + c(x_1)u(x_1) = f(x_1)$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:

$(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

u_0


$= 0$

For $x = x_1$

$$-\frac{u_2 - 2u_1 + u_0}{h^2} + c_1 u_1 = f_1$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$: $u_0 = 0$
 $(x_j)_{j \in \{0, \dots, J+1\}}, x_0 = 0, x_{J+1} = 0.$

For $x = x_1$

$$-\frac{u_2 - 2u_1}{h^2} + c_1 u_1 = f_1$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$: $u_0 = 0$
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 0$.

For $x = x_1$

$$\frac{2u_1 - u_2}{h^2} + c_1 u_1 = f_1$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$: $u_0 = 0$
 $(x_j)_{j \in \{0, \dots, J+1\}}, x_0 = 0, x_{J+1} = 0.$

For $x = x_1$

$$\left(\frac{2}{h^2} + c_1 \right) u_1 - \frac{1}{h^2} u_2 = f_1$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1 \right) u_1 - \frac{1}{h^2} u_2 &= f_1 \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \end{aligned}$$

For $x = x_2$

$$-u''(x_2) + c(x_2)u(x_2) = f(x_2)$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \end{aligned}$$

For $x = x_2$

$$-\frac{u_3 - 2u_2 + u_1}{h^2} + c_2 u_2 = f_2$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \end{aligned}$$

For $x = x_2$

$$-\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 = f_2$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$:

$(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$:

$(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


For $x = x_j$ with $j \in \{2, \dots, J-1\}$

$$-u''(x_j) + c(x_j)u(x_j) = f(x_j)$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ &\vdots \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


For $x = x_j$ with $j \in \{2, \dots, J-1\}$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ &\vdots \end{aligned}$$

$$-\frac{u_{j+1} - 2u_j + u_{j-1}}{h^2} + c_j u_j = f_j$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ &\vdots \end{aligned}$$

For $x = x_j$ with $j \in \{2, \dots, J-1\}$

$$-\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} = f_j$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \end{aligned}$$


Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \end{aligned}$$


$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$


For $x = x_I$

$$\begin{array}{rcl} u_0 & & = 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 & & = f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 & & = f_2 \\ & \vdots & \vdots \\ & \vdots & \vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} & & = f_j \\ & \vdots & \vdots \end{array}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


For $x = x_j$

$$-u''(x_j) + c(x_j)u(x_j) = f(x_j)$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


For $x = x_j$

$$-\frac{u_{j+1} - 2u_j + u_{j-1}}{h^2} + c_j u_j = f_j$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


For $x = x_j$

$$\frac{2u_j - u_{j-1}}{h^2} + c_j u_j = f_j$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.


For $x = x_j$

$$-\frac{1}{h^2}u_{j-1} + \left(\frac{2}{h} + c_j\right)u_j = f_j$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right)u_1 - \frac{1}{h^2}u_2 &= f_1 \\ -\frac{1}{h^2}u_1 + \left(\frac{2}{h^2} + c_2\right)u_2 - \frac{1}{h^2}u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2}u_{j-1} + \left(\frac{2}{h^2} + c_j\right)u_j - \frac{1}{h^2}u_{j+1} &= f_j \\ &\vdots \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$




Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \\ -\frac{1}{h^2} u_{J-1} + \left(\frac{2}{h^2} + c_J\right) u_J &= f_J \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$



Consider a uniform mesh on $[0, 1]$:
 $(x_j)_{j \in \{0, \dots, J+1\}}$, $x_0 = 0$, $x_{J+1} = 1$.

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1 \right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2 \right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j \right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \\ -\frac{1}{h^2} u_{J-1} + \left(\frac{2}{h^2} + c_J \right) u_J &= f_J \\ u_{J+1} &= 0 \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

$$U_h = [u_1, \dots, u_J]^*$$

$$F_h = [f_1, \dots, f_J]^*$$

$$A_h = \frac{1}{h^2} \begin{pmatrix} 2 & -1 & & & 0 \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ 0 & & -1 & 2 & -1 \\ & & & -1 & 2 \end{pmatrix}$$

$$C_h = \begin{pmatrix} c_1 & & & & 0 \\ & c_2 & & & \\ & & \ddots & & \\ & & & c_{J-1} & \\ 0 & & & & c_J \end{pmatrix}$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \\ -\frac{1}{h^2} u_{J-1} + \left(\frac{2}{h^2} + c_J\right) u_J &= f_J \\ u_{J+1} &= 0 \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

$$U_h = [u_1, \dots, u_J]^*$$

$$F_h = [f_1, \dots, f_J]^*$$

$$A_h = \frac{1}{h^2} \begin{pmatrix} 2 & -1 & & & 0 \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ 0 & & -1 & 2 & -1 \\ & & & -1 & 2 \end{pmatrix}$$

$$C_h = \begin{pmatrix} c_1 & & & & 0 \\ & c_2 & & & \\ & & \ddots & & \\ & & & c_{J-1} & \\ 0 & & & & c_J \end{pmatrix}$$

$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \\ -\frac{1}{h^2} u_{J-1} + \left(\frac{2}{h^2} + c_J\right) u_J &= f_J \\ u_{J+1} &= 0 \end{aligned}$$

Let $f \in C^0([0, 1])$ and $c \in C^0([0, 1], \mathbb{R}^+)$. Consider

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

$$U_h = [u_1, \dots, u_J]^*$$

$$F_h = [f_1, \dots, f_J]^*$$

$$A_h = \frac{1}{h^2} \begin{pmatrix} 2 & -1 & & & 0 \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ 0 & & & -1 & 2 \end{pmatrix}$$

$$C_h = \begin{pmatrix} c_1 & & & & 0 \\ & c_2 & & & \\ & & \ddots & & \\ & & & c_{J-1} & \\ 0 & & & & c_J \end{pmatrix}$$

$$(A_h + C_h)U_h = F_h \text{ and } u_0 = u_{J+1} = 0$$



$$\begin{aligned} u_0 &= 0 \\ \left(\frac{2}{h^2} + c_1\right) u_1 - \frac{1}{h^2} u_2 &= f_1 \\ -\frac{1}{h^2} u_1 + \left(\frac{2}{h^2} + c_2\right) u_2 - \frac{1}{h^2} u_3 &= f_2 \\ &\vdots \\ -\frac{1}{h^2} u_{j-1} + \left(\frac{2}{h^2} + c_j\right) u_j - \frac{1}{h^2} u_{j+1} &= f_j \\ &\vdots \\ -\frac{1}{h^2} u_{J-1} + \left(\frac{2}{h^2} + c_J\right) u_J &= f_J \\ u_{J+1} &= 0 \end{aligned}$$

Theorem VII.2.21

For all $J \geq 1$, the linear system

$$(A_h + C_h)U_h = F_h$$

has a unique solution U_h .

The proof was given in the lab session an hour ago
(Exercise E.V.2)

Theorem VII.2.21

For all $J \geq 1$, the linear system

$$(A_h + C_h)U_h = F_h$$

has a unique solution U_h .

The proof was given in the lab session an hour ago
(Exercise E.V.2)

Definition VII.2.22

The matrix A_h is the **FDM-Matrix of the Laplacian in dimension 1**.

Maximum Principle

Definition VII.2.23

We say that a vector v (resp. a matrix M) is non-negative if all of its components are non-negative. We note $v \geq 0$ (resp. $M \geq 0$).

Definition VII.2.24

*A matrix $M \in \mathcal{M}_q(\mathbb{R})$ is **monotone** if*

$$\forall v \in \mathbb{R}^q, Mv \geq 0 \Rightarrow v \geq 0$$

Maximum Principle

Definition VII.2.23

We say that a vector v (resp. a matrix M) is non-negative if all of its components are non-negative. We note $v \geq 0$ (resp. $M \geq 0$).

Definition VII.2.24

*A matrix $M \in \mathcal{M}_q(\mathbb{R})$ is **monotone** if*

$$\forall v \in \mathbb{R}^q, Mv \geq 0 \Rightarrow v \geq 0$$

Lemma VII.2.25

A monotone matrix is non-singular.

Maximum Principle

Definition VII.2.23

We say that a vector v (resp. a matrix M) is non-negative if all of its components are non-negative. We note $v \geq 0$ (resp. $M \geq 0$).

Definition VII.2.24

A matrix $M \in \mathcal{M}_q(\mathbb{R})$ is **monotone** if

$$\forall v \in \mathbb{R}^q, Mv \geq 0 \Rightarrow v \geq 0$$

Lemma VII.2.25

A monotone matrix is non-singular.

Lemma VII.2.26

A matrix M is monotone iff $M^{-1} > 0$.

Maximum Principle

Lemma VII.2.27

$A_h + C_h$ is monotone.

Proof: Lab session to come.

Maximum Principle

Lemma VII.2.27

$A_h + C_h$ is monotone.

Proof: Lab session to come.

If $F_h \geq 0$

Maximum Principle

Lemma VII.2.27

$A_h + C_h$ is monotone.

Proof: Lab session to come.

If $F_h \geq 0$ then $(A_h + C_h)U_h \geq 0$

Maximum Principle

Lemma VII.2.27

$A_h + C_h$ is monotone.

Proof: Lab session to come.

If $F_h \geq 0$ then $(A_h + C_h)U_h \geq 0$ then $U_h \geq 0$.

Maximum Principle

Lemma VII.2.27

$A_h + C_h$ is monotone.

Proof: Lab session to come.

If $F_h \geq 0$ then $(A_h + C_h)U_h \geq 0$ then $U_h \geq 0$.

Theorem VII.2.28 (Maximum Principle)

If $f \geq 0$ then U_h has only non-negative components.

Maximum Principle

Lemma VII.2.27

$A_h + C_h$ is monotone.

Proof: Lab session to come.

If $F_h \geq 0$ then $(A_h + C_h)U_h \geq 0$ then $U_h \geq 0$.

Theorem VII.2.28 (Maximum Principle)

If $f \geq 0$ then U_h has only non-negative components.

Compare with Theorem V.2.6.

VII.2.4. Convergence

Let's get rid of a possible misconception

U_h is **not** a discretization of the solution.

.

Let's get rid of a possible misconception

U_h is **not** a discretization of the solution.

U_h does **not** hold the values of the u at the nodes of the grid:

.

Let's get rid of a possible misconception

U_h is **not** a discretization of the solution.

U_h does **not** hold the values of the u at the nodes of the grid:
 u_j approximates $u(x_j)$ (or so is our wish).

Let's get rid of a possible misconception

U_h is **not** a discretization of the solution.

U_h does **not** hold the values of the u at the nodes of the grid:
 u_j approximates $u(x_j)$ (or so is our wish).

We have discretized the problem, not the solution.

Let's get rid of a possible misconception

U_h is **not** a discretization of the solution.

U_h does **not** hold the values of the u at the nodes of the grid:
 u_j approximates $u(x_j)$ (or so is our wish).

We have discretized the problem, not the solution.

Now, we need to compare u_j and $u(x_j)$
and verify if the former approximates the latter.

Projection onto the mesh

Define the projection Π_h of a function on the mesh by its pointwise evaluation on the nodes (x_j) :

$$\begin{aligned}\Pi_h : C([0, 1]) &\text{ to } \mathbb{R}^J \\ u &\mapsto [u(x_j)]_{j \in \{1, \dots, J\}}\end{aligned}$$

Projection onto the mesh

Define the projection Π_h of a function on the mesh by its pointwise evaluation on the nodes (x_j) :

$$\begin{aligned}\Pi_h : C([0, 1]) &\text{ to } \mathbb{R}^J \\ u &\mapsto [u(x_j)]_{j \in \{1, \dots, J\}}\end{aligned}$$

We have $F_h = \Pi_h(f)$

Projection onto the mesh

Define the projection Π_h of a function on the mesh by its pointwise evaluation on the nodes (x_j) :

$$\begin{aligned}\Pi_h : C([0, 1]) &\text{ to } \mathbb{R}^J \\ u &\mapsto [u(x_j)]_{j \in \{1, \dots, J\}}\end{aligned}$$

We have $F_h = \Pi_h(f)$ and $C_h = \Pi_h(c)$.

Projection onto the mesh

Define the projection Π_h of a function on the mesh by its pointwise evaluation on the nodes (x_j) :

$$\begin{aligned}\Pi_h : C([0, 1]) &\text{ to } \mathbb{R}^J \\ u &\mapsto [u(x_j)]_{j \in \{1, \dots, J\}}\end{aligned}$$

We have $F_h = \Pi_h(f)$ and $C_h = \Pi_h(c)$.

We don't have $U_h = \Pi_h(u)$ in general

Projection onto the mesh

Define the projection Π_h of a function on the mesh by its pointwise evaluation on the nodes (x_j) :

$$\begin{aligned}\Pi_h : C([0, 1]) &\text{ to } \mathbb{R}^J \\ u &\mapsto [u(x_j)]_{j \in \{1, \dots, J\}}\end{aligned}$$

We have $F_h = \Pi_h(f)$ and $C_h = \Pi_h(c)$.

We don't have $U_h = \Pi_h(u)$ in general, let us measure the gap:

$$E_h = U_h - \Pi_h u = \begin{pmatrix} u_1 - u(x_1) \\ \vdots \\ u_J - u(x_J) \end{pmatrix}$$

Convergence

Definition VII.2.29

*The numerical method is **convergent** if*

$$\lim_{h \rightarrow 0} \|E_h\|_h = 0$$

where $\|\cdot\|_h$ is a norm on \mathbb{R}^J .

Convergence

Definition VII.2.29

*The numerical method is **convergent** if*

$$\lim_{h \rightarrow 0} \|E_h\|_h = 0$$

where $\|\cdot\|_h$ is a norm on \mathbb{R}^J .

Furthermore,

Convergence

Definition VII.2.29

*The numerical method is **convergent** if*

$$\lim_{h \rightarrow 0} \|E_h\|_h = 0$$

where $\|\cdot\|_h$ is a norm on \mathbb{R}^J .

*Furthermore, if $\|E_h\|_h = O(h^p)$ the method is said to be of **order** p*

Consistency

Definition VII.2.30

Define

$$\mathcal{E}_h = A_h \Pi_h u - F_h \in \mathbb{R}^J$$

*We say the Finite Difference Method is **consistent** with the PDE if*

$$\lim_{h \rightarrow 0} \max_{j \in \{1, \dots, J\}} |(\mathcal{E}_h)_j| = \lim_{h \rightarrow 0} \|\mathcal{E}_h\|_\infty = 0$$

Consistency

Definition VII.2.30

Define

$$\mathcal{E}_h = A_h \Pi_h u - F_h \in \mathbb{R}^J$$

We say the Finite Difference Method is **consistent** with the PDE if

$$\lim_{h \rightarrow 0} \max_{j \in \{1, \dots, J\}} |(\mathcal{E}_h)_j| = \lim_{h \rightarrow 0} \|\mathcal{E}_h\|_\infty = 0$$

Furthermore,

Consistency

Definition VII.2.30

Define

$$\mathcal{E}_h = A_h \Pi_h u - F_h \in \mathbb{R}^J$$

We say the Finite Difference Method is **consistent** with the PDE if

$$\lim_{h \rightarrow 0} \max_{j \in \{1, \dots, J\}} |(\mathcal{E}_h)_j| = \lim_{h \rightarrow 0} \|\mathcal{E}_h\|_\infty = 0$$

Furthermore, the consistency is of order p if $\|\mathcal{E}_h\|_\infty = O(h^p)$

Consistency

Definition VII.2.30

Define

$$\mathcal{E}_h = A_h \Pi_h u - F_h \in \mathbb{R}^J$$

We say the Finite Difference Method is **consistent** with the PDE if

$$\lim_{h \rightarrow 0} \max_{j \in \{1, \dots, J\}} |(\mathcal{E}_h)_j| = \lim_{h \rightarrow 0} \|\mathcal{E}_h\|_\infty = 0$$

Furthermore, the consistency is of order p if $\|\mathcal{E}_h\|_\infty = O(h^p)$

What follows makes no mathematical sense but tries to describe the meaning of this definition:

$$\text{consistency} \iff \lim_{\text{mesh size} \rightarrow 0} (\text{PDE} - \text{FDM}) = 0$$

Consistency

Example

In our earlier problem

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

Consistency

Example

In our earlier problem

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

take $f \in C^2(0, 1)$ and $c = 0$ (Poisson's equation in dimension 1).

Consistency

Example

In our earlier problem

$$\begin{cases} -u''(x) + c(x)u(x) = f(x) & x \in]0, 1[\\ u(0) = u(1) = 0 \end{cases}$$

*take $f \in C^2(0, 1)$ and $c = 0$ (Poisson's equation in dimension 1).
Then $u \in C^4(0, 1)$.*

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0,1)$. The solution $u \in C^4(0,1)$.

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0,1)$. The solution $u \in C^4(0,1)$.

The second-order central difference quotient is a finite difference approximation of the second derivative of order 2:

$$\frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) = O(h^2)$$

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0,1)$. The solution $u \in C^4(0,1)$.

The second-order central difference quotient is a finite difference approximation of the second derivative of order 2:

$$\frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) = O(h^2)$$

Using the Taylor-Lagrange inequality, we could be more precise:

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0, 1)$. The solution $u \in C^4(0, 1)$.

The second-order central difference quotient is a finite difference approximation of the second derivative of order 2:

$$\frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) = O(h^2)$$

Using the Taylor-Lagrange inequality, we could be more precise:

$$\left| \frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) \right| \leq \|u^{(4)}\|_{\infty} \frac{h^2}{12}$$

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0, 1)$. The solution $u \in C^4(0, 1)$.

The second-order central difference quotient is a finite difference approximation of the second derivative of order 2:

$$\frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) = O(h^2)$$

Using the Taylor-Lagrange inequality, we could be more precise:

$$\max_{j \in \{1, \dots, J\}} \left| \frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) \right| \leq \|u^{(4)}\|_{\infty} \frac{h^2}{12}$$

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0, 1)$. The solution $u \in C^4(0, 1)$.

The second-order central difference quotient is a finite difference approximation of the second derivative of order 2:

$$\frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) = O(h^2)$$

Using the Taylor-Lagrange inequality, we could be more precise:

$$\max_{j \in \{1, \dots, J\}} \left| \frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) \right| \leq \|u^{(4)}\|_{\infty} \frac{h^2}{12}$$

The FDM is consistent with the PDE in the order 2.

Consistency

Example

Consider the Poisson equation in dimension 1 with a right hand side $f \in C^2(0, 1)$. The solution $u \in C^4(0, 1)$.

The second-order central difference quotient is a finite difference approximation of the second derivative of order 2:

$$\frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) = O(h^2)$$

Using the Taylor-Lagrange inequality, we could be more precise:

$$\max_{j \in \{1, \dots, J\}} \left| \frac{u(x_{j+1}) - 2u(x_j) + u(x_{j-1}))}{h^2} - u''(x_j) \right| \leq \|u^{(4)}\|_{\infty} \frac{h^2}{12}$$

The FDM is consistent with the PDE in the order 2.

Note: The remainder vanishes if $u \in \mathbb{R}_3[X]$.

Stability

Definition VII.2.31

Consider a numerical method $A_h U_h = F_h$ where

- U_h is the the unknown,
- F_h is the input data and,
- A_h is a non-singular matrix.

Consider $\| \cdot \|$ the matrix norm induced by the norm $\| \cdot \|$ on \mathbb{R}^J .

The numerical method is **stable** if there exists a constant C independent of J such that

$$\|A_h^{-1}\| \leq C.$$

Stability

Definition VII.2.31

Consider a numerical method $A_h U_h = F_h$ where

- U_h is the unknown,
- F_h is the input data and,
- A_h is a non-singular matrix.

Consider $\| \cdot \|$ the matrix norm induced by the norm $\| \cdot \|$ on \mathbb{R}^J .

The numerical method is **stable** if there exists a constant C independent of J such that

$$\|A_h^{-1}\| \leq C.$$

Stability means small errors in the input data F_h don't get out of hand.

Stability

The convergence in \mathbb{R}^d does not depend on the norm we choose, since \mathbb{R}^d has a finite dimension.

However, since C must **not** depend on h (i.e. must not depend on $d = J$), we can have stability for a norm and not for another norm. Therefore, it is necessary to say which norm $\| \cdot \|$ we take for stability (and later for convergence).

Convergence

$$E_h = U_h - \Pi_h u$$

Convergence

$$A_h E_h = A_h U_h - A_h \Pi_h u$$

Convergence

$$A_h E_h = F_h - A_h \Pi_h u$$

Convergence

$$A_h E_h = -\mathcal{E}_h$$

Convergence

$$E_h = -A_h^{-1} \mathcal{E}_h$$

Convergence

$$\|E_h\| = \| -A_h^{-1} \mathcal{E}_h \|$$

Convergence

$$\|E_h\| = \|A_h^{-1}\mathcal{E}_h\|$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$(\text{Stability}) \quad \|A_h^{-1}\| \leq C$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\text{(Stability)} \quad \|A_h^{-1}\| \leq C$$

$$\text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\left. \begin{array}{l} \text{(Stability)} \quad \|A_h^{-1}\| \leq C \\ \text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0 \end{array} \right\} \Rightarrow$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\left. \begin{array}{l} \text{(Stability)} \quad \|A_h^{-1}\| \leq C \\ \text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0 \end{array} \right\} \Rightarrow \lim_{h \rightarrow 0} \|E_h\| = 0 \text{ (Convergence)}$$

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\left. \begin{array}{l} \text{(Stability)} \quad \|A_h^{-1}\| \leq C \\ \text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0 \end{array} \right\} \Rightarrow \lim_{h \rightarrow 0} \|E_h\| = 0 \text{ (Convergence)}$$

Theorem VII.2.32

*A finite difference method that is both **consistent** and **stable** is **convergent**.*

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\left. \begin{array}{l} \text{(Stability)} \quad \|A_h^{-1}\| \leq C \\ \text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0 \end{array} \right\} \Rightarrow \lim_{h \rightarrow 0} \|E_h\| = 0 \text{ (Convergence)}$$

Theorem VII.2.32

*A finite difference method that is both **consistent** and **stable** is **convergent**.*

PDE \longrightarrow FDM \longrightarrow Solution

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\left. \begin{array}{l} \text{(Stability)} \quad \|A_h^{-1}\| \leq C \\ \text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0 \end{array} \right\} \Rightarrow \lim_{h \rightarrow 0} \|E_h\| = 0 \text{ (Convergence)}$$

Theorem VII.2.32

*A finite difference method that is both **consistent** and **stable** is **convergent**.*

PDE \longrightarrow FDM \longrightarrow Solution

Convergence

$$\|E_h\| \leq \|A_h^{-1}\| \|\mathcal{E}_h\|$$

$$\left. \begin{array}{l} \text{(Stability)} \quad \|A_h^{-1}\| \leq C \\ \text{(Consistency)} \quad \lim_{h \rightarrow 0} \|\mathcal{E}_h\| = 0 \end{array} \right\} \Rightarrow \lim_{h \rightarrow 0} \|E_h\| = 0 \text{ (Convergence)}$$

Theorem VII.2.32

*A finite difference method that is both **consistent** and **stable** is **convergent**.*

PDE \longrightarrow FDM \longrightarrow Solution

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let $[A_h^{-1}]_{ij}$ be the component of A_h^{-1} on line i and column j .

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let $[A_h^{-1}]_{ij}$ be the component of A_h^{-1} on line i and column j .
We have established: (last lecture)

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J |[A_h^{-1}]_{ij}|$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let $[A_h^{-1}]_{ij}$ be the component of A_h^{-1} on line i and column j .
We know that A_h is monotone so $A_h^{-1} \geq 0$ therefore

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij}$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let i be an integer in $[1, J]$

and $g \in \mathbb{R}^J$ be a vector for which each component $g_j = 1$.

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

We can look for a solution in $\mathbb{R}_2[x]$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Let $u(x) = \frac{1}{2}(-x^2 + x)$. It is a solution.

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Let $u(x) = \frac{1}{2}(-x^2 + x)$. It is a solution.

We have $1(x_i) = g_i$ so $A_h w = g$ is the discretization of $-u'' = 1$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Let $u(x) = \frac{1}{2}(-x^2 + x)$. It is a solution.

We have $1(x_i) = g_i$ so $A_h w = g$ is the discretization of $-u'' = 1$

Since $u \in \mathbb{R}_2[x]$, we have $u''(x_i) = \frac{u(x_{i+1}) - 2u(x_i) + u(x_{i-1}))}{h^2}$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Let $u(x) = \frac{1}{2}(-x^2 + x)$. It is a solution.

We have $1(x_i) = g_i$ so $A_h w = g$ is the discretization of $-u'' = 1$

Since $u \in \mathbb{R}_2[x]$, we have $u''(x_i) = \frac{u(x_{i+1}) - 2u(x_i) + u(x_{i-1}))}{h^2}$

It follows that $w_i = u(x_i)$.

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Let $u(x) = \frac{1}{2}(-x^2 + x)$. It is a solution.

We have $1(x_i) = g_i$ so $A_h w = g$ is the discretization of $-u'' = 1$

Since $u \in \mathbb{R}_2[x]$, we have $u''(x_i) = \frac{u(x_{i+1}) - 2u(x_i) + u(x_{i-1}))}{h^2}$

It follows that $w_i = u(x_i)$. Since $A_h w = g$ we have $w = A_h^{-1} g$.

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Consider $-u'' = 1$ with the initial condition $u(0) = u(1) = 0$.

Let $u(x) = \frac{1}{2}(-x^2 + x)$. It is a solution.

We have $1(x_i) = g_i$ so $A_h w = g$ is the discretization of $-u'' = 1$

Since $u \in \mathbb{R}_2[x]$, we have $u''(x_i) = \frac{u(x_{i+1}) - 2u(x_i) + u(x_{i-1}))}{h^2}$

It follows that $w_i = u(x_i)$. Since $A_h w = g$ we have $w = A_h^{-1} g$.

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$w_i = u(x_i)$.

$w = A_h^{-1}g$.

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$w_i = u(x_i)$.

$w = A_h^{-1}g. \Rightarrow \sum_{j=1}^J [A_h^{-1}]_{ij} g_j = w_i$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$w_i = u(x_i)$.

$$\sum_{j=1}^J [A_h^{-1}]_{ij} g_j = w_i$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

$$\|A_h\|_{\infty} = \max_{1 \leq i \leq J} \sum_{j=1}^J [A_h^{-1}]_{ij} g_j$$

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$$\sum_{j=1}^J [A_h^{-1}]_{ij} g_j = u(x_i)$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$$\|A\|_{\infty} = \max_{1 \leq i \leq J} u(x_i)$$

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$$\|A\|_{\infty} = \max_{1 \leq i \leq J} u(x_i)$$

u reaches its maximum at $\frac{1}{2}$ and $u(\frac{1}{2}) = \frac{1}{8}$ thus

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Proof:

Let $u(x) = \frac{1}{2}(-x^2 + x)$.

$$\|A\|_{\infty} = \max_{1 \leq i \leq J} u(x_i)$$

u reaches its maximum at $\frac{1}{2}$ and $u(\frac{1}{2}) = \frac{1}{8}$ thus

$$\|A\|_{\infty} \leq \frac{1}{8}$$

QED

Stability

Theorem VII.2.33

Let A_h be the matrix associated to the Poisson's equation in dimension 1.

$$\forall J \geq 1, \quad \|A_h^{-1}\|_{\infty} \leq \frac{1}{8}.$$

Corollary VII.2.34

The finite difference method for the Poisson's problem in dimension 1 converges in $\|\cdot\|_{\infty}$. The order is 2.

Programming the FDM for $f = 1$

```
import numpy as np
from scipy import sparse
from scipy.sparse.linalg import dsolve
import matplotlib
import matplotlib.pyplot as plt
```

```
J = 10
h = 1.0/(J+1)
```

```
# Building Ah
```

```
diagonal = np.ones(J)*2.0
side_diagonal = np.ones(J-1)*(-1.0)
h2_Ah = sparse.diags([side_diagonal, diagonal, side_diagonal],
                    [-1,0,1], format="csr")
Ah = h2_Ah*(1/(h**2))
```

```
print("h2_Ah_\n", h2_Ah.A)
print("Ah_\n", Ah.A)
```

```

h2_Ah =
[[ 2. -1.  0.  0.  0.  0.  0.  0.  0.  0.]
 [-1.  2. -1.  0.  0.  0.  0.  0.  0.  0.]
 [ 0. -1.  2. -1.  0.  0.  0.  0.  0.  0.]
 [ 0.  0. -1.  2. -1.  0.  0.  0.  0.  0.]
 [ 0.  0.  0. -1.  2. -1.  0.  0.  0.  0.]
 [ 0.  0.  0.  0. -1.  2. -1.  0.  0.  0.]
 [ 0.  0.  0.  0.  0. -1.  2. -1.  0.  0.]
 [ 0.  0.  0.  0.  0.  0. -1.  2. -1.  0.]
 [ 0.  0.  0.  0.  0.  0.  0. -1.  2. -1.]
 [ 0.  0.  0.  0.  0.  0.  0.  0. -1.  2.]]


Ah =
[[ 242. -121.  0.  0.  0.  0.  0.  0.  0.  0.]
 [-121.  242. -121.  0.  0.  0.  0.  0.  0.  0.]
 [ 0. -121.  242. -121.  0.  0.  0.  0.  0.  0.]
 [ 0.  0. -121.  242. -121.  0.  0.  0.  0.  0.]
 [ 0.  0.  0. -121.  242. -121.  0.  0.  0.  0.]
 [ 0.  0.  0.  0. -121.  242. -121.  0.  0.  0.]
 [ 0.  0.  0.  0.  0. -121.  242. -121.  0.  0.]
 [ 0.  0.  0.  0.  0.  0. -121.  242. -121.  0.]
 [ 0.  0.  0.  0.  0.  0.  0. -121.  242. -121.]
 [ 0.  0.  0.  0.  0.  0.  0.  0. -121.  242.]]
    
```

Programming the FDM for $f = 1$

```
# Building b  
b = np.ones(J)  
  
# Solving for u  
u = dsolve.spsolve(Ah, b)  
  
# Plotting the solution  
x = np.linspace(0.0, 1.0, num=J+2)  
u = np.concatenate(([0], u, [0]))  
fig = plt.figure()  
ax = fig.gca()  
ax.plot(x, u)  
plt.show()
```


Approximation of the solution

$$J = 10$$




Approximation of the solution

$$J = 100$$



Approximation of the solution

$$J = 500$$




Approximation of the solution



Program available for download at

<https://cagnol.link/fdmp1>



Estimating the error



Estimating the error





Estimating the error



Slope of the line on the log-scale graph: 1.97

Estimating the error



Slope of the line on the log-scale graph: 1.97

Program available for download at <https://cagnol.link/fdmp1cr>

VII.3. FDM in dimension $d > 1$

VII.3.1. Meshing

Structured vs. Unstructured meshes

For **Finite Element Method** we often mesh using simplices (triangles in 2D).

The mesh does not follow a pattern structure.

Structured vs. Unstructured meshes

For **Finite Element Method** we often mesh using simplices (triangles in 2D).

The mesh does not follow a pattern structure.

For **Finite Difference Method**, we often mesh using orthotopes (rectangles in 2D)

The mesh follows a pattern structure: we tessellate Ω .

Structured vs. Unstructured meshes

For **Finite Element Method** we often mesh using simplices (triangles in 2D).

The mesh does not follow a pattern structure.

For **Finite Difference Method**, we often mesh using orthotopes (rectangles in 2D)

The mesh follows a pattern structure: we tessellate Ω .

Definition VII.3.1

A **structured mesh** is a mesh that can be produced by replicating an elementary cell.

A mesh that is not structured is called **unstructured**.

With a structured mesh, every vertex can be easily numbered by a d -tuple: (i, j) if $d = 2$ and (i, j, k) if $d = 3$.

Example of a structured mesh


Consider a square $\Omega \subset \mathbb{R}^2$:

We tessellate Ω with $(J + 1)^2$ elementary squares.

$J + 2$ points subdivide $[0, 1]$ on the x -axis.

$J + 2$ points subdivide $[0, 1]$ on the y -axis.

There are J^2 points (in red) that are not on $\partial\Omega$. Points $P_n = (x_j, y_j)$ are numbered from 1 to J^2 with $n = J(i - 1) + j$




$J = 9$

44/62


Example of a structured mesh

For a more complicated $\Omega \subset \mathbb{R}^2$



Example of a structured mesh

An example for $\Omega \subset \mathbb{R}^3$



Credit: CFDyna

VII.3.2. Finite Difference Approximation of the Partial Derivatives

First order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions.

First order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions.

$$\frac{\partial f}{\partial x}(x, y) = \frac{f(x+h, y) - f(x, y)}{h} + O(h) \quad (\text{forward})$$

$$\frac{\partial f}{\partial x}(x, y) = \frac{f(x, y) - f(x-h, y)}{h} + O(h) \quad (\text{backward})$$

$$\frac{\partial f}{\partial x}(x, y) = \frac{f(x+\frac{h}{2}, y) - f(x-\frac{h}{2}, y)}{h} + O(h^2) \quad (\text{central})$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{f(x, y+k) - f(x, y)}{k} + O(k) \quad (\text{forward})$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{f(x, y) - f(x, y-k)}{k} + O(k) \quad (\text{backward})$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{f(x, y+\frac{k}{2}) - f(x, y-\frac{k}{2})}{k} + O(k^2) \quad (\text{central}))$$

First order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions.
 For $x = x_i$ and $y = y_j$

$$\frac{\partial f}{\partial x}(x_j, y_i) = \frac{f(x_{j+1}, y_i) - f(x_j, y_i)}{x_{j+1} - x_j} + O(h) \quad (\text{forward})$$

$$\frac{\partial f}{\partial x}(x_j, y_i) = \frac{f(x_j, y_i) - f(x_{j-1}, y_i)}{x_i - x_{j-1}} + O(h) \quad (\text{backward})$$

$$\frac{\partial f}{\partial x}(x_j, y_i) = \frac{f(x_{j+1}, y_i) - f(x_{j-1}, y_i)}{x_{j+1} - x_{j-1}} + O(h^2) \quad (\text{central})$$







$$\frac{\partial f}{\partial y}(x_j, y_i) = \frac{f(x_j, y_{i+1}) - f(x_j, y_i)}{y_{i+1} - y_i} + O(k) \quad (\text{forward})$$

$$\frac{\partial f}{\partial y}(x_j, y_i) = \frac{f(x_j, y_i) - f(x_j, y_{i-1})}{y_i - y_{i-1}} + O(k) \quad (\text{backward})$$

$$\frac{\partial f}{\partial y}(x_j, y_i) = \frac{f(x_j, y_{i+1}) - f(x_j, y_{i-1})}{y_{i+1} - y_{i-1}} + O(k^2) \quad (\text{central})$$

First order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions.
For $x = x_i$ and $y = y_j$, the stencils are:

	Fw.	Bw.	Ct.
$\partial f / \partial x$			
$\partial f / \partial y$			

Second order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions.
The central difference quotient for the second derivative are given by:

Second order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions. The central difference quotient for the second derivative are given by:

$$\frac{\partial^2 f}{\partial x^2}(x, y) = \frac{f(x+h, y) - 2f(x, y) + f(x-h, y)}{h^2} + O(h^2)$$

$$\frac{\partial^2 f}{\partial y^2}(x, y) = \frac{f(x, y+k) - 2f(x, y) + f(x, y-k)}{k^2} + O(k^2)$$

$$\begin{aligned} \frac{\partial^2 f}{\partial x \partial y}(x, y) &= \frac{f(x+\frac{h}{2}, y+\frac{k}{2}) + f(x-\frac{h}{2}, y-\frac{k}{2})}{hk} \\ &\quad - \frac{f(x+\frac{h}{2}, y-\frac{k}{2}) + f(x-\frac{h}{2}, y+\frac{k}{2})}{hk} + O(h^2, k^2). \end{aligned}$$

You can adapt these formulas to forward and backward difference quotients.

Second order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions. For $x = x_i$ and $y = y_j$. The central difference quotient for the second derivative are given by:

$$\frac{\partial^2 f}{\partial x^2}(x_j, y_i) = \frac{f(x_{j+1}, y_i) - 2f(x_j, y_i) + f(x_{j-1}, y_i)}{(x_{j+1} - x_{j-1})^2} + O(h^2)$$



$$\frac{\partial^2 f}{\partial y^2}(x_j, y_i) = \frac{f(x_j, y_{i+1}) - 2f(x_j, y_i) + f(x_j, y_{i-1})}{(y_{i+1} - y_{i-1})^2} + O(k^2)$$

$$\begin{aligned} \frac{\partial^2 f}{\partial x \partial y}(x_j, y_i) &= \frac{f(x_{j+1}, y_{i+1}) + f(x_{j-1}, y_{i-1})}{(x_{j+1} - x_{j-1})(y_{i+1} - y_{i-1})} \\ &\quad - \frac{f(x_{j+1}, y_{i-1}) + f(x_{j-1}, y_{i+1})}{(x_{j+1} - x_{j-1})(y_{i+1} - y_{i-1})} + O(h^2, k^2). \end{aligned}$$

You can adapt these formulas to forward and backward difference quotients.

Second order Partial Derivatives for $d = 2$

What we did in dimension one generalizes to higher dimensions.
 For $x = x_i$ and $y = y_j$. The stencils are:



Laplace Operator for $d = 2$

Since $\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$ we have

$$\begin{aligned}\Delta f(x, y) = & \frac{f(x+h, y) - 2f(x, y) + f(x-h, y)}{h^2} \\ & + \frac{f(x, y+k) - 2f(x, y) + f(x, y-k)}{k^2} + O(h^2, k^2)\end{aligned}$$

Laplace Operator for $d = 2$

Since $\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$ we have


$$\begin{aligned}\Delta f(x_j, y_i) &= \frac{f(x_{j+1}, y_i) - 2f(x_j, y_i) + f(x_{j-1}, y_i)}{(x_{j+1} - x_{j-1})^2} \\ &\quad + \frac{f(x_j, y_{i+1}) - 2f(x_j, y_i) + f(x_j, y_{i-1}))}{(y_{i+1} - y_{i-1})^2} + O(h^2, k^2)\end{aligned}$$

Laplace Operator for $d = 2$

Since $\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$ we have

$$\Delta f(x_j, y_i) = \frac{f(x_{j+1}, y_i) - 2f(x_j, y_i) + f(x_{j-1}, y_i)}{(x_{j+1} - x_{j-1})^2} + \frac{f(x_j, y_{i+1}) - 2f(x_j, y_i) + f(x_j, y_{i-1})}{(y_{i+1} - y_{i-1})^2} + O(h^2, k^2)$$


The (five-point) stencil is



VII.3.3. Approximating the PDE


Let us discretize Δu

Let us discretize Δu



$$\frac{1}{h^2} ((u(x_{j+1}, y_i) - 2u(x_j, y_i) + u(x_{j-1}, y_i)) + (u(x_j, y_{i+1}) - 2u(x_j, y_i) + u(x_j, y_{i-1})))$$


Let us discretize Δu



$$\frac{1}{h^2} ((u(x_{j+1}, y_i) - 2u(x_j, y_i) + u(x_{j-1}, y_i)) + (u(x_j, y_{i+1}) - 2u(x_j, y_i) + u(x_j, y_{i-1})))$$


$$\frac{1}{h^2} (u(x_j, y_{i-1}) + u(x_{j-1}, y_i) - 4u(x_j, y_i) + u(x_{j+1}, y_i) + (u(x_j, y_{i+1})))$$

Let us discretize Δu




$$\frac{1}{h^2} (u(x_j, y_{i-1}) + u(x_{j-1}, y_i) - 4u(x_j, y_i) + u(x_{j+1}, y_i) + (u(x_j, y_{i+1})))$$

Let us discretize Δu




$$\frac{1}{h^2} (u(x_j, y_{i-1}) + u(x_{j-1}, y_i) - 4u(x_j, y_i) + u(x_{j+1}, y_i) + (u(x_j, y_{i+1})))$$
$$\frac{1}{h^2} (u_{j,i-1} + u_{j-1,i} - 4u_{j,i} + u_{j+1,i} + u_{j,i+1})$$

Let us discretize Δu



$$\frac{1}{h^2}(u_{j,i-1} + u_{j-1,i} - 4u_{j,i} + u_{j+1,i} + u_{j,i+1})$$

Let us discretize Δu



$$\frac{1}{h^2}(u_{j,i-1} + u_{j-1,i} - 4u_{j,i} + u_{j+1,i} + u_{j,i+1})$$

Going from dual-index to single-index with $(i, j) \mapsto n = J(i-1) + j$

$$\frac{1}{h^2}(u_{j+J(i-2)} + u_{j-1+J(i-1)} - 4u_{j+J(i-1)} + u_{j+1+J(i-1)} + u_{j+Ji})$$

where u_n is replaced by zero for all out-of-range n .

The discretization of $-\Delta u = f$ is given by:

$$\frac{1}{h^2}(-u_{j+J(i-2)} - u_{j-1+J(i-1)} + 4u_{j+J(i-1)} - u_{j+1+J(i-1)} - u_{j+Ji}) = f_{j-1+J(i-1)}$$

The discretization of $-\Delta u = f$ is given by:

$$\frac{1}{h^2}(-u_{j+J(i-2)} - u_{j-1+J(i-1)} + 4u_{j+J(i-1)} - u_{j+1+J(i-1)} - u_{j+Ji}) = f_{j-1+J(i-1)}$$

$$A_h = \frac{1}{h^2} \begin{pmatrix} \mathbf{T} & -\mathbf{I} & 0 & \dots & 0 \\ -\mathbf{I} & \mathbf{T} & -\mathbf{I} & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & -\mathbf{I} & \mathbf{T} & -\mathbf{I} \\ 0 & \dots & 0 & -\mathbf{I} & \mathbf{T} \end{pmatrix} \quad \text{with}$$

$$\mathbf{T} = \begin{pmatrix} 4 & -1 & 0 & \dots & 0 \\ -1 & 4 & -1 & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & -1 & 4 & -1 \\ 0 & \dots & 0 & -1 & 4 \end{pmatrix}$$

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & \dots & 0 \\ 0 & 1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}$$

```
import numpy as np
from scipy import sparse

J = 4

diagonal_T = np.ones(J**2)*4.0

side_diagonal_T = np.ones(J**2-1)*(-1.0)
side_diagonal_T[np.arange(1,J**2)%J==0] = 0


diagonal_I = np.ones(J**2-J)

h2_Ah = sparse.diags([-diagonal_I ,
                     side_diagonal_T ,
                     diagonal_T ,
                     side_diagonal_T ,
                     -diagonal_I],
                    [-J,-1,0, 1,J], format="csr")

print(h2_Ah.A)
```

```
[ [ 4. -1.  0.  0. -1.  0.  0.  0.  0.  0.  0.  0.  0.  0.  0.  0.]
  [-1.  4. -1.  0.  0. -1.  0.  0.  0.  0.  0.  0.  0.  0.  0.  0.]
  [ 0. -1.  4. -1.  0.  0. -1.  0.  0.  0.  0.  0.  0.  0.  0.  0.]
  [ 0.  0. -1.  4.  0.  0.  0. -1.  0.  0.  0.  0.  0.  0.  0.  0.]
  [-1.  0.  0.  0.  4. -1.  0.  0. -1.  0.  0.  0.  0.  0.  0.  0.]
  [ 0. -1.  0.  0. -1.  4. -1.  0.  0. -1.  0.  0.  0.  0.  0.  0.]
  [ 0.  0. -1.  0.  0. -1.  4. -1.  0.  0. -1.  0.  0.  0.  0.  0.]
  [ 0.  0.  0. -1.  0.  0. -1.  4.  0.  0.  0. -1.  0.  0.  0.  0.]
  [ 0.  0.  0.  0. -1.  0.  0.  0.  4. -1.  0.  0. -1.  0.  0.  0.]
  [ 0.  0.  0.  0.  0. -1.  0.  0. -1.  4. -1.  0.  0. -1.  0.  0.]
  [ 0.  0.  0.  0.  0.  0. -1.  0.  0. -1.  4. -1.  0.  0. -1.  0.]
  [ 0.  0.  0.  0.  0.  0.  0. -1.  0.  0. -1.  4.  0.  0.  0. -1.]
  [ 0.  0.  0.  0.  0.  0.  0.  0. -1.  0.  0.  0.  4. -1.  0.  0.]
  [ 0.  0.  0.  0.  0.  0.  0.  0.  0. -1.  0.  0. -1.  4. -1.  0.]
  [ 0.  0.  0.  0.  0.  0.  0.  0.  0.  0. -1.  0.  0. -1.  4. -1.]
  [ 0.  0.  0.  0.  0.  0.  0.  0.  0.  0.  0. -1.  0.  0. -1.  4.]]
```

[[4.	-1.	0.	0.	-1.	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.]]
[[-1.	4.	-1.	0.	0.	-1.	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.]]
[[0.	-1.	4.	-1.	0.	0.	-1.	0.	0.	0.	0.	0.	0.	0.	0.	0.]]
[[0.	0.	-1.	4.	0.	0.	0.	-1.	0.	0.	0.	0.	0.	0.	0.	0.]]
[[-1.	0.	0.	0.	4.	-1.	0.	0.	-1.	0.	0.	0.	0.	0.	0.	0.]]
[[0.	-1.	0.	0.	-1.	4.	-1.	0.	0.	-1.	0.	0.	0.	0.	0.	0.]]
[[0.	0.	-1.	0.	0.	-1.	4.	-1.	0.	0.	-1.	0.	0.	0.	0.	0.]]
[[0.	0.	0.	-1.	0.	0.	-1.	4.	0.	0.	0.	-1.	0.	0.	0.	0.]]
[[0.	0.	0.	0.	-1.	0.	0.	0.	4.	-1.	0.	0.	-1.	0.	0.	0.]]
[[0.	0.	0.	0.	0.	-1.	0.	0.	-1.	4.	-1.	0.	0.	-1.	0.	0.]]
[[0.	0.	0.	0.	0.	0.	-1.	0.	0.	-1.	4.	-1.	0.	0.	-1.	0.]]
[[0.	0.	0.	0.	0.	0.	0.	0.	-1.	0.	0.	0.	4.	-1.	0.	0.]]
[[0.	0.	0.	0.	0.	0.	0.	0.	0.	-1.	0.	0.	-1.	4.	-1.	0.]]
[[0.	0.	0.	0.	0.	0.	0.	0.	0.	0.	-1.	0.	0.	-1.	4.	-1.]]
[[0.	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.	-1.	0.	0.	-1.	4.]]



Approximating the solution to the PDE for $f = 1$

$$A_h = h^2 A_h * (1/(h^2))$$

Approximating the solution to the PDE for $f = 1$

$$A_h = h^2 A_h * (1/(h^2))$$

$$b = \text{np.ones}(J^2)$$

Approximating the solution to the PDE for $f = 1$

```
Ah = h2_Ah*(1/(h**2))
```

```
b = np.ones(J**2)
```

```
# We need to add this at the beginning of the file  
# from scipy.sparse.linalg import dsolve:
```

```
u = dsolve.spsolve(Ah, b)
```

Approximating the solution to the PDE for $f = 1$

```

z = np.empty([J+2, J+2])

for i in range(0, J+2):
    for j in range(0, J+2):

        n = j+J*(i-1)    # Going from two indices to one

        if i==0 or i==J+1:
            z[i, j] = 0.0
        if j==0 or j==J+1:
            z[i, j] = 0.0
        if i>0 and j>0 and i<J+1 and j<J+1:
            z[i, j] = u[n-1]
            # elements of u are numbered starting at 0

```

Approximating the solution to the PDE for $f = 1$



```
# We need to add this at the beginning of the file
# import matplotlib.cm as cm
# from matplotlib import pyplot as plt
# from mpl_toolkits.mplot3d import Axes3D

# First way to plot the data: showing values using colors
plt.imshow(z, cmap=cm.jet, extent=[0.0, 1.0, 0.0, 1.0])
plt.colorbar()
plt.show()

# Second way to plot the data: we create a 3D-plot
fig = plt.figure()
ax = fig.gca(projection='3d')
x = np.linspace(0.0, 1.0, num=J+2)
y = np.linspace(0.0, 1.0, num=J+2)
X, Y = np.meshgrid(x, y)
surf = ax.plot_surface(X, Y, z, linewidth=0, cmap=cm.jet)
plt.show()
```



Approximating the solution to the PDE for $f = 1$

$$J = 4$$





Approximating the solution to the PDE for $f = 1$

$$J = 40$$





Approximating the solution to the PDE for $f = 1$

$$J = 400$$



Approximating the solution to the PDE for $f = 1$

$$J = 800$$



Approximating the solution to the PDE for $f = 1$

Program available for download at

<https://cagnol.link/fdmp2>

VII.4. FEM-FDM comparison

FDM	FEM
Consistency error	Céa Lemma
Consistency	interpolation
L^∞ -stability	L^2 -coercivity