

Information theory

Some basic definitions: The case of continuous random variables

Georgios Ropokis

CentraleSupélec, Campus Rennes


Table of contents

1. Continuous random variables
2. Differential Entropy
3. Joint and conditional differential entropy
4. Relative entropy and mutual information
5. Further properties
6. Transformations of random variables
7. Gaussian random vectors and differential entropy

Continuous random variables

Definition of a continuous random variable

- We define a continuous random variable as a mapping between a random experiment and a sample space that is an infinite and uncountable subset of \mathbb{R} .
- Example: For a javelin throw experiment we can define the event that the javelin lands at a point s . The distance $x(s)$ covered by the javelin is then a continuous random variable.



Examples of continuous random variables

- Measurements obtained by an analogue sensor.
- The received signal strength for a wireless receiver, at a random position.
- Thermal noise at an electric circuit.
- The interference reaching a wireless communications receiver.

Characterizing a continuous random variable: Probability density function

- Assigning a non zero probability of occurrence to each one of the possible values of a continuous random variable X does not lead to a probability sum equal to 1.
- Instead, we assign non-zero probabilities for each one of the subintervals of the range of values of random variable X .
- We define the cumulative distribution function $F_X(x)$ as the function that gives us the probability that $X \leq x$, i.e., as:

$$F_X(x) = \Pr(X \leq x). \quad (1)$$

- Probability density function: If the derivative of the cumulative distribution function exists, then we call this derivative $f_X(\cdot)$, the probability density function of X . Using $f_X(x)$ we can calculate $\Pr\{a \leq x \leq b\}$ as:

$$\Pr\{a \leq x \leq b\} = \int_a^b f_X(x) dx. \quad (2)$$

- Both the probability density function and the cumulative distribution function uniquely characterize random variable X .

Properties of probability density functions

Property 1: Non negativity

$$f_X(x) \geq 0, \quad -\infty < x < \infty. \quad (3)$$

Property 2: Integration property

$$\int_{-\infty}^{\infty} f_X(x) dx = 1. \quad (4)$$

Some well known continuous distributions

Uniform distribution

$$f_X(x) = \begin{cases} \frac{1}{b-a}, & a < x < b \\ 0, & \text{otherwise.} \end{cases} \quad (5)$$

Exponential distribution

$$f_X(x) = \begin{cases} \lambda \exp(-\lambda x), & x > 0 \\ 0, & x \leq 0. \end{cases} \quad (6)$$

Gaussian distribution

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right), -\infty < x < \infty. \quad (7)$$

Implicit characterization of random variables

Expectation of a random variable

We define the expectation of a random variable as the integral:

$$\mathbb{E}\{X\} = \int_{-\infty}^{\infty} x f_X(x) dx. \quad (8)$$

Expectation of a function of a random variable

For a random variable $Y = g(X)$, we define the expectation of Y as:

$$\mathbb{E}\{Y\} = \mathbb{E}\{g(X)\} = \int_{-\infty}^{\infty} g(x) f_X(x) dx. \quad (9)$$

Differential Entropy

Definition of Differential Entropy

Definition: Differential entropy

Let X be a continuous random variable having a non-zero probability density function $f_X(x)$ over a support set \mathcal{D} and a cumulative distribution function $F_X(x)$. We define the differential entropy as:

$$h(X) = - \int_{\mathcal{D}} f_X(x) \log(f_X(x)) dx.$$

Remark: Similar to the case of a discrete random variable, the differential entropy is determined by the probability density function of the considered random variable. For this reason, we can also use notation $h(f)$ instead of $h(X)$.

Differential entropy for some known distribution functions

Uniform distribution

Let us consider a random variable uniformly distributed in the interval (a, b) . We can then calculate its differential entropy as:


$$h(X) = -\mathbb{E} \{ \log(f_X(X)) \} = \int_a^b \frac{1}{b-a} \log(b-a) dx = \log(b-a). \quad (10)$$

Remark: Assuming $b - a < 1$, the differential entropy becomes negative. As a result, unlike the case of the entropy of a discrete random variable positivity is not necessary for the differential entropy.

Exponential distribution (Entropy in nats)

$$\begin{aligned} h(X) &= - \int_0^{\infty} \lambda \exp(-\lambda x) \ln(\lambda \exp(-\lambda x)) dx \\ &= -\ln \lambda + \lambda \int_0^{\infty} x \lambda \exp(-\lambda x) dx = 1 - \ln \lambda \end{aligned} \quad (11)$$

Differential entropy of exponential distribution



As λ increases X becomes more and more limited (with respect to the values taken with significant probability) and the entropy decreases.

Differential entropy is still a measure of uncertainty!

Differential entropy for some known distribution functions

Gaussian distribution (Entropy in nats)


We consider a random variable X following a normal distribution of the form:

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right). \quad (12)$$

The differential entropy (in nats) is then given as:

$$\begin{aligned} h(X) &= - \int_{-\infty}^{+\infty} \frac{\exp\left(-\frac{x^2}{2\sigma^2}\right)}{\sqrt{2\pi\sigma^2}} \ln\left(\frac{\exp\left(-\frac{x^2}{2\sigma^2}\right)}{\sqrt{2\pi\sigma^2}}\right) dx \\ &= \frac{\ln 2\pi\sigma^2}{2} + \frac{1}{2\sigma^2} \int_{-\infty}^{+\infty} \frac{x^2 \exp\left(-\frac{x^2}{2\sigma^2}\right)}{\sqrt{2\pi\sigma^2}} dx \\ &= \frac{\ln 2\pi\sigma^2}{2} + \frac{\mathbb{E}\{X^2\}}{2\sigma^2} = \frac{1}{2} \ln 2\pi e \sigma^2 \end{aligned}$$

Differential entropy of a normal random variable



Regardless of the distribution (e.g. Gaussian or exponential), low entropy indicates that the random variable is confined to a small set of values, while high entropy indicates that the random variable is more dispersed.

Connection between differential entropy and discrete entropy

Quantization and entropy

- Let X be a continuous random variable having a probability density function $f_X(x)$
- Let us also consider dividing the range of values of X in n bins, each one of length $\Delta = \frac{1}{2^n}$.
- From the Mean Value Theorem, we have that for the interval $[i\Delta, (i+1)\Delta)$, there exists an $x_i \in [i\Delta, (i+1)\Delta)$ such that $f_X(x_i) \Delta = \int_{i\Delta}^{(i+1)\Delta} f_X(x) dx$
- Question: By introducing the discrete random variable:

$$X_Q = x_i, \text{ if } i\Delta \leq X < (i+1)\Delta,$$

How is the entropy of X_Q connected to $h(X)$?

Connection between differential entropy and discrete entropy

Quantization and entropy

We start by introducing the probabilities

$p_i = \Pr(i\Delta \leq X < (i+1)\Delta) = \int_{i\Delta}^{(i+1)\Delta} f_X(x) dx = f_X(x_i) \Delta$. We can then calculate the entropy of X_Q as:

$$\begin{aligned} H(X_Q) &= \sum_{i=-\infty}^{+\infty} p_i \log p_i = - \sum_{i=-\infty}^{+\infty} f_X(x_i) \Delta \log(f_X(x_i) \Delta) \\ &= - \sum_{i=-\infty}^{+\infty} f_X(x_i) \Delta \log(f_X(x_i)) - \sum_{i=-\infty}^{+\infty} f_X(x_i) \Delta \log(\Delta) \quad (13) \\ &= - \sum_{i=-\infty}^{+\infty} f_X(x_i) \Delta \log(f_X(x_i)) - \log \Delta. \end{aligned}$$

Theorem: If the density $f(x)$ of X is Riemann integrable, then:

$$H(X_Q) + \log \Delta \rightarrow h(X), \text{ as } \Delta \rightarrow 0.$$

Result: For a continuous random variable X , the number of bits to describe X with n -bit accuracy is equal to $h(X) + n$.

Joint and conditional differential entropy

We characterize a random vector $\mathbf{X} = [X_1, \dots, X_N]$ by means of its joint distribution function $F_{\mathbf{X}}(x_1, \dots, x_N)$ defined as:

$$F_{\mathbf{X}}(x_1, \dots, x_N) = \Pr(X_1 \leq x_1, \dots, X_N \leq x_N). \quad (14)$$

We also define the joint probability distribution function as:

$$f_{\mathbf{X}}(x_1, \dots, x_N) = \frac{\partial^N F_{\mathbf{X}}(x_1, \dots, x_N)}{\partial x_1 \cdots \partial x_N} \quad (15)$$

Expectation operators for random vectors

Definition: Expectation of a function of a random vector

Let $g(\mathbf{X}) : \mathbb{R}^N \rightarrow \mathbb{R}$ be a multivariate function of \mathbf{X} . Let $\mathcal{D}_{\mathbf{X}}$ be the domain of support for the joint probability density function of \mathbf{X} . We then define its expectation as the following N -dimensional integral:

$$\mathbb{E}\{g(\mathbf{X})\} = \int_{\mathcal{D}_{\mathbf{X}}} g(x_1, \dots, x_N) f_{\mathbf{X}}(x_1, \dots, x_N) dx_1 \dots x_N \quad (16)$$

Some important moments

Expectation of a vector: $\mathbb{E}\{\mathbf{X}\} = [\mathbb{E}\{X_1\}, \dots, \mathbb{E}\{X_N\}]^T$

Covariance matrix: $\mathbf{\Sigma} = \mathbb{E}\{(\mathbf{X} - \mathbb{E}\{\mathbf{X}\})(\mathbf{X} - \mathbb{E}\{\mathbf{X}\})^T\}$

Correlation matrix: $\mathbf{C} = \mathbb{E}\{\mathbf{X}\mathbf{X}^T\}$.

Conditional distributions and statistics

Definition: Conditional distribution

Assuming knowledge of random variables X_1, \dots, X_k , we define the conditional distribution of x_{k+1}, \dots, x_N as:

$$f_{\mathbf{X}}(x_{k+1}, \dots, x_N | x_1, \dots, x_k) = \frac{f_{\mathbf{X}}(x_1, \dots, x_k, \dots, x_N)}{f_{X_1, \dots, X_k}(x_1, \dots, x_k)}. \quad (17)$$

$$\text{where } f_{X_1, \dots, X_k}(x_1, \dots, x_k) = \int_{\mathcal{D}_{k+1}} \cdots \int_{\mathcal{D}_N} f_{\mathbf{X}}(x_1, \dots, x_N) dx_{k+1} \cdots dx_N. \quad (18)$$

and \mathcal{D}_i is the domain of support of X_i .

Definition: Conditional expectation

Assuming knowledge of random variables X_1, \dots, X_k , we define the conditional expectation of $g(x_1, \dots, x_N)$ as

$$\mathbb{E}_{|X_1, \dots, X_k} \{g(x_1, \dots, x_N)\} = \int_{\mathcal{D}_{k+1}} \cdots \int_{\mathcal{D}_N} g(x_1, \dots, x_N) f_{\mathbf{X}}(x_{k+1}, \dots, x_N | x_1, \dots, x_k) dx_{k+1} \cdots dx_N. \quad (19)$$

The Multivariate Gaussian Distribution and its properties

Definition: Jointly Gaussian random variables & Gaussian vectors

A collection $X_i, i \in I$ of random variables is said to have a joint Gaussian distribution if every linear combination of X_i s is a Gaussian random variable.

A random vector \mathbf{x} is said to be a Gaussian random vector if its elements are jointly Gaussian distributed.

Definition: The multivariate Gaussian distribution

Let \mathbf{X} be a Gaussian random vector, having a mean value $\boldsymbol{\mu}$ and a non-singular covariance matrix \mathbf{K} . The joint distribution function of \mathbf{X} is then defined as

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{n/2} |\mathbf{K}|^{1/2}} \exp \left(-\frac{(\mathbf{x} - \boldsymbol{\mu})^T \mathbf{K}^{-1} (\mathbf{x} - \boldsymbol{\mu})}{2} \right). \quad (20)$$

Definitions of joint and conditional differential entropy

Definition: Differential entropy

Definition: Let X_1, \dots, X_n be a set of n random variables defined on a support set $\mathcal{D}_X \subseteq \mathbb{R}^n$. We define the differential entropy of X_1, \dots, X_n as:

$$h(X_1, \dots, X_n) = - \int_{\mathcal{D}_X} f_X(x_1, \dots, x_n) \log f_X(x_1, \dots, x_n) dx_1 \cdots dx_n. \quad (21)$$

Definition: Conditional differential entropy

Let X and Y be random variables having a joint distribution $f_{X,Y}(x,y)$. We define the conditional differential entropy $h(X|Y)$ as:

$$h(X|Y) = - \int_{\mathcal{D}_X} \int_{\mathcal{D}_Y} f_{X,Y}(x,y) \log f_X(x|y) dx dy. \quad (22)$$

where $\mathcal{D}_X, \mathcal{D}_Y$ are the domains of support of X and Y respectively.

Remark: Given that $f_{X,Y}(x,y) = f_X(x|y) f_Y(y)$, we can equivalently write the conditional differential entropy as:

$$h(X|Y) = h(X, Y) - h(Y). \quad (23)$$

Entropy of Gaussian distribution (in nats)

Theorem: Entropy of a multivariate Gaussian vector (Entropy in nats)

The entropy of a multivariate Gaussian vector with a mean vector μ and covariance matrix \mathbf{K} is written as:

$$h(\mathbf{X}) = \frac{1}{2} \ln(2\pi e)^n |\mathbf{K}|.$$

Proof: Recall that:

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{n/2} |\mathbf{K}|^{1/2}} \exp\left(-\frac{(\mathbf{x} - \mu)^T \mathbf{K}^{-1} (\mathbf{x} - \mu)}{2}\right), \mathbf{x} \in \mathbb{R}^n. \quad (24)$$

We then have that:

$$h(\mathbf{X}) = - \int_{\mathbb{R}^n} f_{\mathbf{X}}(\mathbf{x}) \left(-\frac{1}{2} (\mathbf{x} - \mu)^T \mathbf{K}^{-1} (\mathbf{x} - \mu) - \ln\left((2\pi)^{n/2} |\mathbf{K}|^{1/2}\right) \right) d\mathbf{x} \quad (25)$$

Entropy of Gaussian distribution

Proof continued

$$\begin{aligned}h(\mathbf{X}) &= - \int_{\mathbb{R}^n} f_{\mathbf{X}}(\mathbf{x}) \left(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \mathbf{K}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right) d\mathbf{x} + \frac{1}{2} \ln((2\pi)^n |\mathbf{K}|) \\&= \frac{1}{2} \sum_{j=1}^n \sum_{i=1}^n [\mathbf{K}]_{j,i} [\mathbf{K}^{-1}]_{i,j} + \frac{1}{2} \ln((2\pi)^n |\mathbf{K}|) \\&= \frac{1}{2} \sum_{j=1}^n (\mathbf{K} \mathbf{K}^{-1})_{j,j} + \ln((2\pi)^n |\mathbf{K}|) \\&= \frac{n}{2} + \ln((2\pi)^n |\mathbf{K}|) = \frac{1}{2} \ln((2\pi e)^n |\mathbf{K}|)\end{aligned}\tag{26}$$

Relative entropy and mutual information

Definition:Relative entropy

Given two probability density functions $f_X(\cdot)$ and $z_X(\cdot)$, we define the relative entropy (also called Kullback-Leibler distance) $D(f_X||z_X)$ between the two densities as:

$$D(f_X||z_X) = \int_{\mathcal{D}_X} f_X(x) \log \left(\frac{f_X(x)}{z_X(x)} \right) dx. \quad (27)$$

where \mathcal{D}_X is the domain of support for $f_X(\cdot)$.

Remark: If the domain of support of $f_X(\cdot)$ is not fully contained in the domain of support of $z_X(\cdot)$, then Kullback-Leibler distance is infinite.

Mutual information

Definition: Mutual information

Given two random variables X and Y with a joint density $f_{X,Y}(x,y)$, we define the mutual information as:

$$I(X; Y) = \int_{\mathcal{D}_X} \int_{\mathcal{D}_Y} f_{X,Y}(x,y) \log \left(\frac{f_{X,Y}(x,y)}{f_X(x) f_Y(y)} \right) dx dy. \quad (28)$$

Basic properties

1. $I(X; Y) = h(X) - h(X|Y)$
2. $I(X; Y) = h(Y) - h(Y|X)$
3. $I(X; Y) = h(X) + h(Y) - h(X, Y)$
4. $I(X; Y) = D(f_{X,Y}(x,y) || f_X(x) f_Y(y))$

Further properties

Properties of differential entropy/relative entropy/ mutual information

Theorem: Positivity of relative entropy

Given two densities $f_X(\cdot)$ and $z_X(\cdot)$ it holds that $D(f_X||z_X) \geq 0$ with equality if and only if $f_X = z_X$ almost everywhere.

Proof: Using Jensen inequality and the concavity of the logarithm we have that:

$$\begin{aligned} -D(f_X||z_X) &= \int_{\mathcal{D}_X} f_X(x) \log\left(\frac{z_X(x)}{f_X(x)}\right) dx \leq \log\left(\int_{\mathcal{D}_X} f_X(x) \frac{z_X(x)}{f_X(x)} dx\right) \\ &= \log\left(\int_{\mathcal{D}_X} \log z_X(x) dx\right) \leq \log 1 = 0. \end{aligned} \tag{29}$$

where \mathcal{D}_X is the domain of support of $f_X(\cdot)$. Moreover, we note that equality can only be satisfied if the Jensen inequality is satisfied with equality. This only occurs if $f(x) = g(x)$ almost everywhere.

What does the above theorem imply?

Consequences of the theorem

- $I(X; Y) \geq 0$ where equality holds if and only if X and Y are independent.
- $h(X|Y) \leq h(X)$ where equality holds if and only if X and Y are independent.

Transformations of random variables

Translation and scaling of random variables

Theorem: Differential entropy and invariance to translations

$$h(X + c) = h(X) \quad (30)$$

Proof: This result is a direct consequence of the definition of differential entropy.

Theorem: Differential entropy and scaling

$$h(aX) = h(X) + \log |a| \quad (31)$$

Proof: Use the fact that $f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y}{a}\right)$, as well as the definition of differential entropy.

Remark: As $|a|$ increases, the uncertainty increases.

Gaussian random vectors and differential entropy

Maximizing the differential entropy

Theorem: Upper bound on the differential entropy

Let $\mathbf{X} \in \mathbb{R}^n$ be a zero mean random vector having a covariance matrix $\mathbf{K} = \mathbb{E}\{\mathbf{X}\mathbf{X}^T\}$. It then holds that $h(\mathbf{X}) \leq \frac{1}{2} \log((2\pi e)^n |\mathbf{K}|)$ where equality holds if and only if $\mathbf{X} \sim \mathcal{N}(\mathbf{0}, \mathbf{K})$.

Proof: Let us start by using notation $\phi_{\mathbf{K}}(\cdot)$ for the probability density function of a vector distributed according to $\mathcal{N}(\mathbf{0}, \mathbf{K})$. It is easy then to see that the logarithm of $\phi_{\mathbf{K}}(\mathbf{x})$ is a quadratic form and that $\int x_i x_j \phi_{\mathbf{K}}(\mathbf{x}) d\mathbf{x} = [\mathbf{K}]_{i,j}$. We now use the non negativity property for the differential entropy between any distribution $g(\cdot)$ having a covariance matrix \mathbf{K} and a Gaussian distribution having the same covariance matrix, and obtain that:

$$\begin{aligned} 0 \leq D(g||\phi_{\mathbf{K}}) &= \int_{\mathbb{R}^n} g(\mathbf{x}) \log(g(\mathbf{x})/\phi_{\mathbf{K}}(\mathbf{x})) d\mathbf{x} \\ &= -h(g) - \int_{\mathbb{R}^n} g(\mathbf{x}) \log \phi_{\mathbf{K}}(\mathbf{x}) d\mathbf{x} \\ &= -h(g) - \int \phi_{\mathbf{K}}(\mathbf{x}) \log \phi_{\mathbf{K}}(\mathbf{x}) = -h(g) + h(\phi_{\mathbf{K}}) \end{aligned} \tag{32}$$

Note: In our proof we have used the fact that term $\int g \log \phi_{\mathbf{K}} = \int \phi_{\mathbf{K}} \log \phi_{\mathbf{K}}$, since $\phi_{\mathbf{K}}$ is a quadratic form, it is solely determined by the covariance matrix.