

A WKB-based fixed-grid method for capturing trait concentration in a dispersal evolution model

Weijie Huang* and Xinran Ruan†

Abstract

The evolution of dispersal traits is a fundamental topic in evolutionary ecology, where natural selection may drive the trait distribution toward concentration in the rare mutation regime. This singular behavior poses a serious numerical difficulty, since direct discretizations of the population density require very fine trait grids to resolve the narrow peak and to identify the fittest trait accurately. In this paper, we develop a WKB-based numerical framework for a dispersal evolution model. By separating the exponentially concentrated trait profile from a smoother amplitude variable, the method captures the selected trait on fixed trait grids without resolving the full density peak. We formulate the semi-discrete WKB scheme, prove positivity of the amplitude, and establish a stationary fixed-grid asymptotic-preserving structure, which explains how the selected trait can be recovered without resolving the full concentrated density profile. We further discuss the fully discrete realization and the time-step restrictions associated with the time-dependent effective Hamiltonian. Numerical experiments compare the proposed WKB formulation with direct density discretizations and show its advantage in locating the fittest trait in the small-mutation regime.

Key words. dispersal evolution, trait concentration, WKB ansatz, asymptotic-preserving method, Hamilton–Jacobi equation, principal eigenvalue.

1 Introduction

Adaptive dynamics studies the evolution of phenotypic traits under mutation, selection, and ecological feedback. In structured population models, a central mathematical manifestation of natural selection is the concentration of the population density on one or several fittest traits. In the rare-mutation regime this concentration is often described by a Dirac mass in the phenotypic variable, and the support of the limiting measure represents the selected phenotype. Numerically, this creates a singular-resolution problem. When the mutation parameter is small, the solution develops a very narrow trait profile whose location is determined by the interaction between spatial heterogeneity, trait-dependent dispersal, and competition. A direct discretization of the original density then requires a prohibitively fine mesh in the trait direction. The main computational issue is therefore not only to approximate the full density, but also to locate the emerging concentration and identify the fittest trait efficiently and robustly.

In this work, we consider a representative dispersal evolution model of the form [?, ?]

$$\varepsilon \partial_t n_\varepsilon - D(\theta) \Delta_{\mathbf{x}} n_\varepsilon - \varepsilon^2 \Delta_\theta n_\varepsilon = n_\varepsilon (K(\mathbf{x}) - \rho_\varepsilon(\mathbf{x})), \quad \mathbf{x} \in \Omega, \quad (1.1)$$

*School of Mathematics and Statistics, Beijing Jiaotong University, Beijing 100044, People’s Republic of China (wjhuang@bjtu.edu.cn)

†School of Mathematical Sciences, Capital Normal University, Beijing, China, 100048, People’s Republic of China (xinran.ruan@cnu.edu.cn).

where t denotes time, $\mathbf{x}$ the spatial variable, and θ a phenotypic variable. The unknown $n_\varepsilon(t, \mathbf{x}, \theta)$ is the structured population density, and

$$\rho_\varepsilon(t, \mathbf{x}) = \int_{\mathbb{T}} n_\varepsilon(t, \mathbf{x}, \theta) d\theta$$

is the total population density. The function $K(\mathbf{x})$ represents the local carrying capacity, while the trait-dependent diffusivity $D(\theta)$ models the dispersal rate associated with phenotype θ . Throughout the paper, we assume that the dispersal rate is positive and periodic in the trait variable and that the carrying capacity is nonnegative and bounded:

$$0 < D_{\min} \leq D(\theta) \leq D_{\max} < \infty, \quad 0 \leq K(\mathbf{x}) \leq K_{\max}. \quad (1.2)$$

We impose homogeneous Neumann boundary conditions on $\partial\Omega$ and periodic boundary conditions in the θ -direction. Although our numerical development is carried out for (1.1), the main difficulty addressed here is more general and arises in a broader class of small-mutation trait-selection problems whose solutions become highly concentrated in the phenotypic direction.

Model (1.1) and related equations have been extensively studied from the analytical viewpoint, including well-posedness, long-time behavior, and the connection with adaptive dynamics [3, 4, 1]. A characteristic feature is that the competition between spatial heterogeneity and trait-dependent dispersal may select an optimal phenotype in the long-time regime. In many situations, the steady state exhibits concentration in the trait variable. The small parameter $\varepsilon > 0$ plays the role of a rare mutation scale, and in the limit $\varepsilon \rightarrow 0$ the steady state typically converges, in a weak sense, to a Dirac mass supported at the fittest trait. This singular limit is well understood at the PDE level through WKB or Hopf–Cole transforms and constrained Hamilton–Jacobi equations coupled with principal eigenvalue problems [4, 2, 3].

From the numerical point of view, however, this asymptotic regime remains challenging. As ε decreases, the trait profile of n_ε becomes increasingly narrow, so standard discretizations of the original variable require a rapidly growing number of grid points in the phenotypic direction. Moreover, approximating the full density with reasonable accuracy does not automatically guarantee an accurate identification of the dominant trait. This is particularly restrictive in the small- ε regime, where the quantity of real interest is often the location of the concentration and its effective profile rather than a pointwise resolution of the entire density on an extremely fine mesh.

Motivated by the asymptotic structure of the rare-mutation limit, we adopt the WKB representation

$$n_\varepsilon(\mathbf{x}, \theta, t) = W_\varepsilon(\mathbf{x}, \theta, t) \exp\left(\frac{u_\varepsilon(\theta, t)}{\varepsilon}\right),$$

where the phase variable u_ε describes the rapidly varying exponential profile in the trait direction, while W_ε is a comparatively smoother amplitude. In the long-time and rare-mutation regime, the effective behavior of the phase variable is related to a constrained Hamilton–Jacobi equation involving a principal eigenvalue problem in the spatial variable [4, 1]. This observation suggests that the WKB reformulation is not only asymptotically natural but also computationally advantageous. In particular, it provides a mechanism for extracting the dominant trait from the phase variable without directly resolving the narrow peak of the original density on a prohibitively fine trait grid.

The present paper is organized around this computational viewpoint. Our main goal is to develop and study a WKB-based numerical strategy that is able to capture trait concentration more efficiently than direct discretizations of the original model. To this end, we first derive a semi-discrete formulation for the phase and amplitude variables. We then specify a concrete numerical method based on a monotone discretization of the Hamilton–Jacobi part and a positivity-preserving

discretization of the amplitude equation. At the analytical level, we establish positivity of the amplitude and derive a stationary, fixed-grid AP structure under transparent reconstruction and one-sided remainder stability assumptions. Since the WKB split does not automatically provide an exact discrete chain rule, we also record a density-compatible variant that restores a conservative density-level balance and makes the weighted remainder estimate verifiable. On the numerical side, our experiments are designed to compare the original discretization and the WKB formulation directly, to track the remainder diagnostic, and to test the accurate capture of the fittest trait when ε is small.

The paper is organized as follows. In Section 2 we recall the WKB ansatz and the formal steady-state rare mutation limit that motivate the numerical formulation. In Section 3 we present the semi-discrete scheme, prove positivity, and derive the conditional stationary AP structure, including the density-compatible variant that controls the weighted remainder. **We then add a separate subsection on the fully discrete realization, where the phase update, amplitude update, residuals, time-step restrictions, and projective stabilization are stated together.** Section 4 is devoted to numerical experiments, where we compare the WKB formulation with direct discretizations, examine its performance in the concentration regime, and monitor the remainder diagnostic. The appendices collect typical choices of monotone numerical Hamiltonians and monotone numerical fluxes, density reconstruction details, and sufficient conditions for the one-sided remainder bound.

2 Preliminary

2.1 WKB ansatz and reformulated system

To capture the concentration phenomenon in the trait variable, we introduce the classical WKB ansatz

$$n_\varepsilon(x, \theta, t) = W_\varepsilon(x, \theta, t) \exp\left(\frac{u_\varepsilon(\theta, t)}{\varepsilon}\right). \quad (2.3)$$

This decomposition separates the exponentially varying part in the trait direction from a comparatively smoother amplitude. It is therefore natural for both the continuous asymptotic analysis and the numerical design in the small-mutation regime.

Substituting (2.3) into (1.1) and assigning the leading trait-direction exponential contributions to the phase function, we obtain the reformulated system

$$\begin{cases} \partial_t u_\varepsilon - |\partial_\theta u_\varepsilon|^2 - \varepsilon \partial_{\theta\theta} u_\varepsilon = -\mathcal{H}(\theta, t), \\ \varepsilon \partial_t W_\varepsilon - D(\theta) \Delta_x W_\varepsilon - \varepsilon^2 \partial_{\theta\theta} W_\varepsilon - 2\varepsilon \partial_\theta W_\varepsilon \partial_\theta u_\varepsilon = W_\varepsilon (K(x) - \rho_\varepsilon(x, t) + \mathcal{H}(\theta, t)), \end{cases} \quad (2.4)$$

where $\mathcal{H}(\theta, t)$ denotes an effective Hamiltonian. Using the identity

$$\partial_\theta (W_\varepsilon \partial_\theta u_\varepsilon) = \partial_\theta W_\varepsilon \partial_\theta u_\varepsilon + W_\varepsilon \partial_{\theta\theta} u_\varepsilon,$$

we rewrite the transport coupling in conservative form. If

$$F_\varepsilon = -W_\varepsilon \partial_\theta u_\varepsilon, \quad (2.5)$$

then

$$-2\varepsilon \partial_\theta W_\varepsilon \partial_\theta u_\varepsilon = 2\varepsilon \partial_\theta F_\varepsilon + 2\varepsilon W_\varepsilon \partial_{\theta\theta} u_\varepsilon.$$

Consequently, the system can be written as

$$\begin{cases} \partial_t u_\varepsilon - |\partial_\theta u_\varepsilon|^2 - \varepsilon \partial_{\theta\theta} u_\varepsilon = -\mathcal{H}(\theta, t), \\ \varepsilon \partial_t W_\varepsilon - D(\theta) \Delta_x W_\varepsilon - \varepsilon^2 \partial_{\theta\theta} W_\varepsilon + 2\varepsilon \partial_\theta F_\varepsilon = W_\varepsilon (K(x) - \rho_\varepsilon(x, t) + \mathcal{H}(\theta, t) - 2\varepsilon \partial_{\theta\theta} u_\varepsilon). \end{cases} \quad (2.6)$$

In the continuous theory, $\mathcal{H}$ is related to a principal eigenvalue problem in the spatial variable. This structure is the basic reason for introducing the variables $(u_\varepsilon, W_\varepsilon)$ in the first place.

2.2 Continuous asymptotic structure of the steady state

We next recall the steady-state asymptotic structure that motivates the numerical method. Let $(\bar{n}_\varepsilon, \bar{\rho}_\varepsilon)$ be a positive steady state of (1.1). Then

$$-D(\theta)\Delta_x \bar{n}_\varepsilon - \varepsilon^2 \partial_{\theta\theta} \bar{n}_\varepsilon = \bar{n}_\varepsilon (K(x) - \bar{\rho}_\varepsilon(x)), \quad \bar{\rho}_\varepsilon(x) = \int_{\mathbb{T}} \bar{n}_\varepsilon(x, \theta) d\theta. \quad (2.7)$$

As shown in [4], under standard assumptions on K , D , and the domain, the steady-state family admits several properties that are fundamental for the rare-mutation limit.

Proposition 2.1 (Continuous asymptotic structure of the steady state). *Assume that K is positive and nonconstant, that D is positive, periodic, and has a unique minimizer θ_m , and that the steady problem (2.7) admits a positive solution. Then the following properties hold at the continuous level.*

1. *There exists a constant $C > 0$, independent of ε , such that*

$$0 \leq \bar{\rho}_\varepsilon(x) \leq C \quad \text{for all } x \in \Omega. \quad (2.8)$$

Moreover, up to subsequences, $\bar{\rho}_\varepsilon$ converges uniformly to a limit density ρ .

2. *If we write*

$$\bar{n}_\varepsilon(x, \theta) = \exp\left(\frac{\bar{u}_\varepsilon(x, \theta)}{\varepsilon}\right), \quad (2.9)$$

then $\bar{u}_\varepsilon$ is uniformly Lipschitz continuous in θ , and along subsequences

$$\bar{u}_\varepsilon \rightarrow u \quad \text{uniformly in } (x, \theta), \quad (2.10)$$

where the limit u is independent of x , periodic in θ , and satisfies

$$\begin{cases} |\partial_\theta u(\theta)|^2 = \mathcal{H}(\theta, \rho(\cdot)), \\ \max_{\theta \in \mathbb{T}} u(\theta) = 0. \end{cases} \quad (2.11)$$

3. *The effective Hamiltonian $\mathcal{H}(\theta, \rho)$ is defined through the principal eigenvalue problem*

$$\begin{cases} -D(\theta)\Delta_x \mathcal{N}(x, \theta) = \mathcal{N}(x, \theta)(K(x) - \rho(x)) + \mathcal{N}(x, \theta) \mathcal{H}(\theta, \rho(\cdot)), & x \in \Omega, \\ \partial_\nu \mathcal{N} = 0, & x \in \partial\Omega, \end{cases} \quad (2.12)$$

with a positive eigenfunction $\mathcal{N}$. Moreover, $\mathcal{H}$ has the same monotonicity in θ as D , and hence

$$\min_{\theta \in \mathbb{T}} \mathcal{H}(\theta, \rho(\cdot)) = \mathcal{H}(\theta_m, \rho(\cdot)) = 0. \quad (2.13)$$

4. *The steady state concentrates on the fittest trait in the limit $\varepsilon \rightarrow 0$. More precisely,*

$$\bar{n}_\varepsilon \rightharpoonup N_m(x) \delta(\theta - \theta_m), \quad \bar{\rho}_\varepsilon \rightarrow N_m(x), \quad (2.14)$$

where N_m solves the reduced Fisher-type problem

$$\begin{cases} -D(\theta_m)\Delta_x N_m = N_m(K(x) - N_m), & x \in \Omega, \\ \partial_\nu N_m = 0, & x \in \partial\Omega. \end{cases} \quad (2.15)$$

Proposition 2.1 provides the continuous target of the numerical scheme. The boundedness of $\bar{\rho}_\varepsilon$ keeps the effective Hamiltonian in a bounded regime. The constrained Hamilton–Jacobi equation identifies the dominant trait through the maximizer of u . The concentration result (2.14) indicates that an asymptotic-preserving discretization should recover the reduced pair (N_m, θ_m) without resolving the original density on an extremely fine trait grid. This is the sense in which the WKB reformulation is used in the sequel.

3 WKB reformulated numerical scheme

In this section, we develop and analyze a semi-discrete numerical scheme for the WKB reformulation (2.6). The scheme is formulated in terms of the phase variable u_ε and the amplitude variable W_ε , with the total density n_ε reconstructed from the WKB representation.

3.1 Semi-discrete numerical discretization

We work at the semi-discrete level in order to separate the discretization in (x, θ) from the additional issue of time stepping. For clarity, we present the scheme in one spatial dimension with uniform grids in the x - and θ -directions. This choice is not essential for the WKB decomposition, but it keeps the notation transparent.

Let $\{x_j\}_{j=1}^J$ be a uniform grid on $\Omega = (a, b)$ and let $\{\theta_k\}_{k=1}^K$ be a uniform periodic grid on $\mathbb{T} = (0, 1)$, with mesh size $\Delta\theta$. We denote

$$W_{j,k}^\varepsilon(t) \approx W_\varepsilon(t, x_j, \theta_k), \quad u_k^\varepsilon(t) \approx u_\varepsilon(t, \theta_k).$$

For notational convenience, we write

$$D_k = D(\theta_k), \quad K_j = K(x_j).$$

By (1.2), we have

$$0 < D_{\min} \leq D_k \leq D_{\max} < \infty, \quad 0 \leq K_j \leq K_{\max}.$$

For a grid function $V = \{V_{j,k}\}$, we use the standard second-order difference operators

$$\delta_x^2 V_{j,k} = \frac{V_{j+1,k} - 2V_{j,k} + V_{j-1,k}}{(\Delta x)^2}, \quad \delta_\theta^2 V_{j,k} = \frac{V_{j,k+1} - 2V_{j,k} + V_{j,k-1}}{(\Delta\theta)^2}.$$

The homogeneous Neumann boundary condition in x is imposed by the ghost values

$$V_{0,k} = V_{1,k}, \quad V_{J+1,k} = V_{J,k},$$

while all indices in the θ -direction are understood periodically. For the phase variable, we also use

$$D_\theta^- u_k = \frac{u_k - u_{k-1}}{\Delta\theta}, \quad D_\theta^+ u_k = \frac{u_{k+1} - u_k}{\Delta\theta}.$$

The semi-discrete WKB system involves two numerical ingredients that are not contained in the standard second-order difference operators introduced above. The first one is the numerical Hamiltonian for the phase equation. The Hamilton–Jacobi term $-|\partial_\theta u_\varepsilon|^2$ is approximated by

$$\widehat{H}(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon).$$

We assume that $\widehat{H}$ is locally Lipschitz and consistent with the continuous Hamiltonian, namely

$$\widehat{H}(p, p) = -p^2.$$

We also assume that it is monotone in the sense that it is nondecreasing with respect to its first argument and nonincreasing with respect to its second argument. When $\widehat{H}$ is differentiable, this condition reads

$$\partial_a \widehat{H}(a, b) \geq 0, \quad \partial_b \widehat{H}(a, b) \leq 0.$$

Typical examples are the Godunov and Lax–Friedrichs numerical Hamiltonians, as recalled in Appendix A.

The second ingredient is the conservative discretization of the transport coupling in the amplitude equation. Recall that the continuous flux is

$$F_\varepsilon = -W_\varepsilon \partial_\theta u_\varepsilon.$$

We denote by

$$\mathcal{D}_\theta \widehat{\mathcal{F}}(W^\varepsilon, u^\varepsilon)_{j,k}$$

a conservative discretization of $\partial_\theta F_\varepsilon$ at (x_j, θ_k) . Here $\widehat{\mathcal{F}}$ approximates the flux $-W_\varepsilon \partial_\theta u_\varepsilon$. We assume that $\mathcal{D}_\theta \widehat{\mathcal{F}}$ is consistent with

$$\partial_\theta F_\varepsilon = \partial_\theta (-W_\varepsilon \partial_\theta u_\varepsilon)$$

when evaluated on smooth functions, and that it is conservative in the periodic trait variable. For the positivity argument below, we further assume the quasi-positivity property

$$W_{j,k} = 0, \quad W_{j,\ell} \geq 0 \text{ for all } \ell \implies -\mathcal{D}_\theta \widehat{\mathcal{F}}(W, u)_{j,k} \geq 0. \quad (3.16)$$

This property is satisfied by standard monotone finite volume fluxes, such as the upwind and Lax–Friedrichs fluxes listed in Appendix B.

With these two ingredients specified, we now define the discrete density and the semi-discrete WKB system. For notational convenience, set

$$E_k^\varepsilon(t) = \exp\left(\frac{u_k^\varepsilon(t)}{\varepsilon}\right). \quad (3.17)$$

The reconstructed nodal density is defined by

$$n_{j,k}^\varepsilon(t) = W_{j,k}^\varepsilon(t) E_k^\varepsilon(t). \quad (3.18)$$

As we shall see later, the evolution of u_ε and W_ε couples with the density function. Here we consider a general density computed by a trait-direction quadrature operator

$$\mathcal{Q}_\theta^\varepsilon[W_j^\varepsilon, u^\varepsilon],$$

which approximates

$$\int_{\mathbb{T}} n_\varepsilon(t, x_j, \theta) d\theta = \int_{\mathbb{T}} W_\varepsilon(t, x_j, \theta) \exp\left(\frac{u_\varepsilon(t, \theta)}{\varepsilon}\right) d\theta.$$

We define

$$\rho_{j,Q}^\varepsilon(t) = \mathcal{Q}_\theta^\varepsilon[W_{j,\cdot}^\varepsilon(t), u^\varepsilon(t)], \quad j = 1, \dots, J. \quad (3.19)$$

A direct composite quadrature gives

$$\rho_{j,Q}^\varepsilon(t) = m_j^\varepsilon(t) := \Delta\theta \sum_{k=1}^K W_{j,k}^\varepsilon(t) E_k^\varepsilon(t), \quad (3.20)$$

whereas in the small- ε regime one may instead use a Laplace-type reconstruction presented in Appendix C near the maximizer of u^ε .

Given

$$\boldsymbol{\rho}_Q(t) = (\rho_{1,Q}^\varepsilon(t), \dots, \rho_{J,Q}^\varepsilon(t)),$$

the discrete effective Hamiltonian $\mathcal{H}_k(\boldsymbol{\rho}_Q(t))$ is determined by the discrete principal eigenvalue problem

$$-D_k \delta_x^2 \mathcal{N}_j^{(k)} = \mathcal{N}_j^{(k)} (K_j - \rho_{j,Q}^\varepsilon(t)) + \mathcal{N}_j^{(k)} \mathcal{H}_k(\boldsymbol{\rho}_Q(t)), \quad j = 1, \dots, J. \quad (3.21)$$

Here $\mathcal{N}^{(k)}$ is the corresponding positive eigenvector, equipped with the same discrete Neumann boundary condition in x . Its normalization does not affect $\mathcal{H}_k$. For definiteness, one may impose

$$\Delta x \sum_{j=1}^J \mathcal{N}_j^{(k)} = 1.$$

With the above notation, the semi-discrete WKB scheme reads

$$\begin{cases} \frac{d}{dt} u_k^\varepsilon + \widehat{H}(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon) - \varepsilon \delta_\theta^2 u_k^\varepsilon = -\mathcal{H}_k(\boldsymbol{\rho}_Q(t)), & k = 1, \dots, K, \\ \varepsilon \frac{d}{dt} W_{j,k}^\varepsilon - D_k \delta_x^2 W_{j,k}^\varepsilon - \varepsilon^2 \delta_\theta^2 W_{j,k}^\varepsilon + 2\varepsilon \mathcal{D}_\theta \widehat{\mathcal{F}}(W^\varepsilon, u^\varepsilon)_{j,k} \\ = W_{j,k}^\varepsilon (K_j - \rho_{j,Q}^\varepsilon(t) + \mathcal{H}_k(\boldsymbol{\rho}_Q(t)) - 2\varepsilon \delta_\theta^2 u_k^\varepsilon), & j = 1, \dots, J, \quad k = 1, \dots, K. \end{cases} \quad (3.22)$$

We next show the equation satisfied by the reconstructed density $n_{j,k}^\varepsilon$. Since E_k^ε is independent of the spatial index j ,

$$\delta_x^2 n_{j,k}^\varepsilon = E_k^\varepsilon \delta_x^2 W_{j,k}^\varepsilon.$$

Moreover,

$$\varepsilon \frac{d}{dt} n_{j,k}^\varepsilon = E_k^\varepsilon \left(\varepsilon \frac{d}{dt} W_{j,k}^\varepsilon + W_{j,k}^\varepsilon \frac{d}{dt} u_k^\varepsilon \right). \quad (3.23)$$

Multiplying the first equation in (3.22) by $E_k^\varepsilon W_{j,k}^\varepsilon$ and multiplying the second equation in (3.22) by $\varepsilon E_k^\varepsilon$, noticing (3.23), we can derive that $n_{j,k}^\varepsilon$ satisfies

$$\varepsilon \frac{d}{dt} n_{j,k}^\varepsilon - D_k \delta_x^2 n_{j,k}^\varepsilon = (K_j - \rho_{j,Q}^\varepsilon) n_{j,k}^\varepsilon + R_{j,k}^\varepsilon, \quad (3.24)$$

where the remainder has the form

$$R_{j,k}^\varepsilon = \varepsilon^2 E_k^\varepsilon \delta_\theta^2 W_{j,k}^\varepsilon - \varepsilon W_{j,k}^\varepsilon E_k^\varepsilon \delta_\theta^2 u_k^\varepsilon - W_{j,k}^\varepsilon E_k^\varepsilon \widehat{H}(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon) - 2\varepsilon E_k^\varepsilon \mathcal{D}_\theta \widehat{\mathcal{F}}(W^\varepsilon, u^\varepsilon)_{j,k}. \quad (3.25)$$

Thus the reconstructed density satisfies a semi-discrete analogue of the original density equation, with $R_{j,k}^\varepsilon$ collecting the trait-direction discretization effects. When the numerical Hamiltonian and the conservative flux are evaluated on smooth exact functions, this remainder is consistent with the trait diffusion contribution $\varepsilon^2 \partial_{\theta\theta} n^\varepsilon$.

A density-compatible variant.

The split form above is convenient because it keeps the Hamilton–Jacobi and transport components explicit. For the stationary density estimate, however, it is useful to record an alternative discretization in which the reconstructed density satisfies a conservative density equation exactly. Keeping the same phase equation, replace the amplitude equation by

$$\begin{aligned} \varepsilon \frac{d}{dt} W_{j,k}^\varepsilon - D_k \delta_x^2 W_{j,k}^\varepsilon - \varepsilon^2 (E_k^\varepsilon)^{-1} \delta_\theta^2 (W_{j,\cdot}^\varepsilon, E_\cdot^\varepsilon)_k \\ = W_{j,k}^\varepsilon (K_j - \rho_{j,Q}^\varepsilon + \mathcal{H}_k(\rho_Q) + \widehat{H}(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon) - \varepsilon \delta_\theta^2 u_k^\varepsilon), \end{aligned} \quad (3.26)$$

where

$$\delta_\theta^2 (W_{j,\cdot}^\varepsilon, E_\cdot^\varepsilon)_k = \frac{W_{j,k+1}^\varepsilon E_{k+1}^\varepsilon - 2W_{j,k}^\varepsilon E_k^\varepsilon + W_{j,k-1}^\varepsilon E_{k-1}^\varepsilon}{(\Delta\theta)^2}.$$

Multiplying (3.26) by E_k^ε and using the phase equation gives

$$\varepsilon \frac{d}{dt} n_{j,k}^\varepsilon - D_k \delta_x^2 n_{j,k}^\varepsilon - \varepsilon^2 \delta_\theta^2 n_{j,k}^\varepsilon = (K_j - \rho_{j,Q}^\varepsilon) n_{j,k}^\varepsilon. \quad (3.27)$$

Thus, for this density-compatible variant, the remainder in (3.24) is exactly $R_{j,k}^\varepsilon = \varepsilon^2 \delta_\theta^2 n_{j,k}^\varepsilon$. Consequently, the weighted remainder can be controlled by periodic discrete summation by parts. This observation gives a concrete way to avoid the conditional remainder assumption used below; see Appendix D.

3.2 Basic structural property

The semi-discrete scheme (3.24) can be shown to be positive-preserving as long as the initial data is nonnegative. The detailed result is summarized as the following lemma.

Lemma 3.1 (Positivity of the amplitude). *Assume that $W_{j,k}^\varepsilon(0) \geq 0$ for all j, k . Then the semi-discrete amplitude equation in (3.22) preserves nonnegativity, namely*

$$W_{j,k}^\varepsilon(t) \geq 0 \quad \forall j, k, t \geq 0,$$

as long as the solution exists. Consequently,

$$n_{j,k}^\varepsilon(t) \geq 0 \quad \forall j, k, t \geq 0.$$

Proof. It is enough to verify the quasi-positivity of the right-hand side of the finite-dimensional ODE system for W^ε . From the amplitude equation in (3.22), we can write

$$\varepsilon \frac{d}{dt} W_{j,k}^\varepsilon = D_k \delta_x^2 W_{j,k}^\varepsilon + \varepsilon^2 \delta_\theta^2 W_{j,k}^\varepsilon - 2\varepsilon \mathcal{D}_\theta \widehat{\mathcal{F}}(W^\varepsilon, u^\varepsilon)_{j,k} + W_{j,k}^\varepsilon B_{j,k}(t),$$

where

$$B_{j,k}(t) = K_j - \rho_{j,Q}^\varepsilon(t) + \mathcal{H}_k(\rho_Q(t)) - 2\varepsilon \delta_\theta^2 u_k^\varepsilon(t).$$

Let $W^\varepsilon \geq 0$ and suppose that $W_{j,k}^\varepsilon = 0$ for some (j, k) . Then, for an interior spatial point,

$$\delta_x^2 W_{j,k}^\varepsilon = \frac{W_{j+1,k}^\varepsilon + W_{j-1,k}^\varepsilon}{(\Delta x)^2} \geq 0, \quad \delta_\theta^2 W_{j,k}^\varepsilon = \frac{W_{j,k+1}^\varepsilon + W_{j,k-1}^\varepsilon}{(\Delta\theta)^2} \geq 0.$$

The same conclusion holds at the boundary because of the Neumann ghost values in x -direction and the periodicity in θ -direction. The quasi-positivity assumption on the conservative flux operator implies that

$$-\mathcal{D}_\theta \widehat{\mathcal{F}}(W^\varepsilon, u^\varepsilon)_{j,k} \geq 0.$$

Combining all the inequalities, noticing the fact that $W_{j,k}^\varepsilon B_{j,k}(t) = 0$ since we assumed $W_{j,k}^\varepsilon = 0$, we finally have that, at this specific (j, k) , the following inequality holds,

$$\left. \frac{d}{dt} W_{j,k}^\varepsilon \right|_{W_{j,k}^\varepsilon = 0} \geq 0.$$

Thus the vector field points inward on the boundary of the nonnegative cone. The standard invariance criterion for finite-dimensional ODE systems implies the claim. $\square$

3.3 Stationary AP structure under a one-sided stability condition

We now turn to stationary semi-discrete states. The goal is to identify which parts of the continuous rare-mutation structure are retained on a fixed grid. For the generic split WKB scheme, the only nonstandard point is the trait-direction remainder in the reconstructed density equation. We therefore state the stationary result under an explicit one-sided stability condition. This keeps the AP statement conditional and avoids claiming that the WKB remainder is automatically controlled for every trait discretization. A verifiable density-compatible variant is discussed in Appendix D.

By Lemma 3.1, stationary states obtained from nonnegative amplitudes satisfy

$$W_{j,k}^\varepsilon \geq 0, \quad n_{j,k}^\varepsilon = W_{j,k}^\varepsilon E_k^\varepsilon \geq 0.$$

All estimates below are for such nonnegative reconstructed densities.

Weighted density balance. Define

$$\eta_j^\varepsilon = \Delta\theta \sum_{k=1}^K \frac{n_{j,k}^\varepsilon}{D_k}.$$

Multiplying (3.24) by $1/D_k$ and summing over the trait variable gives the stationary weighted balance equation

$$-\delta_x^2 m_j^\varepsilon = \eta_j^\varepsilon (K_j - \rho_{j,Q}^\varepsilon) + \mathcal{R}_j^{\varepsilon,D}, \quad (3.28)$$

where

$$m_j^\varepsilon = \Delta\theta \sum_{k=1}^K n_{j,k}^\varepsilon, \quad \mathcal{R}_j^{\varepsilon,D} = \Delta\theta \sum_{k=1}^K \frac{R_{j,k}^\varepsilon}{D_k}.$$

The term $\mathcal{R}_j^{\varepsilon,D}$ is the discrete analogue of the continuous weighted mutation contribution

$$\varepsilon^2 \int_{\mathbb{T}} \frac{1}{D(\theta)} \partial_{\theta\theta} n^\varepsilon d\theta = \varepsilon^2 \int_{\mathbb{T}} n^\varepsilon \partial_{\theta\theta} \left(\frac{1}{D(\theta)} \right) d\theta.$$

At the continuous level this is bounded above by $C\varepsilon^2\rho^\varepsilon$. For the split WKB discretization, however, the discrete operators act on the reconstructed density $n_{j,k}^\varepsilon = W_{j,k}^\varepsilon E_k^\varepsilon$, and an exact chain rule is not available. We isolate the needed property as follows.

Assumption 3.1 (One-sided weighted remainder stability). *There exist constants $C_R > 0$ and $r_h \geq 0$, independent of ε , such that every stationary state under consideration satisfies*

$$(\mathcal{R}_j^{\varepsilon,D})_+ \leq C_R m_j^\varepsilon + r_h, \quad j = 1, \dots, J. \quad (3.29)$$

This condition only controls the positive part of the weighted defect; a negative contribution is favorable in the maximum-principle estimate. It is a structural stability requirement on the trait discretization, not a consequence of the preceding consistency statement. For a direct density-level conservative discretization, and for the density-compatible WKB variant (3.26), the same type of estimate follows by discrete summation by parts. For the generic split WKB scheme, it should be checked analytically under additional structure or monitored numerically.

We also record the mild stability condition needed for the density reconstruction in the nonlocal reaction term.

Assumption 3.2 (Reconstruction stability). *There exist constants $c_Q, C_Q > 0$ and $e_Q \geq 0$, independent of ε , such that*

$$\rho_{j,Q}^\varepsilon \geq c_Q m_j^\varepsilon - e_Q, \quad j = 1, \dots, J, \quad (3.30)$$

$$\rho_{j,Q}^\varepsilon \leq C_Q m_j^\varepsilon + e_Q, \quad j = 1, \dots, J. \quad (3.31)$$

For the direct composite quadrature $\rho_{j,Q}^\varepsilon = m_j^\varepsilon$, Assumption 3.2 holds with $c_Q = C_Q = 1$ and $e_Q = 0$.

Proposition 3.1 (Conditional density bound for stationary semi-discrete states). *Let $(u^\varepsilon, W^\varepsilon)$ be a stationary semi-discrete state with nonnegative reconstructed density. Assume that Assumptions 3.1 and 3.2 hold. Then there exist constants $C_m, C_\rho > 0$, independent of ε , such that*

$$0 \leq m_j^\varepsilon \leq C_m, \quad 0 \leq \rho_{j,Q}^\varepsilon \leq C_\rho, \quad j = 1, \dots, J.$$

Proof. Since $n_{j,k}^\varepsilon \geq 0$,

$$\frac{1}{D_{\max}} m_j^\varepsilon \leq \eta_j^\varepsilon \leq \frac{1}{D_{\min}} m_j^\varepsilon.$$

Using (3.28) and (3.29), we obtain

$$-\delta_x^2 m_j^\varepsilon + \eta_j^\varepsilon \rho_{j,Q}^\varepsilon \leq K_{\max} \eta_j^\varepsilon + C_R m_j^\varepsilon + r_h.$$

Let j_* be a point where $M^\varepsilon := \max_j m_j^\varepsilon = m_{j_*}^\varepsilon$. Since $\delta_x^2 m_{j_*}^\varepsilon \leq 0$,

$$\eta_{j_*}^\varepsilon \rho_{j_*,Q}^\varepsilon \leq K_{\max} \eta_{j_*}^\varepsilon + C_R M^\varepsilon + r_h.$$

If $M^\varepsilon \leq e_Q/c_Q$, the bound is immediate. Otherwise, $\rho_{j_*,Q}^\varepsilon \geq c_Q M^\varepsilon - e_Q > 0$, and hence

$$\frac{M^\varepsilon}{D_{\max}} (c_Q M^\varepsilon - e_Q) \leq \left(\frac{K_{\max}}{D_{\min}} + C_R \right) M^\varepsilon + r_h.$$

This quadratic inequality gives a bound for M^ε independent of ε . The upper estimate for $\rho_{j,Q}^\varepsilon$ then follows from (3.31). $\square$

Stationary phase estimate. The density bound controls the coefficients in the discrete eigenvalue problem and hence the stationary phase equation.

Proposition 3.2 (Stationary phase estimate on a fixed trait grid). *Assume that*

$$0 \leq \rho_{j,Q}^\varepsilon \leq \bar{\rho}, \quad j = 1, \dots, J,$$

where $\bar{\rho}$ is independent of ε . Then the discrete effective Hamiltonian defined by (3.21) satisfies

$$|\mathcal{H}_k(\boldsymbol{\rho}_Q)| \leq C_{\mathcal{H}}, \quad k = 1, \dots, K,$$

where $C_{\mathcal{H}}$ is independent of ε .

Assume further that the stationary phase equation

$$\widehat{H}_G(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon) - \varepsilon \delta_\theta^2 u_k^\varepsilon = -\mathcal{H}_k(\boldsymbol{\rho}_Q) \quad (3.32)$$

holds with the Godunov numerical Hamiltonian

$$\widehat{H}_G(p^-, p^+) = -((p^-)_-)^2 - ((p^+)_+)^2.$$

Then

$$\Delta\theta \sum_{k=1}^K |D_\theta^+ u_k^\varepsilon|^2 \leq C_{\mathcal{H}}.$$

Consequently,

$$\max_k |D_\theta^+ u_k^\varepsilon| \leq \frac{\sqrt{C_{\mathcal{H}}}}{\sqrt{\Delta\theta}}.$$

If the phase is normalized by

$$\max_k u_k^\varepsilon = 0,$$

then

$$\max_k |u_k^\varepsilon| \leq \sqrt{C_{\mathcal{H}}}.$$

All constants are independent of ε , although they may depend on the fixed trait mesh.

Proof. The discrete eigenvalue problem is associated with the symmetric operator

$$-D_k \delta_x^2 - (K_j - \rho_{j,Q}^\varepsilon).$$

Its principal eigenvalue has the Rayleigh representation

$$\mathcal{H}_k(\boldsymbol{\rho}_Q) = \min_{\varphi \neq 0} \frac{D_k \Delta x \sum_j |\delta_x^+ \varphi_j|^2 - \Delta x \sum_j (K_j - \rho_{j,Q}^\varepsilon) \varphi_j^2}{\Delta x \sum_j \varphi_j^2}.$$

Since $0 \leq K_j \leq K_{\max}$ and $0 \leq \rho_{j,Q}^\varepsilon \leq \bar{\rho}$, we have

$$\mathcal{H}_k(\boldsymbol{\rho}_Q) \geq -K_{\max}.$$

Testing the quotient with a constant vector gives $\mathcal{H}_k(\boldsymbol{\rho}_Q) \leq \bar{\rho}$. Thus

$$|\mathcal{H}_k(\boldsymbol{\rho}_Q)| \leq \max\{K_{\max}, \bar{\rho}\} =: C_{\mathcal{H}}.$$

Set $q_k^\varepsilon = D_\theta^+ u_k^\varepsilon$. Then $D_\theta^- u_k^\varepsilon = q_{k-1}^\varepsilon$, and

$$-\widehat{H}_G(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon) = ((q_{k-1}^\varepsilon)_-)^2 + ((q_k^\varepsilon)_+)^2.$$

Summing (3.32) over the periodic trait grid eliminates $\delta_\theta^2 u^\varepsilon$. Therefore

$$\begin{aligned} \Delta\theta \sum_{k=1}^K |q_k^\varepsilon|^2 &= -\Delta\theta \sum_{k=1}^K \widehat{H}_G(D_\theta^- u_k^\varepsilon, D_\theta^+ u_k^\varepsilon) \\ &= \Delta\theta \sum_{k=1}^K \mathcal{H}_k(\rho_Q) \leq C_{\mathcal{H}}. \end{aligned}$$

The pointwise slope bound follows from the fixed-grid inequality $|q_k|^2 \leq (\Delta\theta)^{-1} \Delta\theta \sum_\ell |q_\ell|^2$. If $\max_k u_k^\varepsilon = 0$, connecting any node to a maximizer along the periodic grid gives

$$|u_k^\varepsilon| \leq \Delta\theta \sum_{\ell=1}^K |D_\theta^+ u_\ell^\varepsilon| \leq \left(\Delta\theta \sum_{\ell=1}^K |D_\theta^+ u_\ell^\varepsilon|^2 \right)^{1/2}.$$

This proves the claim. $\square$

Formal AP trait-selection limit. We finally pass formally to the fixed-grid rare-mutation limit. The statement is deliberately fixed-grid and conditional: it identifies the limiting selection mechanism once the density and phase estimates above are available.

Proposition 3.3 (Formal AP trait-selection limit). *Assume that the stationary semi-discrete states satisfy Assumptions 3.1 and 3.2, and that the phase is normalized by*

$$\max_k u_k^\varepsilon = 0.$$

Assume also that the stationary phase estimate of Proposition 3.2 applies. Then, up to a subsequence on the fixed grids,

$$\rho_Q^\varepsilon \rightarrow \rho^0, \quad u^\varepsilon \rightarrow u^0.$$

Moreover, the limiting phase satisfies the discrete constrained Hamilton–Jacobi system

$$\widehat{H}(D_\theta^- u_k^0, D_\theta^+ u_k^0) = -\mathcal{H}_k(\rho^0), \quad \max_k u_k^0 = 0.$$

For the Godunov Hamiltonian $\widehat{H}_G$, every maximizer k_ of u^0 satisfies*

$$\mathcal{H}_{k_*}(\rho^0) = 0.$$

If $\mathcal{H}_k(\rho^0)$ has a unique zero at k_m , then $k_ = k_m$. Hence the selected grid trait is identified by*

$$\theta_{k_m} \in \arg \max_k u_k^0.$$

Proof. Proposition 3.1 and (3.31) give a uniform bound on ρ_Q^ε . Since the grids are fixed, a subsequence converges to some ρ^0 . The normalized phase is also uniformly bounded on the fixed trait grid by Proposition 3.2; hence, up to a further subsequence, $u^\varepsilon \rightarrow u^0$.

Passing to the limit in the stationary phase equation gives the discrete constrained Hamilton–Jacobi system, because $\varepsilon \delta_\theta^2 u_k^\varepsilon \rightarrow 0$ on the fixed grid. For the Godunov Hamiltonian, $\widehat{H}_G \leq 0$, and therefore $\mathcal{H}_k(\rho^0) = -\widehat{H}_G(D_\theta^- u_k^0, D_\theta^+ u_k^0) \geq 0$. At a maximizer k_* of u^0 ,

$$D_\theta^- u_{k_*}^0 \geq 0, \quad D_\theta^+ u_{k_*}^0 \leq 0,$$

so $\widehat{H}_G(D_\theta^- u_{k_*}^0, D_\theta^+ u_{k_*}^0) = 0$. Hence $\mathcal{H}_{k_*}(\rho^0) = 0$. If this zero is unique, the maximizer must be k_m . $\square$

The preceding results should be read as a stationary fixed-grid AP structure rather than a complete AP convergence theorem. In particular, they do not provide a fully discrete time-AP estimate for the evolutionary problem. No stability bound uniform in ε is proved for a time-marching method with a time step independent of ε . This distinction is important because the time-dependent equation is a singular relaxation problem as $\varepsilon \rightarrow 0$, while the limiting stationary structure is an elliptic eigenvalue and constrained Hamilton–Jacobi structure. The constants above are uniform in ε for fixed spatial and trait grids. A simultaneous mesh-refinement and rare-mutation limit would require additional estimates on the reconstruction, the phase slopes, and the WKB remainder.

3.4 Fully discrete realization and time-step restrictions

We now collect the fully discrete formulas used in the computations and explain their practical restrictions. This subsection is separated from the semi-discrete analysis because the latter concerns stationary fixed-grid AP structure, whereas the former concerns a singular time-dependent relaxation problem. The fully discrete formulas below are implementation choices and numerical stabilization devices. They are not used in the proof of Propositions 3.1–3.3.

Let $\tau_n = t_{n+1} - t_n$, and let ρ^n and $H_k^n = \mathcal{H}_k(\rho^n)$ be computed from the current reconstructed density. A standard frozen-H phase update is

$$\frac{u_k^{n+1} - u_k^n}{\tau_n} + \widehat{H}(D_\theta^- u_k^n, D_\theta^+ u_k^n) - \varepsilon \delta_\theta^2 u_k^{n+1} = -H_k^n. \quad (3.33)$$

This formula is consistent with the semi-discrete phase equation, but it replaces

$$\int_{t_n}^{t_{n+1}} H_k(\rho(s)) ds$$

by $\tau_n H_k(\rho^n)$. This distinction is essential in the rare-mutation regime. Since $H_k(\rho(t))$ is coupled to the evolving density, it is generally time dependent. An $O(\tau_n)$ error in the phase may be harmless for u itself but can be amplified in the density reconstruction through $\exp(u/\varepsilon)$. This is one of the main reasons why a frozen-H time-marching scheme is not automatically AP in time.

For the split WKB amplitude equation, a fully implicit linear update is

$$\begin{aligned} & \varepsilon \frac{W_{j,k}^{n+1} - W_{j,k}^n}{\tau_n} - D_k \delta_x^2 W_{j,k}^{n+1} - \varepsilon^2 \delta_\theta^2 W_{j,k}^{n+1} + 2\varepsilon D_\theta \widehat{F}(W^{n+1}, u^{n+1})_{j,k} \\ & = W_{j,k}^{n+1} (K_j - \rho_j^n + H_k^n - 2\varepsilon \delta_\theta^2 u_k^{n+1}). \end{aligned} \quad (3.34)$$

At the fully discrete level, positivity of this linear update is tied to an M-matrix condition. A typical sufficient source restriction has the form

$$\frac{\tau_n}{\varepsilon} \max_{j,k} (K_j - \rho_j^n + H_k^n - 2\varepsilon \delta_\theta^2 u_k^n)_+ < 1, \quad (3.35)$$

up to the additional CFL-type restriction associated with any explicitly treated transport part. Thus the semi-discrete positivity result does not imply unconditional positivity for an arbitrary fully discrete time step.

The monotone explicit treatment of the Hamilton–Jacobi part also has its own time-step restriction. For a differentiable monotone numerical Hamiltonian, a sufficient CFL condition is

$$\frac{\tau_n}{\Delta\theta} \max_k \left(\partial_a \widehat{H}(D_\theta^- u_k^n, D_\theta^+ u_k^n) - \partial_b \widehat{H}(D_\theta^- u_k^n, D_\theta^+ u_k^n) \right) \leq 1. \quad (3.36)$$

For the Godunov Hamiltonian this condition can be written in the usual one-sided form in terms of the discrete slopes. In the numerical code this quantity is monitored as a phase CFL diagnostic.

A useful device for reducing the time-discrete WKB chain-rule defect is a time-direction integrating factor. Define

$$E_k^n = \exp(u_k^n/\varepsilon), \quad q_k^{n+1} = \frac{E_k^n}{E_k^{n+1}} = \exp\left(\frac{u_k^n - u_k^{n+1}}{\varepsilon}\right). \quad (3.37)$$

Then one may update the amplitude by

$$\begin{aligned} & \varepsilon \frac{W_{j,k}^{n+1} - q_k^{n+1} W_{j,k}^n}{\tau_n} - D_k \delta_x^2 W_{j,k}^{n+1} - \varepsilon^2 (E_k^{n+1})^{-1} \delta_\theta^2 (W_{j,\cdot}^{n+1} E_{\cdot,k}^{n+1})_k \\ & = (K_j - \rho_j^n) W_{j,k}^{n+1}. \end{aligned} \quad (3.38)$$

Multiplying (3.38) by E_k^{n+1} gives the density-level identity

$$\varepsilon \frac{n_{j,k}^{n+1} - n_{j,k}^n}{\tau_n} - D_k \delta_x^2 n_{j,k}^{n+1} - \varepsilon^2 \delta_\theta^2 n_{j,k}^{n+1} = (K_j - \rho_j^n) n_{j,k}^{n+1}. \quad (3.39)$$

This identity removes one source of discrete chain-rule error between u , W , and n . It does not remove the time-dependence of $H_k(\rho(t))$ in the phase equation and therefore does not by itself prove time-AP stability.

For positivity with respect to the reaction term, the code also provides a Patankar production–destruction option. With

$$r_j^n = K_j - \rho_j^n, \quad (r_j^n)_+ = \max(r_j^n, 0), \quad (r_j^n)_- = \max(-r_j^n, 0),$$

the density-level reaction part may be written as

$$\varepsilon \frac{n_{j,k}^{n+1} - n_{j,k}^n}{\tau_n} - D_k \delta_x^2 n_{j,k}^{n+1} - \varepsilon^2 \delta_\theta^2 n_{j,k}^{n+1} = (r_j^n)_+ n_{j,k}^n - (r_j^n)_- n_{j,k}^{n+1}. \quad (3.40)$$

Equivalently, in the WKB amplitude variable,

$$\begin{aligned} & \varepsilon \frac{W_{j,k}^{n+1} - q_k^{n+1} W_{j,k}^n}{\tau_n} - D_k \delta_x^2 W_{j,k}^{n+1} - \varepsilon^2 (E_k^{n+1})^{-1} \delta_\theta^2 (W_{j,\cdot}^{n+1} E_{\cdot,k}^{n+1})_k \\ & = (r_j^n)_+ q_k^{n+1} W_{j,k}^n - (r_j^n)_- W_{j,k}^{n+1}. \end{aligned} \quad (3.41)$$

This form improves positivity of the reaction update, but it should be regarded as a fully discrete stabilization rather than a new semi-discrete AP argument.

Because W changes under the max-gauge normalization, a residual based directly on $W^{n+1} - W^n$ may be misleading. The code therefore reports a gauge-invariant time-difference residual. With

$$\tilde{u}_k^n = u_k^n - \max_{\ell} u_{\ell}^n, \quad n_{j,k}^n = W_{j,k}^n \exp(u_k^n / \varepsilon),$$

one diagnostic residual is

$$E_{\text{gi}}^n = \max \left\{ \frac{\|\tilde{u}^{n+1} - \tilde{u}^n\|_{\infty}}{\tau_n}, \frac{\|n^{n+1} - n^n\|_{\infty}}{\tau_n(1 + \|n^{n+1}\|_{\infty})} \right\}. \quad (3.42)$$

In very small mutation regimes, even this residual should be interpreted with care, since the phase may have already identified the selected trait while the amplitude still relaxes on a slower numerical scale.

A practical adaptive time-step safeguard is

$$\tau_n = \min \left\{ \tau_{\max}, \eta \frac{\varepsilon}{\max\{1, \Lambda_r^n, \Lambda_H^n\}} \right\}, \quad \eta \in (0, 1), \quad (3.43)$$

where $\Lambda_H^n = \max_k |H_k^n|$. For the split update one may take

$$\Lambda_r^n = \max_{j,k} (K_j - \rho_j^n + H_k^n - 2\varepsilon \delta_{\theta}^2 u_k^n)_+,$$

whereas for the time-density-compatible update one may take $\Lambda_r^n = \max_j (K_j - \rho_j^n)_+$. This is a sufficient numerical safeguard for the stiff source scale, not a sharp necessary condition.

Finally, the code contains an experimental projective WKB integrator for time evolution. For one macro step τ_n , it first performs M micro steps with $\delta t_{\text{mic}} = c_{\text{mic}} \varepsilon$, and then extrapolates the slow phase variable to the macro time. In the safer implementation only u is extrapolated, while W is corrected by additional micro steps if desired. This projective integrator is designed to resolve the initial $O(\varepsilon)$ relaxation layer and then advance the selected-trait dynamics with a larger macro time step. Numerical tests indicate that it is effective for tracking θ_{WKB} , although the amplitude residual may decay more slowly than the phase residual. Therefore it should be presented as a computational stabilization for the evolutionary calculation, not as part of the stationary AP proof.

The main message is that the semi-discrete analysis proves a stationary fixed-grid AP structure. A fully discrete time-marching method with τ independent of ε requires additional stability mechanisms and is not proved here to be AP in time. When $H_k(\rho(t))$ changes appreciably during one step, a frozen-H update may produce a phase error that is magnified by $\exp(u/\varepsilon)$. Thus one should either use a sufficiently small time step, use a stabilization such as projective integration, or interpret the computation primarily through the phase-based selected trait rather than through a fully converged density profile.

4 Numerical experiments

This section describes the numerical experiments used to test the WKB formulation. The tests are organized to match the main assumptions and claims of the method: the structural assumptions used in the stationary AP argument, the emergence of concentration in the trait direction, the accuracy of the reconstructed density when the grid is sufficiently fine, the advantage of the WKB phase on coarse trait grids, standard accuracy checks, and a two-dimensional spatial experiment.

4.1 Common computational setting

Unless otherwise stated, the one-dimensional tests use

$$\Omega = (0, 1), \quad \mathbb{T} = [0, 1),$$

with homogeneous Neumann boundary conditions in the spatial variable and periodic boundary conditions in the trait variable. The baseline coefficients are

$$K(x) = K_0 + K_1 \cos(2\pi x), \quad K_0 = 1, \quad K_1 = 0.5, \quad (4.44)$$

$$D(\theta) = D_{\min} + D_1(1 - \cos(2\pi(\theta - \theta_m))), \quad D_{\min} = 0.2, \quad D_1 = 0.4, \quad \theta_m = 0.35. \quad (4.45)$$

Thus the dispersal rate has a unique minimum at θ_m , which is the expected selected trait in the rare-mutation limit. The default initial data are

$$u_k^0 = -\alpha_0(1 - \cos(2\pi(\theta_k - \theta_0))), \quad \alpha_0 = 0.05, \quad \theta_0 = 0.70, \quad (4.46)$$

$$W_{j,k}^0 = 1 + 0.1 \cos(2\pi x_j), \quad n_{j,k}^0 = W_{j,k}^0 \exp(u_k^0/\varepsilon). \quad (4.47)$$

The initial maximizer θ_0 is deliberately placed away from θ_m , so the computation tests selection rather than preservation of a well-prepared peak.

For the WKB computation we use the phase equation in (3.22) together with either the split amplitude update or the density-compatible update. The default density in the nonlocal reaction term is the log-stabilized direct quadrature

$$\rho_{j,Q}^n = \exp(s^n/\varepsilon) \Delta\theta \sum_{k=1}^K W_{j,k}^n \exp\left(\frac{u_k^n - s^n}{\varepsilon}\right), \quad s^n = \max_k u_k^n. \quad (4.48)$$

A Laplace reconstruction near the maximizer of u^n is also used as a diagnostic and, in selected tests, as an alternative reconstruction; its implementation is described in Appendix C. After each time step, we use the gauge normalization

$$c^n = \max_k u_k^n, \quad u_k^n \leftarrow u_k^n - c^n, \quad W_{j,k}^n \leftarrow W_{j,k}^n \exp(c^n/\varepsilon), \quad (4.49)$$

which preserves the reconstructed density $W_{j,k}^n \exp(u_k^n/\varepsilon)$. **The fully discrete time updates, residuals, and time-step restrictions used in the code are summarized in Subsection 3.4.**

The trait marginal, selected traits, and concentration width are computed as

$$P_k = \int_{\Omega} n(x, \theta_k) dx, \quad \theta_{\text{dir}} = \theta_{\arg \max_k P_k}, \quad \theta_{\text{WKB}} = \theta_{\arg \max_k u_k}, \quad (4.50)$$

$$d_{\mathbb{T}}(a, b) = \min_{q \in \mathbb{Z}} |a - b + q|, \quad (4.51)$$

$$\sigma_{\theta} = \left(\frac{\sum_{k=1}^K d_{\mathbb{T}}(\theta_k, \theta_{\text{WKB}})^2 P_k}{\sum_{k=1}^K P_k} \right)^{1/2}. \quad (4.52)$$

For the weighted remainder assumption, we monitor

$$\Gamma_R^{\varepsilon} = \max_{1 \leq j \leq J} \frac{(R_j^{\varepsilon, D})_+}{m_j^{\varepsilon} + 10^{-14}}, \quad m_j^{\varepsilon} = \Delta\theta \sum_{k=1}^K n_{j,k}^{\varepsilon}, \quad R_j^{\varepsilon, D} = \Delta\theta \sum_{k=1}^K \frac{R_{j,k}^{\varepsilon}}{D_k}. \quad (4.53)$$

A bounded value of Γ_R^{ε} as ε decreases is not a proof of Assumption 3.1, but it is a useful numerical diagnostic for the positive part of the weighted defect. **In addition to Γ_R^{ε} , the fully discrete diagnostics described in Subsection 3.4 are reported. In particular, the phase residual, amplitude residual, density residual, and the selected trait are monitored separately, since the WKB phase may converge to the selected trait before the full amplitude has reached a numerical steady state.** For the direct density solver, the analogous residual is computed with $n^{n+1} - n^n$.

4.2 Test 1: numerical verification of the structural assumptions

The first test checks the structural quantities that appear in the stationary AP argument. We use

$$J = 200, \quad K = 64, \quad \varepsilon \in \{5 \times 10^{-2}, 2 \times 10^{-2}, 10^{-2}, 5 \times 10^{-3}\}.$$

Both the split WKB update and the density-compatible update are run. The following quantities are reported:

- $\min_{j,k} W_{j,k}$, to monitor positivity of the amplitude;
- $\max_j \rho_{j,Q}$, to monitor the density bound;
- Γ_R^ε , to test the one-sided weighted remainder condition;
- $\min_k H_k(\rho_Q)$ and the location of this minimum, to check consistency with the selected dispersal minimizer;
- $d_T(\theta_{\text{WKB}}, \theta_m)$, to check trait selection.

The density-compatible update is expected to keep Γ_R^ε controlled by the discrete second difference of $1/D_k$, as explained in Appendix D. For the generic split update, this table is a diagnostic for the conditional assumption.

Table 1: Test 1. Numerical diagnostics for the structural assumptions.

variant	ε	$\min W$	$\max \rho_Q$	Γ_R^ε	θ_{WKB}	$d_T(\theta_{\text{WKB}}, \theta_m)$
split	5×10^{-2}	—	—	—	—	—
split	2×10^{-2}	—	—	—	—	—
split	10^{-2}	—	—	—	—	—
split	5×10^{-3}	—	—	—	—	—
density-compatible	5×10^{-2}	—	—	—	—	—
density-compatible	2×10^{-2}	—	—	—	—	—
density-compatible	10^{-2}	—	—	—	—	—
density-compatible	5×10^{-3}	—	—	—	—	—

4.3 Test 2: long-time concentration in the trait direction

This test checks whether the solution actually concentrates in the trait direction after sufficiently large time. We use

$$J = 200, \quad K = 128, \quad \varepsilon \in \{5 \times 10^{-2}, 2 \times 10^{-2}, 10^{-2}\},$$

and store snapshots at several times, for example

$$t \in \{1, 5, 10, 20, 40, 80\}.$$

For each snapshot, we plot the trait marginal $P_k(t)$, the phase $u_k(t)$, and record $\theta_{\text{WKB}}(t)$, $\theta_{\text{dir}}(t)$, and $\sigma_\theta(t)$. The expected behavior is that the maximizer of u moves toward θ_m , while the concentration width σ_θ decreases with time and with ε .

Table 2: Test 2. Long-time concentration diagnostics.

ε	t	$\theta_{\text{WKB}}(t)$	$\theta_{\text{dir}}(t)$	$\sigma_\theta(t)$	$\min_k H_k(\rho_Q(t))$
5×10^{-2}	—	—	—	—	—
2×10^{-2}	—	—	—	—	—
10^{-2}	—	—	—	—	—

4.4 Test 3: fine-grid comparison between direct density and WKB reconstruction

When the trait grid is sufficiently fine, the direct density solution and the WKB reconstruction should be close at finite ε . We therefore run the direct solver and the WKB solver on the same fine grid,

$$J = 200, \quad K \in \{256, 512\}, \quad \varepsilon \in \{2 \times 10^{-2}, 10^{-2}\}.$$

The reconstructed WKB density is

$$n_{j,k}^{\text{WKB}} = W_{j,k} \exp(u_k/\varepsilon),$$

and it is compared with the direct density $n_{j,k}^{\text{dir}}$ using

$$e_1 = \frac{\Delta\theta \sum_k \int_\Omega |n^{\text{WKB}}(x, \theta_k) - n^{\text{dir}}(x, \theta_k)| dx}{\Delta\theta \sum_k \int_\Omega |n^{\text{dir}}(x, \theta_k)| dx}, \quad (4.54)$$

$$e_2 = \frac{(\Delta\theta \sum_k \int_\Omega |n^{\text{WKB}}(x, \theta_k) - n^{\text{dir}}(x, \theta_k)|^2 dx)^{1/2}}{(\Delta\theta \sum_k \int_\Omega |n^{\text{dir}}(x, \theta_k)|^2 dx)^{1/2}}. \quad (4.55)$$

We also compare the trait marginals and selected traits. This test is not meant to show the main advantage of WKB; rather, it verifies that the WKB variables reconstruct the same finite- ε density when the direct grid is fine enough.

Table 3: Test 3. Fine-grid comparison of direct density and WKB reconstruction.

ε	K	e_1	e_2	θ_{dir}	θ_{WKB}	$d_{\mathbb{T}}(\theta_{\text{dir}}, \theta_{\text{WKB}})$
2×10^{-2}	256	—	—	—	—	—
2×10^{-2}	512	—	—	—	—	—
10^{-2}	256	—	—	—	—	—
10^{-2}	512	—	—	—	—	—

4.5 Test 4: coarse trait grids and AP advantage of the WKB phase

This test demonstrates the main computational advantage of the WKB formulation. We keep ε small and vary the number of trait grid points:

$$J = 200, \quad \varepsilon = 5 \times 10^{-3}, \quad K \in \{16, 24, 32, 48, 64, 128\}.$$

A high-resolution direct computation or, when this is too expensive, the limiting trait θ_m , is used as the reference. We compare the selected traits

$$e_{\text{dir}} = d_{\mathbb{T}}(\theta_{\text{dir}}, \theta_{\text{ref}}), \quad e_{\text{WKB}} = d_{\mathbb{T}}(\theta_{\text{WKB}}, \theta_{\text{ref}}).$$

The expected outcome is that the direct density solver requires a much finer trait grid to localize the narrow peak, whereas the WKB phase remains a smoother object and identifies the concentration location on substantially coarser trait grids.

Table 4: Test 4. Coarse trait-grid comparison and AP advantage.

K	θ_{ref}	θ_{dir}	θ_{WKB}	e_{dir}	e_{WKB}	CPU ratio
16	—	—	—	—	—	—
24	—	—	—	—	—	—
32	—	—	—	—	—	—
48	—	—	—	—	—	—
64	—	—	—	—	—	—
128	—	—	—	—	—	—

4.6 Test 5: accuracy and convergence tests

The fifth test checks standard numerical accuracy. Since closed-form solutions are not available, we compute a reference solution on a refined grid and compare coarser computations with it. For example, use

$$\varepsilon = 2 \times 10^{-2}, \quad (J_{\text{ref}}, K_{\text{ref}}, \tau_{\text{ref}}) = (400, 256, 2.5 \times 10^{-4}),$$

and compare against coarser triples such as

$$(J, K, \tau) \in \{(100, 64, 10^{-3}), (200, 128, 5 \times 10^{-4}), (300, 192, 3.33 \times 10^{-4})\}.$$

The reported quantities are the errors in ρ , the trait marginal P_k , the phase u_k up to the gauge $\max_k u_k = 0$, and the selected trait. If different trait grids are used, the reference phase and marginal are interpolated periodically in θ . We also perform a time-step sensitivity test for the WKB evolution. This test records E_{gi}^n , its phase and amplitude components, the selected trait θ_{WKB} , and the diagnostics associated with the projective time integrator. The purpose is to separate two questions. The first one is whether the WKB phase correctly captures the fittest trait on a coarse trait grid. The second one is whether the full amplitude and density variables have reached a fully converged steady state. In the very small mutation regime, the first may hold before the second.

Table 5: Test 5. Accuracy against a refined WKB reference solution.

J	K	τ	error in ρ	error in P	trait error
100	64	10^{-3}	—	—	—
200	128	5×10^{-4}	—	—	—
300	192	3.33×10^{-4}	—	—	—

4.7 Test 6: two-dimensional spatial experiment

The last test checks that the WKB strategy is not restricted to one spatial dimension. We solve the model on

$$\Omega = (0, 1)^2$$

using a triangular finite-element discretization in $x = (x_1, x_2)$ and the same periodic finite-difference discretization in θ . A representative two-dimensional landscape is

$$K(x_1, x_2) = 1 + 0.35 \cos(2\pi x_1) \cos(2\pi x_2) + 0.15 \cos(4\pi x_1), \quad (4.56)$$

with the same baseline dispersal function (4.45). Use

$$K_\theta \in \{32, 64\}, \quad \varepsilon \in \{2 \times 10^{-2}, 10^{-2}\}.$$

The outputs are the spatial total density $\rho(x_1, x_2)$, the trait marginal P_k , the phase u_k , and the selected trait θ_{WKB} . This test is mainly qualitative: it should show that the spatial heterogeneity is resolved by the finite-element amplitude/eigenvalue solves, while the concentration location is still captured through the one-dimensional phase variable.

Table 6: Test 6. Two-dimensional spatial WKB experiment.

ε	K_θ	mesh size	θ_{WKB}	$d_{\mathbb{T}}(\theta_{\text{WKB}}, \theta_m)$
2×10^{-2}	32	—	—	—
2×10^{-2}	64	—	—	—
10^{-2}	32	—	—	—
10^{-2}	64	—	—	—

A Numerical Hamiltonians and one-sided derivative reconstruction

For the monotone theory in Section 3, the numerical Hamiltonian $\hat{H}(a, b)$ is assumed to be consistent with $-p^2$, nondecreasing in its first argument, and nonincreasing in its second argument. Two standard choices are the Lax–Friedrichs Hamiltonian

$$\hat{H}_{\text{LF}}(a, b) = -\left(\frac{a+b}{2}\right)^2 + \frac{\alpha}{2}(a-b), \quad (1.57)$$

where α is chosen no smaller than the relevant characteristic speed, and the Godunov Hamiltonian

$$\hat{H}_G(a, b) = -(a^-)^2 - (b^+)^2, \quad a^- := \min\{a, 0\}, \quad b^+ := \max\{b, 0\}. \quad (1.58)$$

The stationary phase estimate in Proposition 3.2 uses $\hat{H}_G$.

Some numerical runs use a high-order one-sided WENO reconstruction for the derivatives entering the Hamiltonian. This is an implementation option, not the monotone Hamiltonian used in the proof. For a periodic grid function u_k , define the left-biased quantities

$$a = u_{k-3} - 2u_{k-2} + u_{k-1}, \quad b = u_{k-2} - 2u_{k-1} + u_k,$$

$$c = u_{k-1} - 2u_k + u_{k+1}, \quad d = u_k - 2u_{k+1} + u_{k+2}.$$

Set

$$I_0 = 13(a-b)^2 + 3(a-3b)^2, \quad I_1 = 13(b-c)^2 + 3(b+c)^2,$$

$$I_2 = 13(c-d)^2 + 3(3c-d)^2,$$

and

$$\alpha_0 = \frac{1}{(\epsilon_0 + I_0)^2}, \quad \alpha_1 = \frac{6}{(\epsilon_0 + I_1)^2}, \quad \alpha_2 = \frac{3}{(\epsilon_0 + I_2)^2}, \quad \omega_r = \frac{\alpha_r}{\alpha_0 + \alpha_1 + \alpha_2}.$$

The left derivative reconstruction used in the implementation is

$$p_k^- = \frac{1}{\Delta\theta} \left[\frac{-(u_{k-1} - u_{k-2}) + 7(u_k - u_{k-1}) + 7(u_{k+1} - u_k) - (u_{k+2} - u_{k+1})}{12} - \frac{\omega_0}{3}(a - 2b + c) - \frac{\omega_2 - 1/2}{6}(b - 2c + d) \right]. \quad (1.59)$$

The right-biased derivative p_k^+ is obtained by applying the same formula to the reversed stencil. Equivalently, with

$$a^+ = u_{k+3} - 2u_{k+2} + u_{k+1}, \quad b^+ = u_{k+2} - 2u_{k+1} + u_k, \\ c^+ = u_{k+1} - 2u_k + u_{k-1}, \quad d^+ = u_k - 2u_{k-1} + u_{k-2},$$

and the corresponding weights ω_0^+, ω_2^+ ,

$$p_k^+ = \frac{1}{\Delta\theta} \left[\frac{-(u_{k-1} - u_{k-2}) + 7(u_k - u_{k-1}) + 7(u_{k+1} - u_k) - (u_{k+2} - u_{k+1})}{12} + \frac{\omega_0^+}{3}(a^+ - 2b^+ + c^+) + \frac{\omega_2^+ - 1/2}{6}(b^+ - 2c^+ + d^+) \right]. \quad (1.60)$$

The WENO option then uses a Roe-type entropy-fix numerical Hamiltonian for $H(p) = -p^2$ with the reconstructed values (p_k^+, p_k^-) . Since this high-order reconstruction is not monotone in the sense assumed in Section 3, the numerical experiments report it separately from the monotone Godunov/Lax–Friedrichs choices when needed.

B Conservative fluxes for the split amplitude equation

The continuous conservative flux in the split amplitude equation is

$$F = -W\partial_\theta u.$$

On a uniform periodic trait grid, write

$$D_\theta \widehat{F}(W, u)_{j,k} = \frac{\widehat{F}_{j,k+1/2} - \widehat{F}_{j,k-1/2}}{\Delta\theta}, \quad a_{k+1/2} = -\frac{u_{k+1} - u_k}{\Delta\theta}.$$

The upwind flux is

$$\widehat{F}_{j,k+1/2}^{\text{up}} = a_{k+1/2}^+ W_{j,k} + a_{k+1/2}^- W_{j,k+1}, \quad a^+ = \max\{a, 0\}, \quad a^- = \min\{a, 0\}. \quad (2.61)$$

A Lax–Friedrichs flux is

$$\widehat{F}_{j,k+1/2}^{\text{LF}} = \frac{1}{2} a_{k+1/2} (W_{j,k} + W_{j,k+1}) - \frac{1}{2} \alpha_{k+1/2} (W_{j,k+1} - W_{j,k}), \quad \alpha_{k+1/2} \geq |a_{k+1/2}|. \quad (2.62)$$

Both fluxes are conservative in the periodic trait variable. The upwind flux also satisfies the quasi-positivity condition used in Lemma 3.1.

C Density reconstruction and gauge normalization

The default reconstruction of ρ in the code is the log-stabilized quadrature (4.48). This is algebraically identical to the composite quadrature

$$\rho_{j,Q}^n = \Delta\theta \sum_k W_{j,k}^n \exp(u_k^n/\varepsilon),$$

but avoids underflow or overflow when ε is small. The max-gauge (4.49) is applied after each WKB time step. It leaves $n_{j,k} = W_{j,k} \exp(u_k/\varepsilon)$ unchanged and keeps $\max_k u_k = 0$. **The fully discrete time updates and the corresponding time-step restrictions are stated together in Subsection 3.4; they are no longer dispersed in this appendix.**

For diagnostic purposes, a Laplace reconstruction can be used near a nondegenerate maximizer. Let k_* be an index at which u_k is maximal. The periodic stencil is represented in local coordinates

$$z_{k_*-1} = -\Delta\theta, \quad z_{k_*} = 0, \quad z_{k_*+1} = \Delta\theta,$$

even if the stencil crosses the endpoint of $[0, 1)$. With

$$d_{\theta\theta}u_{k_*} = \frac{u_{k_*+1} - 2u_{k_*} + u_{k_*-1}}{(\Delta\theta)^2},$$

the parabolic vertex is

$$z_* = -\frac{\Delta\theta}{2} \frac{u_{k_*+1} - u_{k_*-1}}{u_{k_*+1} - 2u_{k_*} + u_{k_*-1}}. \quad (3.63)$$

The reconstructed trait location is $\theta_* = (\theta_{k_*} + z_*) \bmod 1$. Interpolating W and u quadratically on this local stencil gives $W(x_j, \theta_*)$ and $u(\theta_*)$, and

$$\rho_j^{\text{Lap}} = W(x_j, \theta_*) \exp\left(\frac{u(\theta_*)}{\varepsilon}\right) \left(\frac{2\pi\varepsilon}{|d_{\theta\theta}u_{k_*}|}\right)^{1/2}. \quad (3.64)$$

If $d_{\theta\theta}u_{k_*} \geq 0$ or the curvature is too small, the implementation falls back to the log-stabilized direct quadrature. In practice, the exponential factor is evaluated in a logarithmic form when ε is very small.

D A sufficient condition for the one-sided remainder bound

For the density-compatible variant, the reconstructed density satisfies the conservative density equation exactly at the semi-discrete level. Hence, in the notation of (3.24),

$$R_{j,k}^\varepsilon = \varepsilon^2 \delta_\theta^2 n_{j,k}^\varepsilon.$$

Periodic summation by parts gives

$$R_j^{\varepsilon,D} = \varepsilon^2 \Delta\theta \sum_k \frac{\delta_\theta^2 n_{j,k}^\varepsilon}{D_k} = \varepsilon^2 \Delta\theta \sum_k n_{j,k}^\varepsilon \delta_\theta^2 \left(\frac{1}{D_k}\right). \quad (4.65)$$

Therefore, for $0 < \varepsilon \leq 1$,

$$(R_j^{\varepsilon,D})_+ \leq C_{D,h} m_j^\varepsilon, \quad C_{D,h} := \max_k \left| \delta_\theta^2 \left(\frac{1}{D_k}\right) \right|. \quad (4.66)$$

This verifies Assumption 3.1 with $C_R = C_{D,h}$ and $r_h = 0$. If D is smooth and the standard centered periodic difference is used, $C_{D,h}$ remains bounded under regular trait-grid refinement.

For a generic split WKB discretization, the same conclusion is not automatic because the reconstructed density contains discrete chain-rule defects. A simple sufficient condition is that there exists a constant C_E , independent of ε , such that

$$\exp\left(\frac{u_{k+1}^\varepsilon - u_k^\varepsilon}{\varepsilon}\right) + \exp\left(\frac{u_{k-1}^\varepsilon - u_k^\varepsilon}{\varepsilon}\right) \leq C_E, \quad k = 1, \dots, K, \quad (4.67)$$

and that the positive part of the local reconstruction defect satisfies

$$(R_{j,k}^\varepsilon)_+ \leq C_0 \sum_{\ell \in \{k-1, k, k+1\}} n_{j,\ell}^\varepsilon + r_{j,k,h}, \quad \Delta\theta \sum_k \frac{r_{j,k,h}}{D_k} \leq r_h. \quad (4.68)$$

Then positivity of n^ε , periodicity in k , and the lower bound on D_k imply Assumption 3.1. Condition (4.67) is stronger than a uniform bound on the ordinary discrete derivative of u^ε , which is why the main theorem states the split-WKB AP structure conditionally.

E Direct density solver and two-dimensional finite-element realization

For comparison, the original density equation can be discretized by the old linear implicit update

$$\varepsilon \frac{n_{j,k}^{n+1} - n_{j,k}^n}{\tau} - D_k \delta_x^2 n_{j,k}^{n+1} - \varepsilon^2 \delta_\theta^2 n_{j,k}^{n+1} = n_{j,k}^{n+1} (K_j - \rho_j^n), \quad \rho_j^n = \Delta\theta \sum_k n_{j,k}^n. \quad (5.69)$$

In the revised code, the direct density solver also accepts the Patankar reaction option

$$\varepsilon \frac{n_{j,k}^{n+1} - n_{j,k}^n}{\tau} - D_k \delta_x^2 n_{j,k}^{n+1} - \varepsilon^2 \delta_\theta^2 n_{j,k}^{n+1} = (K_j - \rho_j^n)_+ n_{j,k}^n - (K_j - \rho_j^n)_- n_{j,k}^{n+1}. \quad (5.70)$$

This option is used to match the positivity-oriented WKB update when direct and WKB runs are compared. The same spatial and trait grids are used for the direct and WKB solvers in the one-dimensional comparison tests.

For the two-dimensional spatial experiment, the spatial Laplacian and the eigenvalue problem are discretized by linear finite elements on a triangular mesh. For each trait node θ_k , the discrete effective Hamiltonian is computed from the generalized eigenvalue problem

$$A_k(\rho_Q) N^{(k)} = H_k(\rho_Q) M N^{(k)}, \quad (5.71)$$

where M is the finite-element mass matrix and

$$(A_k)_{ab} = D_k \int_\Omega \nabla \varphi_a \cdot \nabla \varphi_b \, dx - \int_\Omega (K(x) - \rho_Q(x)) \varphi_a \varphi_b \, dx. \quad (5.72)$$

The Neumann boundary condition is natural in this finite-element formulation. The phase equation remains one-dimensional in θ , and the amplitude equation is solved for all spatial finite-element degrees of freedom and trait nodes.

References

- [1] L. Almeida, B. Perthame, and X. Ruan, An asymptotic preserving scheme for capturing concentrations in age-structured models arising in adaptive dynamics, *Journal of Computational Physics*, 464 (2022), 111335.
- [2] G. Barles and B. Perthame, Concentrations and constrained Hamilton–Jacobi equations arising in adaptive dynamics, *Contemporary Mathematics*, 439 (2007), 57–68.
- [3] S. Mirrahimi and B. Perthame, Asymptotic analysis of a selection model with space, *Journal de Mathématiques Pures et Appliquées*, 104 (2015), 1108–1118.
- [4] B. Perthame and P. E. Souganidis, Rare mutations limit of a steady state dispersal evolution model, *Mathematical Modelling of Natural Phenomena*, 11 (2016), 154–166.