An asymptotic preserving scheme for kinetic equations describing bacterial traveling pulses

Canan Akkoyunlu* Jose A. Carrillo† Gissell Estrada-Rodriguez‡
Xinran Ruan§

Abstract

We consider a kinetic model for bacterial chemotaxis that describes the propagation of traveling pulses driven by a nutrient and a self-produced chemoattractant. In the diffusive regime, the model converges to a macroscopic drift–diffusion system with bounded chemotactic fluxes. We construct a fully discrete asymptotic preserving (AP) finite difference scheme based on a micro–macro decomposition of the kinetic equation. The method is stable uniformly with respect to the scaling parameter and, as this parameter tends to zero, reduces to a consistent discretization of the limiting macroscopic equations. We prove positivity and mass conservation for the limiting scheme, and use a positivity-preserving correction for the kinetic solver. Numerical simulations reproduce the experimentally observed traveling pulses at both the kinetic and the macroscopic level, validate the theoretical traveling wave speed, and illustrate the convergence of the kinetic model to its macroscopic limit.

1 Introduction

Chemotaxis is the directed movement of organisms in response to chemical fluctuations in the environment. This behavior is essential for survival [9] and plays a crucial role in many biological processes in multicellular organisms, such as wound healing, fertilization and inflammation [16], while unicellular organisms use it as a foraging mechanism [10, 18]. Many bacteria, in particular, navigate toward favorable conditions by sensing gradients of nutrients or chemoattractants, and away from harmful substances.

*Department of Mathematics and Computer Science, İstanbul Kültür University, 34158 Bakırköy, İstanbul, Türkiye (c.kaya@iku.edu.tr)

†Mathematical Institute, University of Oxford, Woodstock Road, Oxford, OX2 6GG, UK (jose.carrillo@maths.ox.ac.uk)

‡Universitat Politecnica de Catalunya, Barcelona, 08034, Spain. Centre de Recerca Matematica, 08193 Bellaterra, Barcelona, Spain (gissell.estrada@upc.edu)

§School of Mathematical Sciences, Capital Normal University, Beijing, China, 100048 (xinran.ruan@cnu.edu.cn).

Although bacterial chemotaxis has been studied since the end of the 19th century, the first precise mathematical model was proposed by Keller and Segel [12]. They introduced a coupled system of partial differential equations (PDEs) for the cell density $\rho(t,x)$ and the chemical concentration $S(t,x)$:

$$\frac{\partial \rho}{\partial t} = \nabla \cdot (D_\rho(\rho,S)\nabla\rho - \chi(\rho,S)\nabla S) + f(\rho,S), \tag{1.1a}$$

$$\frac{\partial S}{\partial t} = D_S \Delta S + g(\rho,S) - h(\rho,S)S. \tag{1.1b}$$

Here $D_\rho(\rho,S)$ denotes the bacterial diffusivity, $\chi(\rho,S)$ the chemotactic sensitivity, and f, g, and h describe cell growth and death, chemical production, and degradation, respectively.

We consider the migration of *E. coli* in response to a nutrient N and a self-produced chemoattractant S, motivated by the microchannel experiments in [17]. In these experiments, approximately 10^5 cells are initially confined near the left end of a 2 cm channel, while nutrient is supplied from the right. Within about 100 s, the bacteria organize into a dense pulse that propagates toward the nutrient source with nearly constant speed and mass. The pulse exhibits a characteristic asymmetric profile, with a diffuse front and a steep rear, which is maintained during propagation. Since the duration of the experiment is short compared with the cell division cycle, bacterial growth is neglected.

Following the kinetic descriptions of Alt [1] and Dunbar and Othmer [15], we model the bacterial population by the mesoscopic density $f(t,x,v)$, where $x \in \mathbb{R}$ and $v \in V$ denotes the velocity. Since we are interested in planar traveling waves, the dynamics are governed by the one-dimensional kinetic equation

$$\partial_t f(t,x,v) + v\partial_x f(t,x,v) = \int_V \Big(T(t,x,v,v')f(t,x,v') - T(t,x,v',v)f(t,x,v) \Big)\,\mathrm{d}v' \ . \tag{1.2}$$

The turning kernel T describes the change in direction during the tumbling phase of the movement. T depends on the space and time variables through the chemical signals in the medium, and $T(\cdot,v,v')$ represents the change of velocity from v' to v. For simplicity we are going to assume that T only depends on the current velocity with the condition that $\int_V T(t,x,v)\,\mathrm{d}v = 1$.

Following [17], we consider the influence of two chemical species: a chemotactic signal $S(t,x)$ released by the bacteria, and a nutriet $N(t,x)$ consumed by the them. We assume that both chemical cues contribute additively to the turning frequency. More precisely, we define

$$\widetilde{\psi}_1(t,x,v) = \chi_S \phi\left(\frac{DS}{Dt}\Big|_v \right) + \chi_N \phi\left(\frac{DN}{Dt}\Big|_v \right), \qquad \frac{D}{Dt} = \partial_t + v\partial_x,$$

where ϕ is the intracellular response function and χ_S, χ_N are the chemotactic sensitivities. Since the velocity average of $\widetilde{\psi}_1$ does not vanish in general, we introduce the centered perturbation

$$\psi_1(t,x,v) = \widetilde{\psi}_1(t,x,v) - \frac{1}{|V|}\int_V \widetilde{\psi}_1(t,x,w)\,dw, \tag{1.3}$$

which satisfies $\int_V \psi_1(t,x,v)\,dv = 0$. The rescaled turning kernel is then defined by

$$T_\varepsilon(t,x,v) = |V|\,\frac{1+\varepsilon\widetilde{\psi}_1(t,x,v)}{\displaystyle\int_V \big(1+\varepsilon\widetilde{\psi}_1(t,x,w)\big)\,dw},$$

and admits the first-order expansion

$$T_\varepsilon(t,x,v) = 1 + \varepsilon\psi_1(t,x,v) + O(\varepsilon^2), \tag{1.4}$$

with $\int_V T_\varepsilon(t,x,v)\,dv = |V|$.

The chemoattractant and nutrient concentrations satisfy the reaction–diffusion equations

$$\partial_t S = D_S\partial_{xx}S - \alpha S + \beta\rho, \tag{1.5a}$$
$$\partial_t N = D_N\partial_{xx}N - \gamma\rho N, \tag{1.5b}$$

where

$$\rho(t,x) = \int_V f(t,x,v)\,dv\ ,$$

denotes the macroscopic bacterial density. Here the chemoattractant is produced by the bacteria at rate β and naturally degraded at rate α, while the nutrient is consumed at rate γ.

The main objective of this paper is to construct and validate a numerical scheme that accurately captures the dynamics of the kinetic model for all values of the scaling parameter ε, and recovers the corresponding macroscopic drift–diffusion system in the diffusive limit $\varepsilon \to 0$. In particular, we investigate whether the traveling pulses observed at the macroscopic level are consistently reproduced by the kinetic description. To this end, we develop an asymptotic preserving (AP) scheme [8, 11], which remains stable as $\varepsilon \to 0$ and, for fixed discretization parameters, automatically reduces to a consistent discretization of the limiting macroscopic equations. Our approach builds on the micro–macro decomposition introduced in [13] (see Section 3). Other AP approaches for kinetic chemotaxis models can be found in [4, 14, 5] and the references therein.

In the diffusive limit, the kinetic model formally converges to the macroscopic system

$$\partial_t\rho = D_\rho\Delta\rho - \nabla\cdot\big(\rho u_S + \rho u_N\big), \tag{1.6a}$$
$$\partial_t S = D_S\Delta S - \alpha S + \beta\rho, \tag{1.6b}$$
$$\partial_t N = D_N\Delta N - \gamma\rho N, \tag{1.6c}$$

where the drift velocities u_S and u_N are derived from the microscopic response function and take the form

$$u_S = \chi_S J_\phi(-\varepsilon\partial_t S, |\nabla S|)\frac{\nabla S}{|\nabla S|}, \qquad u_N = \chi_N J_\phi(-\varepsilon\partial_t N, |\nabla N|)\frac{\nabla N}{|\nabla N|}. \tag{1.7}$$

Here ϕ is the so-called response function (Section 2), which describes how a single bacterium reacts to changes in the concentration of the chemoattractant S (or of the nutrient N) along its trajectory. The function J_ϕ encodes, at the macroscopic level, these microscopic features of the individual response. The parameter ε is the ratio between the pulse speed and the speed of individual bacteria; in the experiments $\varepsilon \approx 0.1$, so the pulse travels one order of magnitude slower than each individual.

In contrast with the classical Keller–Segel fluxes, these drift terms remain uniformly bounded. This provides a more realistic description of chemotactic motion and allows an analytical characterization of traveling pulses [2]: the flux u_S aggregates the cells into a pulse, while u_N drives this pulse toward higher nutrient concentrations. In the stiff-response limit, where ϕ is a step function, the chemotactic drift reduces to $\chi\,\mathrm{sign}(\partial_x S)$. For this case, in [3] the authors studied the one-dimensional model without nutrient and with S in quasi-stationary equilibrium, and proved the nonlinear stability of its stationary states, which are exponentially decaying clusters, i.e. zero-speed pulses.

The paper is organized as follows. Section 2 derives the macroscopic drift–diffusion system from the underlying kinetic chemotaxis model and recalls the theoretical traveling wave speed. Section 3 introduces the micro–macro decomposition used to construct the asymptotic preserving formulation. In Section 4, a fully discrete AP finite difference scheme is developed, and its asymptotic preserving, positivity, and mass conservation properties are analyzed. Section 5 presents numerical experiments validating the traveling pulse dynamics, the dependence of the wave speed on the chemotactic sensitivities, and the convergence of the kinetic model towards its macroscopic limit as the scaling parameter tends to zero. Finally, Section 6 summarizes the main results and discusses possible directions for future work.

2 Macroscopic system of equations

In this section we derive the macroscopic system that is obtained from (1.2) following [17]. We introduce two relevant parameters: the maximum speed of a single bacterium $c = \max\{|v| : v \in V\}$ and the mean turning frequency λ_0. The main unknown is the speed of the traveling pulse, denoted by σ. We start by rescaling the kinetic model (1.2) into a non-dimensional form using the non-dimensional variables, denoted with tildes, as

$$\tilde{t} = \frac{t}{\bar{t}}\,, \quad \tilde{x} = \frac{x}{\bar{x}}\,, \quad \tilde{V} = \frac{V}{c}\,, \quad \tilde{T} = \frac{T}{\lambda_0}\,,$$

to model the traveling wave with $\bar{x} = \sigma\bar{t}$. According to the experiments, the pulse speed σ is much lower than the speed of a single bacterium c, and this difference is measured by the ratio $\varepsilon = \sigma/c$, where $\varepsilon \ll 1$.

The kinetic equation can be written in the following form, where we drop the tilde for simplicity of notation

$$\varepsilon\partial_t f^\varepsilon + v\partial_x f^\varepsilon = \frac{\mu}{\varepsilon}\Big[\int_V T_\varepsilon(t,x,v')f^\varepsilon(t,x,v')\,\mathrm{d}v' - |V|T_\varepsilon(t,x,v)f^\varepsilon(t,x,v)\Big]\,, \tag{2.1}$$

where $\mu/\varepsilon = \lambda_0 \bar{x}/c$ and with velocity space $V = [-1, 1]$ (one-dimensional case). Here

$$T_\varepsilon(t, x, v) = 1 + \varepsilon \psi_1(t, x, v). \tag{2.2}$$

is the first-order approximation of (1.4) and ψ_1 denotes the centered first-order perturbation defined in (1.3). We consider a regular expansion of f^ε in ε up to second order

$$f^\epsilon(t, x, v) = f^0(t, x, v) + \varepsilon f^1(t, x, v) + \varepsilon^2 f^2(t, x, v) + \mathcal{O}(\varepsilon^3) \ ,$$

where $\int_V f^\epsilon(t, x, v)\, \mathrm{d}v = \int_V f^0(t, x, v)\, \mathrm{d}v = \rho(t, x)$, therefore $\int_V f^i(t, x, v)\, \mathrm{d}v = 0$, $\forall i \geq 1$.

Substituting this ansatz into (2.1) and collecting powers of ε

$$\varepsilon^0 : \quad f^0(t, x, v) = \frac{\rho(t, x)}{|V|} \ , \tag{2.3}$$

$$\varepsilon^1 : \quad v\partial_x f^0 = \mu \left(\int_V \psi_1(v') f^0(v')\, dv' - |V| \psi_1(v) f^0(v) - |V| f^1(v) \right) \ , \tag{2.4}$$

$$\varepsilon^2 : \quad \partial_t f^0 + v\partial_x f^1 = \mu \Big(\int_V \psi_1(v') f^1(v')\, \mathrm{d}v' - |V| \psi_1(v) f^1(v) - |V| f^2(v) \Big) \ . \tag{2.5}$$

Next, after replacing f^0 given by (2.3), we multiply (2.4) by v and we integrate again with respect to v to obtain

$$\begin{aligned} \mu |V| \int_V v f^1(t, x, v)\, \mathrm{d}v = & - \partial_x \frac{\rho(t, x)}{|V|} \int_V v^2\, \mathrm{d}v - \mu \rho(t, x) \chi_S \int_V \phi(\varepsilon \partial_t S + v \partial_x S) v\, \mathrm{d}v \\ & - \mu \rho(t, x) \chi_N \int_V \phi(\varepsilon \partial_t N + v \partial_x N) v\, \mathrm{d}v. \end{aligned} \tag{2.6}$$

We define

$$\begin{gathered} D_\rho = \frac{1}{\mu |V|^2} \int_V v^2\ \mathrm{d}v, \qquad u_S = - \frac{\chi_S}{|V|} \int_V \phi(\varepsilon \partial_t S + v \partial_x S)\, v\ \mathrm{d}v, \\ u_N = - \frac{\chi_N}{|V|} \int_V \phi(\varepsilon \partial_t N + v \partial_x N)\, v\ \mathrm{d}v. \end{gathered} \tag{2.7}$$

In the limiting case where the internal response function is $\phi(Y) = -\phi_0 \operatorname{sign}(Y)$, the flux terms reduce, up to the constant factor determined by ϕ_0 and the velocity measure, to [2]

$$u_S = \chi_S \operatorname{sign}(\partial_x S), \qquad u_N = \chi_N \operatorname{sign}(\partial_x N).$$

Finally, integrating (2.5) in v and using (2.7) we obtain

$$\partial_t \rho = D_\rho \partial_{xx} \rho - \partial_x (\rho u_S + \rho u_N) \ .$$

The diffusion coefficient is determined by the velocity distribution and the turning frequency as in (2.7). In particular, when the normalized scaling $\mu |V| = 1$ is used, this coefficient becomes

$$D_\rho = \frac{1}{|V|} \int_V v^2\, dv. \tag{2.8}$$

For example, when $V = [-1, 1]$, we have $D_\rho = \frac{1}{3}$. We refer to [6, 7] for more details of the derivation. Together with (1.5) for S and N, this system gives a traveling wave solution where the speed is given by the following theorem.

Theorem 2.1 ([2]) *There exist a speed $\sigma > 0$, and two positive values $N_- < N_+$, such that the system (1.6) admits traveling wave solutions $(\rho(z), S(z), N(z))$ for $z = x - \sigma t$ with the following boundary conditions:*

$$\lim_{|z|\to\infty} \rho = \lim_{|z|\to\infty} S = 0, \qquad \lim_{z\to-\infty} N = N_- < N_+ = \lim_{z\to+\infty} N.$$

Assume in addition that $\partial_z S$ changes sign only once, and that $\partial_z N$ is positive everywhere. Then, the speed $\sigma > 0$ is uniquely determined by the following implicit relation,

$$\chi_N - \sigma = \chi_S \frac{\sigma}{\sqrt{4D_S\alpha + \sigma^2}}.$$

The proof in [2] relies on the ansatz that the chemoattractant S has a unique maximum at $z = 0$ and that the nutrient N is increasing. This ansatz decouples system (1.6), so that one can solve explicitly for ρ and then for S; validating the ansatz a posteriori yields the implicit relation for σ stated in Theorem 2.1.

3 Micro-macro decomposition

Throughout this section, we use the scaling $\mu|V| = 1$. To construct an asymptotic preserving scheme consistent with the diffusive limit, we introduce the (constant) equilibrium distribution

$$M(v) = \frac{1}{|V|}, \qquad \int_V M(v)\, dv = 1.$$

Since the leading-order term is given by $f^0 = \rho/|V|$, we decompose the kinetic density as

$$f^\varepsilon(t, x, v) = \frac{\rho^\varepsilon(t, x)}{|V|} + \varepsilon g^\varepsilon(t, x, v), \qquad \int_V g^\varepsilon(t, x, v)\, dv = 0. \tag{3.1}$$

We denote by $\langle\cdot\rangle$ the integration with respect to v over V, namely

$$\langle h\rangle = \int_V h(v)\, dv,$$

and define the projection operator $\Pi h = \langle h\rangle M$. Then $(I - \Pi)g^\varepsilon = g^\varepsilon$, where I is the identity.

Integrating (2.1) with respect to v, and using the conservation property of the T_ε operator together with $\langle vM\rangle = \frac{1}{|V|}\int_V v\, dv = 0$, we obtain the macroscopic equation

$$\partial_t \rho^\varepsilon + \partial_x \langle v g^\varepsilon\rangle = 0.$$

We define the operator

$$\mathcal{Q}_\varepsilon(f) = \int_V T_\varepsilon(t,x,v')f(v')\,dv' - |V|T_\varepsilon(t,x,v)f(v),$$

and using (3.1) we obtain

$$(I-\Pi)\mathcal{Q}_\varepsilon(f^\varepsilon) = -\varepsilon\big(|V|g^\varepsilon + \rho^\varepsilon\psi_1\big) - \varepsilon^2|V|(I-\Pi)(\psi_1 g^\varepsilon).$$

Applying $I-\Pi$ to the kinetic equation (2.1) with $\mathcal{Q}_\varepsilon$ gives

$$\varepsilon^2\partial_t g^\varepsilon + \varepsilon(I-\Pi)(v\partial_x g^\varepsilon) + vM\,\partial_x\rho^\varepsilon = -\mu\big(|V|g^\varepsilon + \rho^\varepsilon\psi_1\big) - \mu\varepsilon|V|(I-\Pi)(\psi_1 g^\varepsilon).$$

Using $M = 1/|V|$ together with the normalization $\mu|V| = 1$, we obtain the micro-macro system

$$\begin{cases} \partial_t\rho^\varepsilon + \partial_x\langle v g^\varepsilon\rangle = 0, \\ \partial_t g^\varepsilon + \dfrac{1}{\varepsilon}(I-\Pi)(v\partial_x g^\varepsilon) = \dfrac{1}{\varepsilon^2}\left(-\dfrac{v}{|V|}\partial_x\rho^\varepsilon - g^\varepsilon - \dfrac{\rho^\varepsilon}{|V|}\psi_1\right) - \dfrac{1}{\varepsilon}(I-\Pi)(\psi_1 g^\varepsilon). \end{cases} \tag{3.2}$$

4 An asymptotic preserving finite difference scheme

In this section we construct an asymptotic preserving finite difference scheme for the micro-macro formulation (3.2). We present the method in one space dimension. The scheme is designed to remain stable as $\varepsilon \to 0$ and to reduce to a consistent discretization of the macroscopic drift-diffusion limit.

Let $\Omega = [x_{\min}, x_{\max}]$ be the computational domain and $x_{j+\frac{1}{2}} = x_{\min} + (j+\frac{1}{2})\Delta x$, $x_j = x_{\min} + j\Delta x$, and $t_n = n\Delta t$. We approximate ρ, S, and N at the half grid points, while the microscopic unknown g is approximated at the integer grid points,

$$\rho^n_{j+\frac{1}{2}} \approx \rho(t_n, x_{j+\frac{1}{2}}), \qquad S^n_{j+\frac{1}{2}} \approx S(t_n, x_{j+\frac{1}{2}}), \qquad N^n_{j+\frac{1}{2}} \approx N(t_n, x_{j+\frac{1}{2}}),$$

and $g^n_{j,k} \approx g(t_n, x_j, v_k)$, $v_k \in V_h$ where $V_h = \{v_k\}$ is a symmetric discrete velocity set satisfying $v_{-k} = -v_k$.

For a half-grid scalar variable $q^n_{j+\frac{1}{2}}$, we use

$$\delta_t^+ q^n_{j+\frac{1}{2}} = \frac{q^{n+1}_{j+\frac{1}{2}} - q^n_{j+\frac{1}{2}}}{\Delta t}, \quad (\delta_x q)^n_j = \frac{q^n_{j+\frac{1}{2}} - q^n_{j-\frac{1}{2}}}{\Delta x}, \quad \text{and} \quad (\delta_x^2 q)^n_{j+\frac{1}{2}} = \frac{q^n_{j+\frac{3}{2}} - 2q^n_{j+\frac{1}{2}} + q^n_{j-\frac{1}{2}}}{\Delta x^2}.$$

Moreover, we define the discrete divergence for $g^n_{j,k}$ by

$$(\delta_x g)^n_{j+\frac{1}{2},k} = \frac{g^n_{j+1,k} - g^n_{j,k}}{\Delta x}.$$

Whenever a half-grid variable is needed at an integer grid point, we use the local average

$$q_j^n = \frac{1}{2}\left(q_{j-\frac{1}{2}}^n + q_{j+\frac{1}{2}}^n\right).$$

The value $(\delta_x^2 q)_j^n$ is defined in the same way by averaging the two neighboring half-grid values.

We define

$$\langle \eta \rangle_h = \sum_k \omega_k \eta_k, \qquad \omega_k > 0, \qquad |V|_h = \sum_k \omega_k, \qquad \Pi_h \eta = \frac{\langle \eta \rangle_h}{|V|_h}.$$

We assume $\langle v \rangle_h = 0$. Then $I - \Pi_h$ is the projection onto the discrete zero-average space.

Discrete response functions At each time step, the response functions are evaluated explicitly. For $q \in \{S, N\}$, we use the scaled derivative

$$\mathcal{D}_\varepsilon^v q = \varepsilon \partial_t q + v \partial_x q. \tag{4.1}$$

We approximate $\partial_t q$ using the equations for S and N at time t^n. We define

$$\mathcal{L}_{S,j}^n = D_S(\delta_x^2 S)_j^n - \alpha S_j^n + \beta \rho_j^n, \quad \mathcal{L}_{N,j}^n = D_N(\delta_x^2 N)_j^n - \gamma \rho_j^n N_j^n.$$

The discrete response arguments are

$$\mathcal{D}_{\varepsilon,h}^{v_k} S_j^n = \varepsilon \mathcal{L}_{S,j}^n + v_k (\delta_x S)_j^n, \qquad \mathcal{D}_{\varepsilon,h}^{v_k} N_j^n = \varepsilon \mathcal{L}_{N,j}^n + v_k (\delta_x N)_j^n.$$

All quantities on the right-hand sides are known at time t^n.

The velocity average of $\phi(a + bv)$ does not vanish in general, even when ϕ is odd. We therefore use the normalized discrete responses

$$\psi_{S,j,k}^n = \chi_S \left[\phi\left(\mathcal{D}_{\varepsilon,h}^{v_k} S_j^n\right) - \Pi_h \phi\left(\mathcal{D}_{\varepsilon,h}^{v_k} S_j^n\right)\right], \psi_{N,j,k}^n = \chi_N \left[\phi\left(\mathcal{D}_{\varepsilon,h}^{v_k} N_j^n\right) - \Pi_h \phi\left(\mathcal{D}_{\varepsilon,h}^{v_k} N_j^n\right)\right].$$

We set $\psi_{j,k}^n = \psi_{S,j,k}^n + \psi_{N,j,k}^n$, and by construction,

$$\langle \psi_{S,j,k}^n \rangle_h = 0, \qquad \langle \psi_{N,j,k}^n \rangle_h = 0, \qquad \langle \psi_{j,k}^n \rangle_h = 0.$$

Consequently, the discrete turning kernel $T_{\varepsilon,j,k}^n = 1 + \varepsilon \psi_{j,k}^n$ satisfies $\langle T_{\varepsilon,j,k}^n \rangle_h = |V|_h$. We also introduce $B_{S,j}^n = \langle v_k \psi_{S,j,k}^n \rangle_h$, $B_{N,j}^n = \langle v_k \psi_{N,j,k}^n \rangle_h$. Then, the corresponding discrete drift velocities are defined by

$$u_{S,j}^n = -\frac{B_{S,j}^n}{|V|_h}, \qquad u_{N,j}^n = -\frac{B_{N,j}^n}{|V|_h}.$$

Fully discrete micro-macro scheme The fully discrete AP scheme is given by

$$\delta_t^+ \rho_{j+\frac{1}{2}}^n + \left\langle v_k (\delta_x g)_{j+\frac{1}{2},k}^{n+1} \right\rangle_h = 0 \ , \tag{4.2a}$$

$$\delta_t^+ g_{j,k}^n + \frac{1}{\varepsilon}(I - \Pi_h)\mathcal{A}_{j,k}^n = \frac{1}{\varepsilon^2}\left(K_{j,k}^{n,n+1} - g_{j,k}^{n+1}\right) - \frac{1}{\varepsilon}(I - \Pi_h)(\psi_{j,k}^n g_{j,k}^n) \ , \tag{4.2b}$$

$$\delta_t^+ S_{j+1/2}^n = D_S(\delta_x^2 S^{n+1})_{j+1/2} - \alpha S_{j+1/2}^{n+1} + \beta \rho_{j+1/2}^{n+1} \ , \tag{4.2c}$$

$$\delta_t^+ N_{j+\frac{1}{2}}^n = D_N(\delta_x^2 N^{n+1})_{j+1/2} - \gamma \rho_{j+1/2}^{n+1} N_{j+1/2}^{n+1} \ . \tag{4.2d}$$

Here the microscopic transport is treated explicitly by the upwind approximation

$$\mathcal{A}_{j,k}^n = v_k^+ (\delta_x g)_{j-\frac{1}{2},k}^n - v_k^- (\delta_x g)_{j+\frac{1}{2},k}^n,$$

where $v_k^+ = \max\{v_k, 0\},\ v_k^- = \max\{-v_k, 0\}$. The local equilibrium part is defined by

$$K_{j,k}^{n,n+1} = -\frac{v_k}{|V|_h}(\delta_x \rho)_j^{n+1} - \frac{\rho_j^{[S],n+1}}{|V|_h}\psi_{S,j,k}^n - \frac{\rho_j^{[N],n+1}}{|V|_h}\psi_{N,j,k}^n. \tag{4.3}$$

The two upwind values $\rho_j^{[S],n+1}$ and $\rho_j^{[N],n+1}$ are chosen according to the effective velocities $u_{S,j}^n$ and $u_{N,j}^n$,

$$\rho_j^{[S],n+1} = \begin{cases} \rho_{j-\frac{1}{2}}^{n+1}, & u_{S,j}^n \geq 0, \\ \rho_{j+\frac{1}{2}}^{n+1}, & u_{S,j}^n < 0, \end{cases} \quad \rho_j^{[N],n+1} = \begin{cases} \rho_{j-\frac{1}{2}}^{n+1}, & u_{N,j}^n \geq 0, \\ \rho_{j+\frac{1}{2}}^{n+1}, & u_{N,j}^n < 0. \end{cases} \tag{4.4}$$

This choice is consistent with the drift part of the limiting flux since

$$-\frac{B_{S,j}^n}{|V|_h}\rho_j^{[S],n+1} - \frac{B_{N,j}^n}{|V|_h}\rho_j^{[N],n+1} = u_{S,j}^n \rho_j^{[S],n+1} + u_{N,j}^n \rho_j^{[N],n+1}.$$

Taking the discrete velocity average of the second equation in (4.2) and using $\langle K_{j,k}^{n,n+1} \rangle_h = 0$ gives

$$\langle g_{j,k}^{n+1} \rangle_h = \frac{\varepsilon^2}{\varepsilon^2 + \Delta t} \langle g_{j,k}^n \rangle_h.$$

Thus $\langle g_{j,k}^n \rangle_h = 0$ is propagated in time if it holds initially.

Decoupled solution strategy Following the micro-macro strategy in [13], the scheme can be solved in a decoupled way. We first introduce an auxiliary unknown $\widetilde{g}_{j,k}^{n+1}$ by

$$\frac{\widetilde{g}_{j,k}^{n+1} - g_{j,k}^n}{\Delta t} + \frac{1}{\varepsilon}(I - \Pi_h)\mathcal{A}_{j,k}^n = \frac{1}{\varepsilon^2}\left(\widetilde{K}_{j,k}^n - \widetilde{g}_{j,k}^{n+1}\right) - \frac{1}{\varepsilon}(I - \Pi_h)(\psi_{j,k}^n g_{j,k}^n). \tag{4.5}$$

where

$$\widetilde{K}^n_{j,k} = -\frac{v_k}{|V|_h}(\delta_x \rho)^n_j - \frac{\rho^{[S],n}_j}{|V|_h}\psi^n_{S,j,k} - \frac{\rho^{[N],n}_j}{|V|_h}\psi^n_{N,j,k}.$$

The upwind values $\rho^{[S],n}_j$ and $\rho^{[N],n}_j$ are defined in the same way as in (4.4), with ρ^{n+1} replaced by ρ^n.

All terms in (4.5) are known except $\widetilde{g}^{n+1}_{j,k}$. Hence

$$\widetilde{g}^{n+1}_{j,k} = \frac{\varepsilon^2}{\varepsilon^2+\Delta t} g^n_{j,k} - \frac{\varepsilon \Delta t}{\varepsilon^2+\Delta t}(I-\Pi_h)\mathcal{A}^n_{j,k} + \frac{\Delta t}{\varepsilon^2+\Delta t}\widetilde{K}^n_{j,k} - \frac{\varepsilon \Delta t}{\varepsilon^2+\Delta t}(I-\Pi_h)(\psi^n_{j,k} g^n_{j,k}).$$

Comparing (4.5) with (4.2b), we obtain

$$g^{n+1}_{j,k} = \widetilde{g}^{n+1}_{j,k} + \lambda_\varepsilon \left(K^{n,n+1}_{j,k} - \widetilde{K}^n_{j,k} \right), \tag{4.6}$$

where

$$\lambda_\varepsilon = \frac{\Delta t}{\varepsilon^2 + \Delta t}. \tag{4.7}$$

Substituting (4.6) into (4.2a), we obtain a closed equation for ρ^{n+1}. Define

$$D_{\rho,h} = \frac{\langle v_k^2 \rangle_h}{|V|_h}, \qquad b^n_{S,j} = \frac{B^n_{S,j}}{|V|_h}, \qquad b^n_{N,j} = \frac{B^n_{N,j}}{|V|_h},$$

and let $a^n = \lambda_\varepsilon D_{\rho,h}$, $c^n_{S,j} = \lambda_\varepsilon b^n_{S,j}$, $c^n_{N,j} = \lambda_\varepsilon b^n_{N,j}$. Then the density equation can be written as

$$\frac{\rho^{n+1}_{j+\frac{1}{2}}}{\Delta t} - \delta_x \left(a^n \delta_x \rho^{n+1} \right)_{j+\frac{1}{2}} - \delta_x \left(c^n_S \rho^{[S],n+1} \right)_{j+\frac{1}{2}} - \delta_x \left(c^n_N \rho^{[N],n+1} \right)_{j+\frac{1}{2}} = r^n_{j+\frac{1}{2}}, \tag{4.8}$$

where the known right-hand side is

$$r^n_{j+\frac{1}{2}} = \frac{\rho^n_{j+\frac{1}{2}}}{\Delta t} - \left\langle v_k (\delta_x \widetilde{g})^{n+1}_{j+\frac{1}{2},k} \right\rangle_h - \delta_x \left(a^n \delta_x \rho^n \right)_{j+\frac{1}{2}} - \delta_x \left(c^n_S \rho^{[S],n} \right)_{j+\frac{1}{2}} - \delta_x \left(c^n_N \rho^{[N],n} \right)_{j+\frac{1}{2}}.$$

Thus only a scalar linear system for ρ^{n+1} needs to be solved at each time step. With the upwind choice for the drift terms (4.4), the resulting matrix is tridiagonal. Once ρ^{n+1} is obtained, g^{n+1} is recovered explicitly from (4.6). We impose zero numerical density flux at both boundaries. This condition is preserved by the positivity correction described in Appendix A. The corrected density is then used to update S^{n+1} and N^{n+1} independently by (4.2c) and (4.2d), respectively.

4.1 Asymptotic preserving property

We now verify that the scheme is formally asymptotic preserving. Let Δt and Δx be fixed. Notice that λ_ε in (4.7) converges to 1 as ε goes to zero. Moreover, by the definition of $\widetilde{g}^{n+1}$, $\widetilde{g}^{n+1}_{j,k}$ converges to $\widetilde{K}^n_{j,k}$. Using the expression of $\widetilde{K}^n$, we obtain $\left\langle v_k(\delta_x \widetilde{g})^{n+1}_{j+\frac{1}{2},k} \right\rangle_h$ converges to

$$-\delta_x \left(D_{\rho,h}\delta_x \rho^n\right)_{j+\frac{1}{2}} - \delta_x \left(b_S^n \rho^{[S],n}\right)_{j+\frac{1}{2}} - \delta_x \left(b_N^n \rho^{[N],n}\right)_{j+\frac{1}{2}} .$$

It follows from the definition of $r^n_{j+\frac{1}{2}}$ in (4.8) that $r^n_{j+\frac{1}{2}}$ converges to $\frac{\rho^n_{j+\frac{1}{2}}}{\Delta t}$. Therefore, in the limit as ε goes to zero, (4.8) reduces to

$$\frac{\rho^{n+1}_{j+\frac{1}{2}} - \rho^n_{j+\frac{1}{2}}}{\Delta t} - \delta_x \left(D_{\rho,h}\delta_x \rho^{n+1}\right)_{j+\frac{1}{2}} - \delta_x \left(b_S^n \rho^{[S],n+1}\right)_{j+\frac{1}{2}} - \delta_x \left(b_N^n \rho^{[N],n+1}\right)_{j+\frac{1}{2}} = 0. \quad (4.9)$$

This is a consistent implicit finite difference discretization of the limiting drift-diffusion equation

$$\partial_t \rho - \partial_x \left(D_{\rho,h}\partial_x \rho\right) - \partial_x \left(b_S \rho\right) - \partial_x \left(b_N \rho\right) = 0 \ ,$$

where $u_S = -b_S$ and $u_N = -b_N$. Hence, for fixed Δt and Δx, the fully discrete scheme reduces to a consistent discretization of the macroscopic drift–diffusion limit as $\varepsilon \to 0$, which is the formal AP property in the diffusive regime.

4.2 Bound-preserving property

We now discuss the bound-preserving property of the limiting scheme. In general, the full kinetic scheme is not automatically positivity preserving. In the computation, this issue is handled by a post-processing limiter as explained in Appendix A. For the limiting macroscopic scheme, however, a positivity and boundedness result can be established.

In this subsection, we consider the limiting macroscopic scheme (4.9) with homogeneous no-flux boundary conditions, so that the discrete boundary fluxes vanish.

The limiting scheme can be written in matrix form as

$$M^n \boldsymbol{\rho}^{n+1} = \boldsymbol{\rho}^n \ ,$$

where $\boldsymbol{\rho}^n = (\rho^n_{j+\frac{1}{2}})_j$ denotes the vector of half-grid density values at time level n. The tridiagonal matrix M^n has entries

$$\begin{aligned}
m^n_{j,j-1} &= -D_{\rho,h}\frac{\Delta t}{(\Delta x)^2} - \frac{\Delta t}{\Delta x}(u^n_{S,j})^+ - \frac{\Delta t}{\Delta x}(u^n_{N,j})^+, \\
m^n_{j,j} &= 1 + 2D_{\rho,h}\frac{\Delta t}{(\Delta x)^2} + \frac{\Delta t}{\Delta x}\left[(u^n_{S,j+1})^+ + (u^n_{S,j})^-\right] + \frac{\Delta t}{\Delta x}\left[(u^n_{N,j+1})^+ + (u^n_{N,j})^-\right], \\
m^n_{j,j+1} &= -D_{\rho,h}\frac{\Delta t}{(\Delta x)^2} - \frac{\Delta t}{\Delta x}(u^n_{S,j+1})^- - \frac{\Delta t}{\Delta x}(u^n_{N,j+1})^-.
\end{aligned}$$

Here $(a)^+ = \max\{a, 0\}$ and $(a)^- = \max\{-a, 0\}$. The boundary rows of M^n are modified according to the homogeneous no-flux boundary conditions.

Proposition 4.1 (Mass conservation, positivity, and upper bound) *Assume that $D_{\rho,h} > 0$ and that the effective velocities satisfy*

$$|u^n_{S,j}| \le U_S, \qquad |u^n_{N,j}| \le U_N$$

for all j and n. Suppose further that

$$\frac{\Delta t}{\Delta x} \le \frac{1}{4(U_S + U_N)}. \tag{4.10}$$

If the initial data satisfy $\rho^0_{j+\frac{1}{2}} \ge 0$ for all j, then the limiting scheme satisfies

$$\rho^n_{j+\frac{1}{2}} \ge 0, \qquad \forall j,\ n \ge 0.$$

Moreover, under the homogeneous no-flux boundary conditions, the discrete mass is conserved,

$$\|\boldsymbol{\rho}^{n+1}\|_1 = \|\boldsymbol{\rho}^n\|_1 := \sum_j |\rho^n_{j+\frac{1}{2}}|,$$

and the solution satisfies the uniform bound in time on a fixed spatial grid

$$0 \le \rho^n_{j+\frac{1}{2}} \le \bar{\rho}, \qquad \bar{\rho} = 2\|\boldsymbol{\rho}^0\|_1.$$

Proof.- We first observe that

$$m^n_{j,j-1} + m^n_{j,j} + m^n_{j,j+1} = 1 + \frac{\Delta t}{\Delta x}\left(u^n_{S,j+1} - u^n_{S,j}\right) + \frac{\Delta t}{\Delta x}\left(u^n_{N,j+1} - u^n_{N,j}\right) =: F^n_j.$$

Using the bounds on u^n_S and u^n_N, we have

$$F^n_j \ge 1 - 2(U_S + U_N)\frac{\Delta t}{\Delta x}.$$

Under (4.10), this gives $F^n_j \ge \frac{1}{2}$.

The off-diagonal entries of M^n are nonpositive. Moreover, the above lower bound implies the required diagonal dominance under the CFL condition. Hence M^n is a nonsingular M-matrix and $(M^n)^{-1}$ is nonnegative. Therefore, if $\boldsymbol{\rho}^n \ge 0$, then

$$\boldsymbol{\rho}^{n+1} = (M^n)^{-1}\boldsymbol{\rho}^n \ge 0.$$

Mass conservation follows from the conservative form of the scheme and the homogeneous no-flux boundary conditions. More precisely,

$$\|\boldsymbol{\rho}^{n+1}\|_1 = \mathbf{1}^T\boldsymbol{\rho}^{n+1} = \mathbf{1}^T M^n \boldsymbol{\rho}^{n+1} = \mathbf{1}^T\boldsymbol{\rho}^n = \|\boldsymbol{\rho}^n\|_1,$$

where we used the relation $\mathbf{1}^T M^n = \mathbf{1}^T$.

Finally, let $\bar{\rho} = 2\|\boldsymbol{\rho}^0\|_1$. By mass conservation and positivity,

$$\rho^n_{j+\frac{1}{2}} \le \|\boldsymbol{\rho}^n\|_1 = \|\boldsymbol{\rho}^0\|_1 = \frac{1}{2}\bar{\rho}.$$

Since $F^n_j \ge 1/2$, we obtain

$$(M^n(\bar{\rho}\mathbf{1}))_j = F^n_j \bar{\rho} \ge \frac{1}{2}\bar{\rho} \ge \rho^n_{j+\frac{1}{2}}.$$

Using the nonnegativity of $(M^n)^{-1}$, we conclude that

$$\boldsymbol{\rho}^{n+1} = (M^n)^{-1}\boldsymbol{\rho}^n \le (M^n)^{-1}M^n(\bar{\rho}\mathbf{1}) = \bar{\rho}\mathbf{1}.$$

This proves the desired upper bound.

5 Numerical experiments

In this section, we present numerical experiments to illustrate the traveling pulse behavior of the model and to verify the asymptotic preserving property of the proposed micro-macro scheme. We first validate the macroscopic solver by comparing the numerical traveling pulse with the analytical traveling profile. We then study the dependence of the traveling speed on the chemotactic sensitivities. Finally, we compare the kinetic and macroscopic solutions for different values of ε.

Unless otherwise stated, we consider the one-dimensional computational domain $\Omega = (0, 150)$. For the profile and asymptotic preserving tests, we use $\Delta x = 0.025$, $\Delta t = 10^{-3}$. The velocity domain is $V = [-1, 1]$. It is discretized by a uniform trapezoidal quadrature with $\Delta v = 0.02$. We denote the discrete velocity nodes and weights by $\{v_k, \omega_k\}$, and use

$$\langle h\rangle_h = \sum_k \omega_k h(v_k), \qquad |V|_h = \sum_k \omega_k.$$

The discrete diffusion coefficient in the limiting macroscopic equation is $D_{\rho,h} = \frac{\langle v^2\rangle_h}{|V|_h}$. For the default velocity grid with $\Delta v = 0.02$, this gives $D_{\rho,h} = 0.3334$.

We use the reflecting boundary conditions described in Appendix A for the kinetic variable and homogeneous Neumann conditions for the chemical variables.

The initial data are chosen so that cells are concentrated near the left side of the domain, with no chemoattractant signal and a constant nutrient level. More precisely, we take

$$\rho(0, x) = \rho_0 e^{-\lambda x}, \qquad S(0, x) = 0, \qquad N(0, x) = N_0,$$

where $\rho_0 = 1,\ \lambda = 3,\ N_0 = 10^3$. For the kinetic computations, the initial data are chosen consistently with the micro-macro decomposition,

$$f^\varepsilon(0, x, v) = \frac{\rho(0, x)}{|V|_h} + \varepsilon g(0, x, v), \qquad g(0, x, v) = 0.$$

The chemical parameters are $D_S = 2,\ D_N = 0,\ \alpha = 0.05,\ \beta = \gamma = 1$. Unless otherwise stated, the chemotactic sensitivities in the profile and AP tests are $\chi_S = 0.5,\ \chi_N = 1.1$. For the experiments in Sections 5.1–5.3, we use

$$\phi_h(y) = -s_h \operatorname{sign}(y), \qquad s_h = \frac{|V|_h}{\langle |v| \rangle_h}.$$

With this normalization, the macroscopic drift velocities are $u_S = \chi_S \operatorname{sign}(S_x)$ and $u_N = \chi_N \operatorname{sign}(N_x)$. The response arguments are evaluated with $\varepsilon = 0$ for the macroscopic model and with $\varepsilon > 0$ for the kinetic model. The kinetic computations use the correction described in Appendix A.

5.1 Macroscopic traveling pulse and analytical profile

We first validate the macroscopic solver by comparing the numerical traveling pulse with the analytical traveling profile. Let x_p be the peak position of the numerical density and set

$$z = x - x_p, \qquad z_- = \min\{z, 0\}, \qquad z_+ = \max\{z, 0\}.$$

The analytical traveling profile with the response function $\phi(y) = -\operatorname{sign}(y)$ found in [2] is written as

$$\rho_{\mathrm{tw}}(x) = \rho_p \exp\left(\lambda_L z_- + \lambda_R z_+\right), \tag{5.1}$$

where

$$\lambda_L = \frac{\chi_S + \chi_N - \sigma}{D_{\rho,h}}, \qquad \lambda_R = \frac{\chi_N - \chi_S - \sigma}{D_{\rho,h}}.$$

Here $\lambda_L > 0$ and $\lambda_R < 0$, so that the profile decays on both sides of the peak. The amplitude ρ_p is chosen by discrete mass matching. More precisely, if $z_j = x_{j+\frac{1}{2}} - x_p$, then


$$\rho_p = \frac{\Delta x \sum_j \rho_j}{\Delta x \sum_j \exp\left(\lambda_L (z_j)_- + \lambda_R (z_j)_+\right)}.$$

Figure 1 compares the macroscopic numerical profiles with the analytical traveling profiles at two representative times. The peak position and amplitude of the analytical profile are chosen from the corresponding numerical solution. The close agreement confirms that the macroscopic solver captures the expected traveling wave structure. This macroscopic solution will be used as the reference solution in the kinetic-to-macroscopic comparison below.

5.2 Dependence of the traveling speed on chemotactic sensitivities

We next investigate how the traveling speed depends on χ_S and χ_N. For each pair (χ_S, χ_N), the predicted traveling speed σ is determined by

$$\chi_N - \sigma = \chi_S \frac{\sigma}{\sqrt{4 D_S \alpha + \sigma^2}}. \tag{5.2}$$



(a) $t = 30$ (b) $t = 50$

Figure 1: Comparison between the macroscopic numerical profiles and the analytical traveling profiles at $t = 30$ and $t = 50$.

To estimate the traveling speed numerically, we track the center of mass of the density,


$$X_\rho(t_n) = \frac{\Delta x \sum_j x_{j+\frac{1}{2}} \rho^n_{j+\frac{1}{2}}}{\Delta x \sum_j \rho^n_{j+\frac{1}{2}}}. \tag{5.3}$$

For the parameter study, we use a slightly coarser mesh to reduce the computational cost, i.e. $\Delta x = 0.1, \Delta v = 0.02, \Delta t = 10^{-2}$. The sensitivities χ_S, χ_N are uniformly sampled on $[0, 2] \times [0, 2]$. The computational domain for this parameter study is $\Omega = (0, 300)$. The wave speeds are evaluated at $t = 100$.

Figure 2 shows the dependence of the traveling speed on χ_S and χ_N. The subfigure (a) gives the speed predicted by (5.2), while the subfigures (b) and (c) show the numerical speed obtained from the macroscopic and the kinetic model with $\varepsilon = 0.01$, respectively, by tracking the center of mass. The three panels agree well in the parameter regime where a clear traveling pulse is formed. Some discrepancies are observed near the transition region where the nutrient bias is not sufficiently strong compared with the self-attractant bias. In this region, the pulse formation may require a longer transient time, and the finite-time center-of-mass estimate may underestimate the asymptotic traveling speed.

5.3 Asymptotic preserving convergence

We now compare the kinetic solution with the limiting macroscopic solution for different values of ε. The purpose of this test is to verify that, for fixed Δx, Δv, and Δt, the kinetic AP solution approaches the limiting macroscopic solution as ε decreases. We use the baseline parameter setting described at the beginning of this section and consider $\varepsilon = 0.1,\ \varepsilon = 0.05,\ \varepsilon = 0.01$.



(a) (b) (c)

Figure 2: Dependence of the traveling speed on χ_S and χ_N. Figure (a) shows the speed predicted by (5.2), while Figure (b) and Figure (c) show the numerical speed obtained from the macroscopic and kinetic model with $\varepsilon = 0.01$, respectively.

Figure 3 compares the traveling density pulses obtained from the kinetic model and the limiting macroscopic model at several representative times. As ε decreases, the kinetic profiles become progressively closer to the macroscopic profile. In particular, for $\varepsilon = 0.01$, the kinetic and macroscopic solutions are already nearly indistinguishable at the plotted times. This provides direct evidence that the AP scheme captures the correct diffusive limit.

To quantify the convergence, we use the discrete L^2-error

$$\mathrm{err}_{\rho,\varepsilon}(t) = \left(\Delta x \sum_j \left| \rho^{\varepsilon}_{j+\frac{1}{2}}(t) - \rho_{j+\frac{1}{2}}(t) \right|^2 \right)^{1/2},$$

where ρ^{ε} is reconstructed from the kinetic solution and ρ denotes the solution of the limiting macroscopic model. Figure 4 plots the final-time error $\mathrm{err}_{\rho,\varepsilon}(50)$ for different values of ε. The error decreases monotonically as ε becomes smaller, confirming the convergence of the kinetic AP solution toward the macroscopic one.

Besides the density profile, it is also important to compare the traveling speed. The instantaneous speed is computed from the center-of-mass trajectory $X_\rho(t)$ introduced in (5.3). Figure 5 shows the speeds obtained from the macroscopic solution and from the kinetic solutions for different values of ε. The horizontal dashed line indicates the predicted value $\sigma_{\mathrm{pred}} = 0.724$, obtained from (5.2). After a short transient regime, the propagation speeds become approximately constant and remain close to the predicted value. Moreover, the kinetic wave speed becomes closer to the macroscopic one as ε decreases.

5.4 Traveling pulses with an exponential response

We next consider the exponentially modulated response

$$\phi_{a,h}(y) = -s_h e^{-a|y|}\,\mathrm{sign}(y), \qquad s_h = \frac{|V|_h}{\langle |v| \rangle_h}. \tag{5.4}$$




Figure 3: Comparison of traveling density pulses obtained from the kinetic model and the limiting macroscopic model at $t = 1, 10, 20, 30, 40, 50$. The propagation of the pulse is captured at both levels, and the kinetic profiles approach the macroscopic profiles as ε decreases.




Figure 4: Final-time density error $\mathrm{err}_{\rho,\varepsilon}(50)$ versus ε. The reference lines $O(\varepsilon)$ and $O(\varepsilon^2)$ are included for comparison.




Figure 5: traveling speed comparison between the macroscopic model and the kinetic model for different values of ε. The horizontal dashed line indicates the macroscopic predicted speed $\sigma_{\text{macro}} \approx 0.724$, obtained from (5.2).

Here $a \geq 0$ controls the exponential modulation, and $a = 0$ recovers the normalized sign response. The initial data and all other parameters are the same as in Section 5.3. For each value of a, the macroscopic reference solution is computed using the same response function with $\varepsilon = 0$.

For $a = 0.001$ and $\varepsilon = 0.01$, Figure 6 shows an asymmetric pulse propagating to the right. Its propagation speed becomes approximately constant at late times. At fixed $a = 0.001$, the kinetic density profiles approach the corresponding numerical macroscopic solution as ε decreases, as shown in Figure 7(a). At fixed $\varepsilon = 0.01$, the density profiles approach the sign-response solution as a decreases in Figure 7(b). The fitted propagation speeds also approach the numerical sign-response value, as reported in Table 1.

Table 1: Numerical propagation speeds for the exponential response with $\varepsilon = 0.01$. The speeds are obtained by least-squares linear fits to the center of mass over $t \in [30, 50]$.

a	10^{-3}	5×10^{-4}	10^{-4}	0
Propagation speed	0.690	0.707	0.720	0.724

Overall, these results show that the proposed AP scheme captures the traveling pulse profiles at the kinetic level and converges to the limiting macroscopic dynamics, both in density profiles and in propagation speed, as ε decreases.




Figure 6: Density profiles for the exponential response with $a = 0.001$ and $\varepsilon = 0.01$ at $t = 1, 10, 20, 30, 40, 50$.




Figure 7: Density profiles at $T = 50$. (a) Kinetic solutions with $\varepsilon = 0.1, 0.05, 0.01$ and the corresponding numerical macroscopic solution for $a = 0.001$. (b) Solutions with $a = 0.001, 0.0005, 0.0001, 0$ for $\varepsilon = 0.01$. The case $a = 0$ corresponds to the sign response.

6 Conclusions

In this work, we developed and analyzed an asymptotic preserving finite difference scheme for a kinetic chemotaxis model describing bacterial traveling pulses. The model couples a run-and-tumble kinetic equation with reaction–diffusion equations for a nutrient and a self-produced chemoattractant, motivated by experimental observations of *E. coli* pulse propagation in microchannels.

The proposed numerical method is based on a micro–macro decomposition of the kinetic equation and remains stable in both the kinetic and diffusive regimes. In the limit as the scaling parameter $\varepsilon \to 0$, the scheme naturally reduces to a consistent discretization of the limiting drift–diffusion system. We also established positivity, mass conservation, and uniform boundedness properties for the limiting macroscopic scheme, and introduced a positivity-preserving post-processing procedure for the kinetic solver.

The numerical experiments confirm that the scheme accurately captures the formation and propagation of traveling bacterial pulses for the sign response function. In particular, the simulations reproduce the characteristic asymmetric pulse profile and validate the theoretical prediction for the traveling wave speed found in [2]. Moreover, the kinetic solutions converge toward the macroscopic limit as ε decreases, demonstrating the asymptotic preserving character of the method.

Several directions remain open for future work. A rigorous convergence analysis of the fully discrete AP scheme would be of significant interest. It would also be natural to extend the method to higher-dimensional settings and more general kinetic response functions. Another important direction concerns the development of fully positivity-preserving AP schemes without post-processing corrections. Finally, incorporating additional biological mechanisms, such as population growth, signal adaptation, or interactions between multiple bacterial species, could provide a more realistic description of collective bacterial migration phenomena.

Acknowledgements CA was supported by TÜBİTAK (The Scientific and Technological Research Council of Türkiye) through the International Postdoctoral Research Fellowship Program under Grant No: 1059B191800629. JAC was supported by the Advanced Grant Nonlocal-CPD (Nonlocal PDEs for Complex Particle Dynamics: Phase Transitions, Patterns and Synchronization) of the European Research Council Executive Agency (ERC) under the European Union Horizon 2020 research and innovation programme (grant agreement No. 883363) and partially supported by the EPSRC EP/V051121/1. JAC was partially supported by the "Maria de Maeztu" Excellence Unit IMAG, reference CEX2020-001105-M, funded by MCIN/AEI/10.13039/501100011033/. GER is supported by the Spanish grant PID2022-143012NA-100 and she is a member of the Catalan research group 2021-SGR-00087. XR acknowledges support from the R&D Program of Beijing Municipal Education Commission from China under Grant No. KM202310028016, and the National Natural Science Foundation of China under Grant No. 12201436.

A Boundary treatment and positivity preservation

We use the discrete velocity average

$$\langle h\rangle_h = \sum_k \omega_k h(v_k), \qquad \omega_k > 0, \qquad |V|_h = \sum_k \omega_k, \qquad \psi_0(v_k) = |V|_h^{-1}.$$

Let J be the number of spatial cells, and assume $v_{-k} = -v_k$ and $\omega_{-k} = \omega_k$. We impose reflection at the endpoints and use reflected ghost values,

$$g^n_{j,k} = g^n_{j,-k} \quad (j = 0, J), \qquad g^n_{-1,k} = g^n_{1,-k}, \qquad g^n_{J+1,k} = g^n_{J-1,-k}.$$

Endpoint symmetry is imposed before the kinetic update, on the auxiliary variable $\widetilde{g}^{n+1}$, and on the provisional microscopic variable before checking positivity. The diffusion and drift boundary fluxes are set to zero. Thus the complete numerical density flux vanishes at both endpoints, and the density update conserves mass. Homogeneous Neumann conditions are used for the chemical variables.

For the correction, we assume $\varepsilon > 0$, omit the superscript $n+1$, and denote the provisional solution of (4.2) by $\widehat{\rho}$ and $\widehat{g}$, with $\langle \widehat{g}_{j,\cdot}\rangle_h = 0$. Define

$$\widehat{f}_{j+\frac{1}{2},k} = \widehat{\rho}_{j+\frac{1}{2}}\psi_0(v_k) + \frac{\varepsilon}{2}\big(\widehat{g}_{j,k} + \widehat{g}_{j+1,k}\big).$$

If $\widehat{\rho}_{j+\frac{1}{2}} \geq 0$ and $\widehat{f}_{j+\frac{1}{2},k} \geq 0$ for all j, k, we set $\rho = \widehat{\rho}$ and $g = \widehat{g}$ without further correction. Otherwise, we apply the following procedure. For $\widehat{\rho}_{j+\frac{1}{2}} > 0$, set

$$\theta_{j+\frac{1}{2}} = \min\left\{1, \min_{\widehat{f}_{j+\frac{1}{2},k}<0} \frac{\widehat{\rho}_{j+\frac{1}{2}}\psi_0(v_k)}{\widehat{\rho}_{j+\frac{1}{2}}\psi_0(v_k) - \widehat{f}_{j+\frac{1}{2},k}}\right\},$$

where the inner minimum is omitted if all values are nonnegative, and

$$\rho^*_{j+\frac{1}{2}} = \widehat{\rho}_{j+\frac{1}{2}}, \qquad \bar{g}^*_{j+\frac{1}{2},k} = \frac{\theta_{j+\frac{1}{2}}}{2}\big(\widehat{g}_{j,k} + \widehat{g}_{j+1,k}\big).$$

For $\widehat{\rho}_{j+\frac{1}{2}} \leq 0$, set $\rho^*_{j+\frac{1}{2}} = \bar{g}^*_{j+\frac{1}{2},k} = 0$. Then $\rho^*_{j+\frac{1}{2}}\psi_0(v_k) + \varepsilon\bar{g}^*_{j+\frac{1}{2},k} \geq 0$ and $\langle \bar{g}^*_{j+\frac{1}{2},\cdot}\rangle_h = 0$. Assuming $\Delta x \sum_j \widehat{\rho}_{j+\frac{1}{2}} > 0$, restore the input mass by

$$\alpha = \frac{\sum_j \widehat{\rho}_{j+\frac{1}{2}}}{\sum_j \rho^*_{j+\frac{1}{2}}}, \qquad \rho_{j+\frac{1}{2}} = \alpha\rho^*_{j+\frac{1}{2}}, \qquad \bar{g}_{j+\frac{1}{2},k} = \alpha\bar{g}^*_{j+\frac{1}{2},k}.$$

We return the microscopic variable to interfaces by

$$\widehat{h}_{j,k} = \tfrac{1}{2}\big(\bar{g}_{j-\frac{1}{2},k} + \bar{g}_{j+\frac{1}{2},k}\big), \qquad 1 \leq j \leq J-1,$$
$$\widehat{h}_{0,k} = \tfrac{1}{2}\big(\bar{g}_{\frac{1}{2},k} + \bar{g}_{\frac{1}{2},-k}\big), \qquad \widehat{h}_{J,k} = \tfrac{1}{2}\big(\bar{g}_{J-\frac{1}{2},k} + \bar{g}_{J-\frac{1}{2},-k}\big),$$

and set $h_j = (I - \Pi_h)\widehat{h}_j$. Since this reconstruction need not preserve positivity, we check

$$F_{j+\frac{1}{2},k} = \rho_{j+\frac{1}{2}}\psi_0(v_k) + \frac{\varepsilon}{2}(h_{j,k} + h_{j+1,k}).$$

If all values are nonnegative, we set $g_j = h_j$. Otherwise, define

$$q_j = \frac{\min(\rho_{j-\frac{1}{2}}, \rho_{j+\frac{1}{2}})}{|V|_h} \quad (1 \le j \le J-1), \qquad q_0 = \frac{\rho_{\frac{1}{2}}}{|V|_h}, \qquad q_J = \frac{\rho_{J-\frac{1}{2}}}{|V|_h},$$

$$m_j = \min_k h_{j,k}, \qquad \eta_j = \begin{cases} \min\{1, q_j/(-\varepsilon m_j)\}, & m_j < 0, \\ 1, & m_j \ge 0, \end{cases} \qquad g_{j,k} = \eta_j h_{j,k}.$$

In this case, $\varepsilon g_{j,k} \ge -q_j$, and hence

$$f_{j+\frac{1}{2},k} = \rho_{j+\frac{1}{2}}\psi_0(v_k) + \frac{\varepsilon}{2}(g_{j,k} + g_{j+1,k}) \ge \frac{\rho_{j+\frac{1}{2}}}{|V|_h} - \frac{q_j + q_{j+1}}{2} \ge 0.$$

The interface scaling preserves the zero velocity mean and endpoint reflection and leaves the density unchanged. Together with the conservative density update, this gives

$$\Delta x \sum_j \rho_{j+\frac{1}{2}} = \Delta x \sum_j \widehat{\rho}_{j+\frac{1}{2}} = \Delta x \sum_j \rho^n_{j+\frac{1}{2}}.$$

At fixed grids, this additional interface scaling is inactive for sufficiently small ε if ρ is bounded below by a positive constant independent of ε and h remains bounded.

References

[1] W. Alt. Biased random walk models for chemotaxis and related diffusion approximations. *Journal of Mathematical Biology*, 9:147–177, 1980.

[2] V. Calvez. Chemotactic waves of bacteria at the mesoscale. *Journal of the European Mathematical Society*, 22(2):593–668, 2019.

[3] V. Calvez and F. Hoffmann. Nonlinear stability of chemotactic clustering with discontinuous advection. *Annales de la Faculté des sciences de Toulouse: Mathématiques*, 32(2):287–318, 2023.

[4] J. A. Carrillo and B. Yan. An asymptotic preserving scheme for the diffusive limit of kinetic systems for chemotaxis. *Multiscale Model. Simul.*, 11(1):336–361, 2013.

[5] A. Chertock, A. Kurganov, M. Luká˘ cová Medvi˘ dová, and c. S. N. Özcan. An asymptotic preserving scheme for kinetic chemotaxis models in two space dimensions. *Kinet. Relat. Models*, 12(1):195–216, 2019.

[6] C. Emako, L. N. De Almeida, and N. Vauchelet. Existence and diffusive limit of a two-species kinetic model of chemotaxis. *Kinetic & Related Models*, 8(2), 2015.

[7] C. Emako, C. Gayrard, A. Buguin, L. Neves de Almeida, and N. Vauchelet. Traveling pulses for a two-species chemotaxis model. *PLoS Computational Biology*, 12(4):e1004843, 2016.

[8] F. Filbet and S. Jin. A class of asymptotic-preserving schemes for kinetic equations and related problems with stiff sources. *Journal of Computational Physics*, 229(20):7625–7648, 2010.

[9] J. Franca-Koh, Y. Kamimura, and P. Devreotes. Navigating signaling networks: chemotaxis in dictyostelium discoideum. *Current Opinion in Genetics & Development*, 16(4):333–338, 2006.

[10] J. Franca-Koh, S. S. Willard, and P. N. Devreotes. G-protein signaling in chemotaxis. In *Handbook of Cell Signaling*, pages 1705–1712. Elsevier, 2010.

[11] S. Jin. Efficient asymptotic-preserving (ap) schemes for some multiscale kinetic equations. *SIAM Journal on Scientific Computing*, 21(2):441–454, 1999.

[12] E. F. Keller and L. A. Segel. Model for chemotaxis. *Journal of Theoretical Biology*, 30(2):225–234, 1971.

[13] M. Lemou and L. Mieussens. A new asymptotic preserving scheme based on micro-macro formulation for linear kinetic equations in the diffusion limit. *SIAM Journal on Scientific Computing*, 31(1):334–368, 2008.

[14] J.-G. Liu, L. Wang, and Z. Zhou. Positivity-preserving and asymptotic preserving method for 2D Keller-Segal equations. *Math. Comp.*, 87(311):1165–1189, 2018.

[15] H. G. Othmer, S. R. Dunbar, and W. Alt. Models of dispersal in biological systems. *Journal of Mathematical Biology*, 26(3):263–298, 1988.

[16] P. Rickert, O. D. Weiner, F. Wang, H. R. Bourne, and G. Servant. Leukocytes navigate by compass: roles of pi3kγ and its lipid products. *Trends in cell biology*, 10(11):466–473, 2000.

[17] J. Saragosti, V. Calvez, N. Bournaveas, A. Buguin, P. Silberzan, and B. Perthame. Mathematical description of bacterial traveling pulses. *PLoS Computational Biology*, 6(8):e1000890, 2010.

[18] S. S. Willard and P. N. Devreotes. Signaling pathways mediating chemotaxis in the social amoeba, dictyostelium discoideum. *European Journal of Cell Biology*, 85(9-10):897–904, 2006.