Xinshi Wang 661975305 HW01

Problem 1

- 1. $\{x \text{ is even } \land y = x + 1 \}$
- 2. $\{1 \le x \le 3\}$
- 3. $\{x > 0 \land y > 0\}$
- 4. $\{x \text{ is divisible by } 50\}$
- 5. None
- 6. $abs(result * result x) \le 0.000001$

Problem 2

- 1. Valid;
- 2. Valid;
- 3. Invalid; we know from code that i=i+1 and j=j-1. Adding those up gives us i+j=i+j. From the precondition we know $i+j\neq 0$. Thus to make it correct, the post condition should also be $i+j\neq 0$.
- 4. Invalid; the else case is $x \leq y$ instead of x < y. Thus, the postcondition should be $\{(m = x \land x > y) \lor (m = y \land x \leq y)\}$

Problem 3

- 1. A code E: possibly invalid. There's no correlation between A and E.
- 2. C code D: possibly invalid. C is a weaker precondition compared to F. We want a stronger one.

Problem 4

1.
$$\{x > 0\}$$

 $x=10;$
 $\{x = 10\}$
 $y=20-x;$
 $\{x = 10 \land y = 20 - x\} \implies \{y = 10 \land x = 10\};$
 $z=y+4;$
 $\{z = y + 4 \land y = 10\} \implies \{y = 10 \land x = 10 \land z = 14\};$
 $y=0;$
 $\{y = 0\} \implies \{y = 0 \land x = 10 \land z = 14\};$

2.
$$\{|x| > 11\}$$
 $x = -x;$
 $\{|x| > 11 \land x = -x\} \implies \{x > 11 \lor x < -11\}$
 $x = x * x;$
 $\{x = x * x \land x > 11 \lor x < -11\} \implies \{x > 121\};$
 $x = x + 1;$
 $\{x = x + 1 \land x > 121\} \implies \{x > 122\};$
3. $\{|x| < 5\}$
if $(x > 0)$ $\{(x > 0) \in \{x > 0 \} \implies \{0 < x < 5\}\}$
 $y = x + 2;$
 $\{0 < x < 5 \land y = x + 2\} \implies \{2 < y < 7\}$
 $\{x > 0 < x < 5 \land y = x + 2\} \implies \{2 < y < 7\}$
 $\{x > 0 < x < 5 \land x < 0\} \implies \{-5 < x < 0\}$
 $\{x > 0 < x < 5 \land x < 0\} \implies \{-6 < y < -1\}$
 $\{x > 0 < x < 0 \} \implies \{-6 < y < -1\}$
 $\{x > 0 < x < 0 \} \implies \{-6 < y < -1\}$
 $\{x > 0 < x < 0 \} \implies \{-6 < y < -1\}$

Problem 5

}

```
1. \{wp(x=-5, y>-2x) \implies (y>10)\}
               x = -5:
                   \{wp(z=2*x+y,z>0) \implies (2x+y>0)=(y>-2x)\}
               z = 2 * x + y;
                   \{z > 0\}:
   2. \{wp(code, x > 0 \lor x \le 0) = \{x > 1 \lor x < -3\}\}\
               if (x > 0){
                   \{wp(x = x + 6, x > 7) = (x + 6 > 7) = (x > 1)\}\
                x = x + 6;
               } else {
                   \{wp(x=4-x,x>7)=(4-x>7)=(x<-3)\}
                   \{x > 7\}
   3. \{wp(\text{if else statements and codes}, x > 0) = (x > 3) \lor (x > -4)\}
               if (x > 4){
                   \{wp(x=x-3, x>0) = (x-3>0) = (x>3)\}\
                x = x - 3;
                } else {
                       \{wp((x < -4) \text{ do } x = x + 3 \land (x \ge -4) \text{ do } (x = x + 1), x > 0\} = (x < -4 \land x > 0)
(-3) \lor (x \ge -4 \land x > -1) = (x \ge -4)
                        if (x < -4)
                             \{wp(x = x + 3, x > 0) = (x > -3)\}\
                           x = x + 3
                        }else {
                             \{wp(x = x + 1, x \ge -4) = (x + 1 > 0) = (x > -1)\}\
                           x = x + 1
                   }
                   {x > 0}
   4. \{wp(x = y + 2, x > 2y - 1) = (y + 2 > 2y - 1) = (y < 3)\}
              x = y + 2;
                   \{wp(z = x + 1, z > 2y) = (x + 1 > 2y) = (x > 2y - 1)\}
              z = x + 1;
                   \{z > 2y\};
```

5.
$$\{wp(x \ge 0 \text{ do } z = x \lor x < 0 \text{ do } z = x + 1, z \ne 0) = (x \ge 0 \land x \ne 0) \lor (x < 0 \land x \ne -1) = -1 < x < 0\}$$

if $(x \ge 0)$

$$\{wp(z = x, z \ne 0) = (x \ne 0)\}$$

$$z = x;$$
else
$$\{wp(z = x + 1, z \ne 0) = (x + 1 \ne 0) = (x \ne -1)\}$$

$$z = x + 1;$$

$$\{z \ne 0\}$$

Problem 6

1.
$$\{x_{pre} < y_{pre}\}$$

$$\{wp(w = y, w = y_{pre} = x_{post} \land y_{post} = x_{pre}) = w = y_{pre} = x_{post} = y_{post} = x_{pre}\}$$

$$w = y;$$

$$\{wp(z = x + y - w, w = y_{pre} = x_{post} \land y_{post} = x_{pre}) = (z = w - w + y_{post}) = (z = x_{pre})\}$$

$$z = x + y - w;$$

$$\{wp(x = w, x_{post} = y_{pre} \land y_{post} = x_{pre}) = (w = y_{pre} = x_{post}) \land y_{post} = x_{pre}\}$$

$$x = w;$$

$$\{x_{post} = y_{pre} \land y_{post} = x_{pre}\}$$

In Sufficient, we need to make sure $x_{pre} = y_{pre}$. The given condition is weaker than $x_{pre} = y_{pre}$.

$$\{(x = y) \lor (x \neq y \land y > 0)\}$$

$$\{wp((x_1 == y) \text{ do } x_2 = 0 \lor (x_1! = y) \text{ do } x_2 = x_1 * y, x_2 \le y) = (x_1 = y \land y \ge 0) \lor (x_1 \neq y \land x_1 \le 1)\}$$

$$\text{if } (x == y);$$

$$\{wp(x_2 = 0, x_2 \le y) = (y \ge 0)\}$$

$$x = 0;$$

$$\text{else}$$

$$\{wp(x_2 = x_1 * y, x_2 \le y) = (x_1 * y \le y) = (x_1 \le 1)\}$$

$$x = x * y;$$

$$\{x \le y\}$$

Insufficient. The given condition is weaker. We also need to make sure $\{x \leq 1\}$ and $y \geq 0$.