Math 54 Lec7

```
Recap  \begin{array}{c} \text{Transpose} \\ \text{properties} \\ \text{Proof of } (AB)^T = B^TA^T \\ \text{Identity matrix} \\ \text{Key property} \\ \text{Inverse (For square matrix)} \\ \text{Left inverse of matrix equals to Right inverse of matrix} \\ \text{Inverse is unique} \\ \text{Terminology} \\ \text{Properties} \end{array}
```

Math 54 Lec7

Recap

```
Linear transformations T:\mathbb{R}^n	o\mathbb{R}^m\ (\mathrm{matrix}\ A), S:\mathbb{R}^p	o\mathbb{R}^n\ (\mathrm{matrix}\ B)
```

Then $T\circ S$ has matrix $A\cdot B$

Transpose

Very important counterpart of any matrix A, denoted by A^T

This switches the rows and columns of A (i.e. i-th row of A becomes i-th column of A^T). If A is $m \times n$, then A^T is $n \times m$.

properties

- $\bullet \ (A+B)^T = A^T + B^T$
- $\bullet \ (AB)^T = B^T A^T$

Proof of
$$(AB)^T = B^T A^T$$

Consider the ij-th entry in $(AB)^T$. That is the ji-th entry in AB which is the j-th row of A times the i-th column in B.

Now consider the ij-entry in B^TA^T . That is the i-th row in $B^T(i$ -th column in B) times j-th column in $A^T(j$ -th row in A).

Identity matrix

For $n \times n$ matrix, I_n is the **identity matrix**, which has 1s for all diagonal entries. 0 for all non-diagonal entries.

I for short if context is clear.

Key property

For any $n \times n$ matrix A, we have $A \cdot I_n = I_n \cdot A = A$.

Inverse (For square matrix)

For an n imes n matrix A, the **"inverse"** of A(if it exists) is denoted A^{-1} has the property that

$$A \cdot A^{-1} = A^{-1} \cdot A = I_n$$

Left inverse of matrix equals to Right inverse of matrix

Say we find B for which $BA = I_n$. Is $AB = I_n$ guaranteed?

Without assuming that say there is also some C where $AC=I_n$

Then
$$BAC=BI_n\Rightarrow I_nC=B\Rightarrow C=B$$

Inverse is unique

Say we have $I_n = AB = AC$

Terminology

A square matrix is **invertible/nonsingular** if it has an inverse, otherwise **singular**.

$$A = egin{bmatrix} a & b \ c & d \end{bmatrix}$$
 , then the **determinant** of A is $ad-bc$. And we have

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Naturally requires ad-bc
eq 0

If
$$ad-bc=0
ightarrowrac{a}{b}=rac{c}{d}$$

Properties

- $(A^{-1})^{-1} = A$
- $(AB)^{-1} = B^{-1}A^{-1}$