

Lesson 7: Solutions of the Schrödinger equation (II)


Clara E. Alonso Alonso

February 13, 2012

Potential step	2
.....	3
.....	4
.....	5
.....	6
.....	7
.....	8
.....	9
.....	10
.....	11
.....	12
.....	13
.....	14
.....	15
.....	16
.....	17
.....	18
.....	19
.....	20
.....	21
.....	22
.....	23
.....	24
.....	25
Potential barrier: barrier penetration	26
.....	27
.....	28
.....	29
.....	30
.....	31
.....	32
.....	33
.....	34
.....	35
.....	36
.....	37
.....	38

.....	39
.....	40
.....	41
WKB (Wentzel, Kramers and Brillouin) Approximation	42
.....	43
.....	44
.....	45
.....	46
.....	47
Applications: α decay and nuclear fusion	48
.....	49
.....	50
.....	51
.....	52
.....	53
.....	54
.....	55
Bibliography	56
.....	57

- (a) $E < V_0$



All states are unbound

Classically: an ∞ force acts at $x = 0$ during an infinitesimal time, causing Δp finite

$$F = - \frac{dV(x)}{dx}$$

3 / 57

Quantum study

- we study particles falling into the step from the left
- we divide the x-axis into two regions I and II

$$\circ \text{ (I)} \quad - \frac{\hbar^2}{2m} \phi_I''(x) = E \phi_I(x)$$

$$x \leq 0 \quad ; \quad k_1 = \frac{\sqrt{2mE}}{\hbar}$$

$$\phi_I(x) = A e^{ik_1 x} + B e^{-ik_1 x}$$

4 / 57

• ...

◦ (II) $-\frac{\hbar^2}{2m}\phi_{II}''(x) + V_0\phi_{II}(x) = E\phi_{II}(x)$

$$\phi_{II}''(x) + \bar{k}_2^2 \phi_{II}(x) = 0 \quad \bar{k}_2 = \frac{\sqrt{2m(E-V_0)}}{\hbar}$$

$$\bar{k}_2 = ik_2 = i\frac{\sqrt{2m(V_0-E)}}{\hbar} \quad ; \quad k_2 \text{ real}$$

$$\phi_{II}(x) = C e^{k_2 x} + D e^{-k_2 x}$$

5 / 57

We will determine the relation between A , B , C and D requiring the continuity of the wave function and its derivative. The wave function is finite $\forall x$

$$\begin{aligned} \phi_I(0) &= \phi_{II}(0) \quad ; \quad \phi_I'(0) = \phi_{II}'(0) \rightarrow \\ A &= \frac{D}{2} \left[1 + i\frac{k_2}{k_1} \right] \quad ; \quad B = \frac{D}{2} \left[1 - i\frac{k_2}{k_1} \right] \end{aligned}$$

$C = 0$ since $e^{k_2 x} \rightarrow \infty$ when $x \rightarrow \infty$

$$\phi(x) = \begin{cases} \frac{D}{2} \left[1 + i\frac{k_2}{k_1} \right] e^{ik_1 x} + \frac{D}{2} \left[1 - i\frac{k_2}{k_1} \right] e^{-ik_1 x} & x \leq 0 \\ D e^{-k_2 x} & x \geq 0 \end{cases}$$

incident wave
reflected wave

transmitted wave

$$\Psi(x, t) = \phi(x) e^{-\frac{iEt}{\hbar}}$$

6 / 57

- We get no conditions limiting the possible values of E (unbound state)
- The wave function is a function of a unique constant D . It can not be determined by normalizing the wave function, since it is not normalizable (unbound state)
- The wave function does not represent a particle but a **beam**. $|D|^2$ can be determined from the incident flux

D can be chosen real

We write $D \left[1 + i \frac{k_2}{k_1} \right] = D' e^{i\delta}$ (1)
 D' and δ real

$$\phi(x) = \begin{cases} D' \cos(k_1 x + \delta) & x \leq 0 \\ D e^{-k_2 x} & x \geq 0 \end{cases}$$

7 / 57

We see that $\phi(x)$ is real if we choose D real. D can be solved from (1) to have the wave function as a function of D' and δ only

Probability current or flux (one dimension)

$$j = \frac{\hbar}{2im} \left(\phi^* \frac{d\phi}{dx} - \frac{d\phi^*}{dx} \phi \right) = \frac{\hbar}{m} \text{Re} \left(\frac{1}{i} \phi^* \frac{d\phi}{dx} \right)$$

If ϕ is real $\rightarrow j = 0$

We can calculate the incident flux (with the incident wave function)

$$j_i = \frac{\hbar k_1 |D|^2}{4m} \left[1 + \left(\frac{k_2}{k_1} \right)^2 \right]$$

(incident particles per unit time)

8 / 57

Reflected flux (calculated from the reflected wave function)

$$j_r = -\frac{\hbar k_1 |D|^2}{4m} \left[1 + \left(\frac{k_2}{k_1} \right)^2 \right] \leq 0$$

$$j = j_i + j_r = 0$$

We define **reflection coefficient**

$$R = -\frac{j_r}{j_i}$$

gives the fraction of the beam reflected in the step

Transmitted flux (calculated with the transmitted wave function)

$$j_t = 0 \quad \text{since} \quad \frac{1}{i} \phi_t^* \frac{d\phi_t}{dx} \quad \text{is purely imaginary}$$

9 / 57

We define **transmission coefficient**

$$T = \frac{j_t}{j_i}$$

For $E < V_0$ $R = 1$; $T = 0$ (as classically)

but quantically the probability of finding the particle with $x \geq 0$ is non-zero (\neq classical)

$$|\phi_{II}(x)|^2 = |D|^2 e^{-2k_2 x} \quad ; \quad x \geq 0$$

exponential decay, the probability drops by a factor e at a distance $\frac{1}{2k_2} = \frac{\hbar}{2\sqrt{2m(V_0-E)}}$

- Larger $V_0 - E$ implies less penetration of the wave function in the classically forbidden region

10 / 57

$\frac{1}{2k_2}$ in this distance prob. decays to 37% of the value in $x = 0$

- Macroscopically $\frac{1}{2k_2}$ is very small

Ex.: $m = 1 \text{ kg}$; $E = 1 \text{ J}$; $V_0 = 2 \text{ J} \rightarrow \frac{1}{2k_2} \approx 4 \cdot 10^{-26} \text{ fm}$

- To locate the particle in the classically forbidden region


$$\Delta x < \frac{\hbar}{2\sqrt{2m(V_0 - E)}} \rightarrow \Delta p \approx \frac{\hbar}{2\Delta x} \approx \sqrt{2m(V_0 - E)}$$

$$\rightarrow \Delta E \approx \frac{(\Delta p)^2}{2m} \approx V_0 - E \quad \text{we can not say that}$$

$$V_0 > E \quad (\text{classically forbidden region})$$

11 / 57

- (b) $E > V_0$



- classically:** the particle slows down when going from I to II ($F_x < 0$). There is no reflection
- quantically:** if E is not much larger than V_0 there is probability that the particle is reflected

12 / 57

Ex. photoelectric cell: the e^- at the cathode receives the photon energy and tries to escape. If E is not much larger than V_0 (V_0 potential to which the e^- is subject on the surface of the metal) there is reflection and it does not escape (there is reduction of efficiency in photoelectric cells for light with $\nu \approx \nu_{\text{threshold}}$)

$$\phi_I(x) = A e^{ik_1 x} + B e^{-ik_1 x} ; \quad k_1 = \frac{\sqrt{2mE}}{\hbar} \quad x \leq 0$$

$$\phi_{II}(x) = C e^{ik_2 x} + D e^{-ik_2 x} ; \quad k_2 = \frac{\sqrt{2m(E - V_0)}}{\hbar} \quad x \geq 0$$

$D = 0$ no reflected wave in region II (there are only **reflections** in the **discontinuities** of potential)

$$\phi_I(0) = \phi_{II}(0) \quad ; \quad A + B = C$$

13 / 57

$$\phi'_I(0) = \phi'_{II}(0)$$

$$ik_1 A - ik_1 B = ik_2 C \rightarrow \frac{k_1}{k_2}(A - B) = C$$

$$C = \frac{2k_1}{k_1 + k_2} A$$

$$B = \frac{k_1 - k_2}{k_1 + k_2} A$$

$$\phi(x) = \begin{cases} A \left(\underset{\text{(i)}}{e^{ik_1 x}} + \underset{\text{(r)}}{\frac{k_1 - k_2}{k_1 + k_2} e^{-ik_1 x}} \right) & x \leq 0 \\ A \frac{2k_1}{k_1 + k_2} \underset{\text{(t)}}{e^{ik_2 x}} & x \geq 0 \end{cases}$$

14 / 57

$$\begin{aligned}
j_i &= \frac{\hbar}{m} \operatorname{Re} \left[\frac{1}{i} A^* e^{-ik_1 x} (ik_1) A e^{ik_1 x} \right] \\
&= \frac{\hbar k_1}{m} |A|^2
\end{aligned}$$

$$\begin{aligned}
j_r &= \frac{\hbar}{m} \operatorname{Re} \left[\frac{1}{i} A^* \frac{k_1 - k_2}{k_1 + k_2} e^{ik_1 x} (-ik_1) A \frac{k_1 - k_2}{k_1 + k_2} e^{-ik_1 x} \right] \\
&= \frac{-\hbar k_1}{m} |A|^2 \left(\frac{k_1 - k_2}{k_1 + k_2} \right)^2
\end{aligned}$$

15 / 57

$$R = -\frac{j_r}{j_i} = \left(\frac{k_1 - k_2}{k_1 + k_2} \right)^2 \quad \text{does not change } k_1 \iff k_2$$

$$\begin{aligned}
j_t &= \frac{\hbar}{m} \operatorname{Re} \left[\frac{1}{i} A^* \frac{2k_1}{k_1 + k_2} e^{-ik_2 x} (ik_2) A \frac{2k_1}{k_1 + k_2} e^{ik_2 x} \right] \\
&= \frac{\hbar k_2}{m} |A|^2 \left(\frac{2k_1}{k_1 + k_2} \right)^2
\end{aligned}$$

$$j_i + j_r = j_t \rightarrow R + T = 1$$


$$\begin{aligned}
T &= \frac{j_t}{j_i} = \frac{k_2}{k_1} \left(\frac{2k_1}{k_1 + k_2} \right)^2 \\
&= \frac{4k_1 k_2}{(k_1 + k_2)^2} \quad \text{it does not vary } k_1 \iff k_2
\end{aligned}$$

16 / 57

$$\begin{aligned}
 R + T &= \left(\frac{k_1 - k_2}{k_1 + k_2} \right)^2 + \frac{4k_1 k_2}{(k_1 + k_2)^2} \\
 &= \left(\frac{k_1 + k_2}{k_1 + k_2} \right)^2 = 1
 \end{aligned}$$

17 / 57

- (i) particle coming from the region of negative x




$$|\phi_I(x)|^2 = |A|^2 \left[1 + 2 \frac{k_1 - k_2}{k_1 + k_2} \cos 2k_1 x + \left(\frac{k_1 - k_2}{k_1 + k_2} \right)^2 \right]$$

It takes values between $\frac{4|A|^2 k_1^2}{(k_1 + k_2)^2}$ and $\frac{4|A|^2 k_2^2}{(k_1 + k_2)^2}$
maximum minimum

$$|\phi_I(x)|_{x=0}^2 = \frac{4|A|^2 k_1^2}{(k_1 + k_2)^2} ; |\phi_{II}(x)|_{\forall x}^2 = \frac{4|A|^2 k_1^2}{(k_1 + k_2)^2}$$

18 / 57

- (ii) particle coming from the left




change $k_1 \iff k_2$; in (i)

$$k_1 = \frac{\sqrt{2mE}}{\hbar} ; k_2 = \frac{\sqrt{2m(E-V_0)}}{\hbar} ; k_1 > k_2 \text{ c}$$

$$|\phi_I(0)|^2 = |\phi_{II}(0)|^2 = \frac{4|A|^2 k_2^2}{(k_1 + k_2)^2} \text{ minima in region I}$$

19 / 57


- if the particle enters from the right in (i), the solution is the same as in (ii) changing $x \iff -x$
- if the particle enters from the right in (ii), the solution is the same as in (i) changing $x \iff -x$
- (1)



we have solved it in (b) $k_1 = \frac{\sqrt{2mE}}{\hbar} ; k_2 = \frac{\sqrt{2m(E-V_0)}}{\hbar}$

20 / 57

- (2)




same as (1), changing $x \iff -x$

R and T as in (1)

21 / 57

- (3)




same as (1), changing $k_1 \iff k_2$

R and T do not vary when changing $k_1 \iff k_2$

22 / 57

- (4)



W.F. as in (3) with change $x \iff -x$

R and T as in (1), (2) and (3)

Once set m , E and $V_0 \rightarrow$ R and T are independent on whether the step goes up or down and where the beam comes from

23 / 57


As a function of E and V_0

$$R = \left(\frac{1 - \sqrt{1 - \frac{V_0}{E}}}{1 + \sqrt{1 - \frac{V_0}{E}}} \right)^2$$

where $\frac{E}{V_0} > 1$

The m of the particles of the beam does not appear, R does not depend on it

24 / 57




case (a) $\rightarrow \frac{E}{V_0} < 1$; case (b) $\rightarrow \frac{E}{V_0} > 1$

$$\frac{E}{V_0} \gg 1 \rightarrow R \rightarrow 0$$

25 / 57

Potential barrier: barrier penetration

26 / 57



$$V(x) = \begin{cases} 0; & x < 0 \text{ and } x > a \\ V_0; & 0 < x < a \end{cases}$$

- (i) $E > V_0$

$$\phi_I = Ae^{ik_1x} + Be^{-ik_1x} ; k_1 = \frac{\sqrt{2mE}}{\hbar}$$

$$\phi_{II} = Pe^{ik_2x} + Qe^{-ik_2x} ; k_2 = \frac{\sqrt{2m(E-V_0)}}{\hbar}$$

$$\phi_{III} = Ce^{ik_1x} + De^{-ik_1x}$$

27 / 57

If we assume that the beam comes from the left $\rightarrow D = 0$

$$\phi_I(0) = \phi_{II}(0) ; \phi'_I(0) = \phi'_{II}(0)$$

$$A + B = P + Q ; k_2(P - Q) = k_1(A - B)$$

$$2A = P(1 + \frac{k_2}{k_1}) + Q(1 - \frac{k_2}{k_1}) ; 2B = P(1 - \frac{k_2}{k_1}) + Q(1 + \frac{k_2}{k_1})$$

$$\phi_{II}(a) = \phi_{III}(a) ; \phi'_{II}(a) = \phi'_{III}(a)$$

$$Pe^{ik_2a} + Qe^{-ik_2a} = Ce^{ik_1a} k_2(Pe^{ik_2a} - Qe^{-ik_2a}) = k_1 Ce^{ik_1a}$$

$$P = \frac{1}{2}Ce^{i(k_1-k_2)a}(1 + \frac{k_1}{k_2})$$

$$Q = \frac{1}{2}Ce^{i(k_1+k_2)a}(1 - \frac{k_1}{k_2})$$

28 / 57

$$A = \left[\frac{1}{4}e^{i(k_1-k_2)a} \frac{(k_1+k_2)^2}{k_1k_2} - \frac{1}{4}e^{i(k_1+k_2)a} \frac{(k_1-k_2)^2}{k_1k_2} \right] C$$

$$B = \left[\frac{1}{4}e^{i(k_1-k_2)a} \frac{k_1^2 - k_2^2}{k_1k_2} - \frac{1}{4}e^{i(k_1+k_2)a} \frac{k_1^2 - k_2^2}{k_1k_2} \right] C$$

$$A = \frac{e^{ik_1a}}{4k_1k_2} \left[(k_1+k_2)^2 e^{-ik_2a} - (k_1-k_2)^2 e^{ik_2a} \right] C$$

$$B = \frac{e^{ik_1a}}{4k_1k_2} (k_1^2 - k_2^2) (e^{-ik_2a} - e^{ik_2a}) C$$

29 / 57

$$j = \frac{\hbar}{m} \text{Re} \left[\frac{1}{i} \phi^* \frac{d\phi}{dx} \right]$$

$$\begin{aligned} \frac{C}{A} &= \frac{4k_1 k_2 e^{-ik_1 a}}{(k_1 + k_2)^2 e^{-ik_2 a} - (k_1 - k_2)^2 e^{ik_2 a}} \\ &= \frac{4k_1 k_2 e^{-i(k_1 - k_2)a}}{(k_1 + k_2)^2 - (k_1 - k_2)^2 e^{i2k_2 a}} \end{aligned}$$

$$\begin{aligned} \frac{B}{A} &= \frac{(k_1^2 - k_2^2)(e^{-ik_2 a} - e^{ik_2 a})}{(k_1 + k_2)^2 e^{-ik_2 a} - (k_1 - k_2)^2 e^{ik_2 a}} \\ &= \frac{(k_1^2 - k_2^2)(1 - e^{i2k_2 a})}{(k_1 + k_2)^2 - (k_1 - k_2)^2 e^{i2k_2 a}} \end{aligned}$$

30 / 57

$$\begin{aligned} R &= \left| \frac{B}{A} \right|^2 \\ &= \frac{(k_1^2 - k_2^2)^2 (1 - e^{-i2k_2 a} - e^{i2k_2 a} + 1)}{[(k_1 + k_2)^2 - (k_1 - k_2)^2 e^{i2k_2 a}][(k_1 + k_2)^2 - (k_1 - k_2)^2 e^{-i2k_2 a}]} \\ &= \frac{(k_1^2 - k_2^2)^2 2(1 - \cos 2k_2 a)}{(k_1 + k_2)^4 - (k_1 + k_2)^2 (k_1 - k_2)^2 (e^{i2k_2 a} + e^{-i2k_2 a}) + (k_1 - k_2)^4} \\ &= \frac{4(k_1^2 - k_2^2)^2 \sin^2 k_2 a}{2k_1^4 + 2k_2^4 + 12k_1^2 k_2^2 - (k_1^2 - k_2^2)^2 2 \cos 2k_2 a} \\ &= \frac{4(k_1^2 - k_2^2)^2 \sin^2 k_2 a}{2(k_1^2 - k_2^2)^2 (1 - 2 \cos 2k_2 a) + 16k_1^2 k_2^2} \\ &= \frac{1}{1 + \frac{4k_1^2 k_2^2}{(k_1^2 - k_2^2)^2 \sin^2 k_2 a}} \\ &= \left[1 + \frac{4k_1^2 k_2^2}{(k_1^2 - k_2^2)^2 \sin^2 k_2 a} \right]^{-1} \end{aligned}$$

31 / 57

$$\begin{aligned}
T &= \left| \frac{C}{A} \right|^2 \\
&= \frac{16k_1^2 k_2^2}{(k_1 + k_2)^4 - (k_1 + k_2)^2 (k_1 - k_2)^2 (e^{i2k_2 a} + e^{-i2k_2 a}) + (k_1 - k_2)^4} \\
&= \frac{16k_1^2 k_2^2}{2(k_1^2 - k_2^2)^2 \sin^2 k_2 a + 16k_1^2 k_2^2} \\
&= \left[1 + \frac{(k_1^2 - k_2^2)^2 \sin^2 k_2 a}{4k_1^2 k_2^2} \right]^{-1}
\end{aligned}$$

as a function of E and V_0 : $k_1^2 = \frac{2mE}{\hbar^2}$; $k_2^2 = \frac{2m(E-V_0)}{\hbar^2}$

32 / 57

$$\begin{aligned}
R &= \left[1 + \frac{4E(E - V_0)}{V_0^2 \sin^2 k_2 a} \right]^{-1} \\
T &= \left[1 + \frac{V_0^2 \sin^2 k_2 a}{4E(E - V_0)} \right]^{-1}
\end{aligned}$$

- $k_2 a = n\pi$; $n = 1, 2, \dots \rightarrow \lambda_2 = \frac{2a}{n}$

$$a = n \frac{\lambda_2}{2}$$

If the width of the barrier is an integral number of half-wavelengths \rightarrow **perfect transmission** (the barrier is completely transparent to the incident particles, resonance condition)

Observed in the scattering of electrons by noble-gas atoms, “**Ramsauer effect**”

33 / 57

- (ii) $E < V_0$

$$\phi_{II} = Pe^{-k'_2 x} + Qe^{k'_2 x} \quad ; \quad k'_2 = \frac{\sqrt{2m(V_0 - E)}}{\hbar}$$

Change the previous k_2 by ik'_2

$$R = \left[1 + \frac{4k_1^2(-k'_2{}^2)}{(k_1^2 + k'_2{}^2)^2 \sin^2(ik'_2 a)} \right]^{-1}$$

$$\sin \alpha = -i \sinh(i\alpha)$$

$$\sin^2 \alpha = -\sinh^2(i\alpha)$$

$$\sin^2(i\alpha) = -\sinh^2(\alpha)$$

$$R = \left[1 + \frac{4k_1^2(k'_2{}^2)}{(k_1^2 + k'_2{}^2)^2 \sinh^2 k'_2 a} \right]^{-1}$$

34 / 57

$$\begin{aligned} T &= \left[1 + \frac{(k_1^2 + k'_2{}^2)^2 \sin^2 ik'_2 a}{4k_1^2(-k'_2{}^2)} \right]^{-1} \\ &= \left[1 + \frac{(k_1^2 + k'_2{}^2)^2 \sinh^2 k'_2 a}{4k_1^2(k'_2{}^2)} \right]^{-1} \\ &= \left[1 + \frac{V_0^2 \sinh^2 k'_2 a}{4E(V_0 - E)} \right]^{-1} \neq 0 \end{aligned}$$

⇓

TUNNELING

35 / 57

Some interesting limits

- 1) $E > V_0$

If $E \rightarrow V_0 \Rightarrow k_2 \rightarrow 0$; $\sin^2(k_2 a) \approx (k_2 a)^2$

$$T_{E \rightarrow V_0}^{E > V_0} \rightarrow \left(1 + \frac{m V_0 a^2}{2 \hbar^2}\right)^{-1}$$

- 2) $E < V_0$

If $E \ll V_0$ and a is large / $k'_2 a \gg 1 \Rightarrow \sinh x \approx \cosh x = \frac{e^x}{2}$ if $x \gg 1$

36 / 57

$$\begin{aligned} T_{k'_2 a \gg 1}^{E < V_0} &\rightarrow \left(1 + \frac{V_0^2 e^{2k'_2 a}}{16E(V_0 - E)}\right)^{-1} \\ &= \frac{1}{\frac{V_0^2 e^{2k'_2 a}}{16E(V_0 - E)} \left(1 + \frac{16E(V_0 - E)}{V_0^2 e^{2k'_2 a}}\right)} = \frac{x'}{1 + x'} \end{aligned}$$


with $x' = \frac{16E(V_0 - E)}{V_0^2 e^{2k'_2 a}} \ll 1$

Expanding around $x' = 0$

$$T_{k'_2 a \gg 1}^{E < V_0} \approx x' = \frac{16E(V_0 - E)e^{-2k'_2 a}}{V_0^2}$$

37 / 57


For $\frac{2mV_0a^2}{\hbar^2} = \frac{47}{3}$



$T = 1$ implies perfect transmission

38 / 57


Ex. Barrier with $a = 4$ arbitrary units (a.u.) and $V_0 = 3$ a.u.



39 / 57

$$E/V_0 = 4/3 \quad ; \quad a = 4$$


$$|\Psi|^2 \text{ (a.u.)}$$



40 / 57


$$E/V_0 = 2.7/3 \quad ; \quad a = 4$$

$$|\Psi|^2 \text{ (a.u.)}$$



41 / 57

- For barriers of arbitrary shape
- We will calculate the **penetrability** \mathcal{T}



- For rectangular barriers we obtained $T_{k'_2 a} \gg 1$ ($E < V_0$)


\mathcal{T} is the dominant factor $e^{-2k'_2 a}$

$$T_{k'_2 a} \gg 1 \sim \frac{16E(V_0 - E)}{V_0^2 e^{2k'_2 a}} ; \quad E < V_0$$

$$\frac{16E(V_0 - E)}{V_0^2} \begin{cases} \text{maximum 4} & E < V_0 \\ \sim 1 & \text{unless } E \sim V_0 \text{ then } k'_2 a \text{ is not } \gg 1 \end{cases}$$

The dominant factor is the exponential

We approximate $T_{k'_2 a} \gg 1 \approx e^{-2k'_2 a}$



Approx.: succession of thin rectangular barriers

$$V(x_1) = V(x_2) = E$$

In x_1 and x_2 , $k_2' a$ is not $\gg 1$

45 / 57

$\mathcal{P}(x)$ probability of finding the particle at x ($x_1 \leq x \leq x_2$)

$\mathcal{P}(x + \Delta x)$ (probability of reaching x) \times (probability of crossing the barrier of width Δx)

$$\mathcal{P}(x + \Delta x) = \mathcal{P}(x) e^{-2k(x)|\Delta x|} ; \quad k(x) = \frac{\sqrt{2m(V(x) - E)}}{\hbar}$$

If $\Delta x \rightarrow 0$

$$\mathcal{P}(x + dx) = \mathcal{P}(x) (1 - 2k(x) |dx|)$$

$$\frac{\mathcal{P}(x + dx) - \mathcal{P}(x)}{|dx|} = -2k(x)\mathcal{P}(x)$$

46 / 57

$$\int_{x_1}^x \frac{d\mathcal{P}(x)}{\mathcal{P}(x)} = - \int_{x_1}^x 2k(x)|dx| \Rightarrow \ln \frac{\mathcal{P}(x)}{\mathcal{P}(x_1)} = - \left| \int_{x_1}^x 2k(x)dx \right|$$

$$\mathcal{P}(x) = \mathcal{P}(x_1)e^{-2\left|\int_{x_1}^x k(x)dx\right|}$$

$$\mathcal{T} = \frac{\mathcal{P}(x_2)}{\mathcal{P}(x_1)} = e^{-2\left|\int_{x_1}^{x_2} k(x)dx\right|}$$


The value of \mathcal{T} does not depend on the way the barrier is crossed

47 / 57

Applications: α decay and nuclear fusion

48 / 57

- we assume that the α particle had a continuing existence as distinct object inside the nucleus (${}^4_2\text{He}_2$)
- it is subject to nuclear attractive potential plus Coulomb repulsion
- to exit the nucleus it has to cross the Coulomb barrier



49 / 57

- ${}_{92}^{238}\text{U} \rightarrow$ there is a barrier of ~ 37 MeV at $R \sim 7$ fm from the center of the nucleus
- α of $E_c = 4.3$ MeV are emitted

$$E^{coul} \simeq \frac{ZZ'e^2}{R} \simeq 37 \text{ MeV}$$


$$R \simeq 1.13A^{1/3} \text{ fm} \simeq 7 \text{ fm}$$

Fine structure constant: $\alpha = \frac{e^2}{4\pi\epsilon_0 \hbar c} = \frac{1}{137}$ (Gaussian units $4\pi\epsilon_0 = 1$)

- in the nucleus $E_c(\alpha) \simeq 10 \text{ MeV} \rightarrow v_d \simeq 10^7 \text{ m/s}$
- classical image $\rightarrow \alpha$ particle strikes the nucleus wall every

$$\Delta t = \frac{2R}{v_d} \simeq 10^{-22} \text{ s/collision}$$

50 / 57



51 / 57

- \mathcal{T} chance, for each impact, that it will succeed in tunneling through the barrier to the outside
- probability of emission per unit time $\rightarrow \frac{\mathcal{T}}{\Delta t}$
- average time to cross the barrier $\tau \sim \frac{\Delta t}{\mathcal{T}}$

$$\mathcal{T} = e^{-2 \int_{r_1}^{r_2} k(r) dr}$$

$$k(r) = \frac{\sqrt{2\mu(V^{coul}(r) - E_c)}}{\hbar} ; \quad V^{coul}(r) = \frac{ZZe^2}{r}$$

$$Z' = 2 ; \quad Z = Z_{\text{nucleus}} - 2 \simeq Z_{\text{nucleus}}$$

$$r_1 \simeq R ; \quad r_2 = \frac{ZZe^2}{E_c}$$

$$E_c = \frac{1}{2}m_\alpha v_f^2 ; \quad v_f \rightarrow \text{speed of emitted } \alpha$$

52 / 57

$$\mu \simeq m_\alpha$$

$$\begin{aligned} \int_{r_1}^{r_2} k(r) dr &= \frac{\sqrt{2\mu}}{\hbar} \int_R^{\frac{ZZe^2}{E_c}} dr \sqrt{\frac{ZZe^2}{r} - E_c} \\ &= \frac{2ZZe^2}{\hbar v_f} \left[-\frac{\sqrt{\gamma-1}}{\gamma} + \arccos \frac{1}{\sqrt{\gamma}} \right] \end{aligned}$$

$$\text{where } \gamma = \frac{ZZe^2}{E_c R}$$

$$\mathcal{T} = e^{-\frac{4ZZe^2}{\hbar v_f} \left[-\frac{\sqrt{\gamma-1}}{\gamma} + \arccos \frac{1}{\sqrt{\gamma}} \right]}$$

53 / 57

- if $\frac{ZZ'e^2}{R} \gg E_c \rightarrow \gamma \gg 1$

$$V^{coul}(R) \gg E_c \rightarrow \arccos \frac{1}{\sqrt{\gamma}} = \frac{\pi}{2} ; \frac{\sqrt{\gamma-1}}{\gamma} \rightarrow \frac{1}{\sqrt{\gamma}}$$

$$\mathcal{T} \simeq e^{-\left(\pi \frac{ZZ'e^2}{\hbar} \sqrt{\frac{2\mu}{E_c} - \frac{2}{\hbar} \sqrt{2\mu e^2 ZZ'R}}\right)}$$

Applying it to ^{238}U

$$E_c \simeq 4 \text{ MeV} ; \frac{ZZ'e^2}{R} \simeq 37 \text{ MeV} ; \gamma \gg 1 ; Z = 90 ; Z' = 2 ; \mu \simeq 4 \times 931 \frac{\text{MeV}}{c^2} ; R = 7 \text{ fm}$$

$$\mathcal{T} \sim 10^{-39}$$

$$T_{1/2} \sim \frac{\Delta t}{\mathcal{T}} \simeq \frac{10^{-22}}{10^{-39}} \text{ s} \simeq 10^{10} \text{ y}$$

according to $T_{1/2}$ observed, $4.5 \times 10^9 \text{ y}$

54 / 57

- tunneling in nuclear physics in reverse: penetration of charged particles \rightarrow **FUSION**

55 / 57

- [1] A.P.French y E.F. Taylor, "Introducción a la Física Cuántica"
- [2] S. Gasiorowicz, "Quantum Physics", ed. John Wiley, 2003, Apéndice B
- [3] D.J. Griffiths, "Introduction to Quantum Mechanics", ed. Pearson Education Inc., 2005
- [4] D. Park, "Introduction to the Quantum Theory", ed. McGraw-Hill, 1992
- [5] C. Sánchez del Río, Física cuántica, 2003
- [6] R.C. Greenhow, "Introductory Quantum Mechanics", IoP Publishing, 1995