

Lesson 3: The interpretation of matter waves

Clara E. Alonso Alonso

October 19, 2011

Probability density. Born interpretation of the wave function	2
.....	3
.....	4
.....	5
.....	6
Mean or expected values	7
.....	8
.....	9
Operators in quantum mechanics (rudiments)	10
.....	11
.....	12
.....	13
.....	14
.....	15
Probability current	16
.....	17
.....	18
.....	19
.....	20
.....	21
.....	22
.....	23
.....	24
Bibliography	25
.....	26

$$\Psi(\vec{r}, t)$$

- complex
- not measurable quantities

Connection between $\Psi(\vec{r}, t)$ and associated particle behavior



PROBABILITY DENSITY

$$P(\vec{r}, t) = |\Psi(\vec{r}, t)|^2$$

probability per unit volume of finding the particle in the neighborhood of \vec{r} at time t (is real and positive)

in one dimension $P(x, t) = |\Psi(x, t)|^2 \rightarrow$ prob. per unit length x. .. t

3 / 26

PROBABILITY

$$P(\vec{r}, t) d^3\vec{r}$$


probability of finding the particle in a volume $d^3\vec{r}$ around \vec{r} at time t
(in one dimension $P(x, t) dx$ probab.....in length dx x....t)

We demand the normalization of the wave function

$$\int_{all\ space} P(\vec{r}, t) d^3\vec{r} = 1$$

(the probability of finding the particle in all space is 100 %)

4 / 26



5 / 26

In one dimension, the probability of finding the particle between $x = a$ and $x = b$, $P(a \leq x \leq b)$:

$$\int_a^b P(x, t) dx$$

In three dimensions, the probability of finding the particle in a volume V

$$\int_V P(\vec{r}, t) d^3\vec{r}$$

Quantum predictions do not give the position of the particle in a time (classical), they give the probability of being in a volume (or length)

they are statistical predictions

It is Born's interpretation of the wave function

6 / 26

Characteristic position of a particle with wave function $\Psi(\vec{r}, t)$ can be its mean or expected value

$$\langle \vec{r} \rangle = \bar{\vec{r}} = \frac{\int \vec{r} P(\vec{r}, t) d^3\vec{r}}{\int P(\vec{r}, t) d^3\vec{r}} = \frac{\int \Psi^*(\vec{r}, t) \vec{r} \Psi(\vec{r}, t) d^3\vec{r}}{\int \Psi^*(\vec{r}, t) \Psi(\vec{r}, t) d^3\vec{r}}$$

For a problem in one dimension

$$\langle x \rangle = \bar{x} = \frac{\int_{-\infty}^{\infty} \Psi^*(x, t) x \Psi(x, t) dx}{\int_{-\infty}^{\infty} \Psi^*(x, t) \Psi(x, t) dx}$$

Expected value of $f(x)$

$$\langle f(x) \rangle = \frac{\int_{-\infty}^{\infty} \Psi^*(x, t) f(x) \Psi(x, t) dx}{\int_{-\infty}^{\infty} \Psi^*(x, t) \Psi(x, t) dx}$$

analogue in three dimensions

Expected value of $f(x, t)$

$$\langle f(x, t) \rangle = \frac{\int_{-\infty}^{\infty} \Psi^*(x, t) f(x, t) \Psi(x, t) dx}{\int_{-\infty}^{\infty} \Psi^*(x, t) \Psi(x, t) dx}$$

With functions $\neq f(\vec{r}, t)$. How do you calculate mean values?

↓

operators must be associated

- For example $\vec{p} \neq f(\vec{r})$

Let's consider the plane wave $\Psi(\vec{r}, t) = e^{i(\vec{k} \cdot \vec{r} - \omega t)}$

$$\frac{\partial \Psi(\vec{r}, t)}{\partial x_j} = i k_j \Psi(\vec{r}, t) = i \frac{p_j}{\hbar} \Psi(\vec{r}, t); j = 1, 2, 3$$

$$\vec{p} \Psi(\vec{r}, t) = -i\hbar \vec{\nabla} \Psi(\vec{r}, t)$$

we associate $\vec{p} \rightarrow -i\hbar \vec{\nabla}$ (differential operator)

11 / 26

- For the energy

$$\frac{\partial \Psi(\vec{r}, t)}{\partial t} = -i \omega \Psi(\vec{r}, t) = -i \frac{E}{\hbar} \Psi(\vec{r}, t)$$

$$E \Psi(\vec{r}, t) = i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t}$$

we associate $E \rightarrow i\hbar \frac{\partial}{\partial t}$ (differential operator)

- If the particle is not free

$$\frac{p^2}{2m} + V(\vec{r}, t) = E$$

we substitute the associated operators

$$-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{r}, t) = i\hbar \frac{\partial}{\partial t} = H$$

12 / 26

We obtain the operator equation

We call **Hamiltonian** the operator associated with the energy, H

Therefore

$$-\frac{\hbar^2}{2m} \nabla^2 \Psi(\vec{r}, t) + V(\vec{r}, t) \Psi(\vec{r}, t) = i \hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t}$$

SCHRÖDINGER EQUATION

ASSOCIATED OPERATORS ARE COMPATIBLE WITH SCHRÖDINGER EQUATION

13 / 26

If we assume a normalized $\Psi(\vec{r}, t)$ $\int |\Psi(\vec{r}, t)|^2 d^3\vec{r} = 1$

$$\begin{aligned} \langle \vec{p} \rangle &= \int d^3\vec{r} \Psi^*(\vec{r}, t) \left(-i\hbar \vec{\nabla} \right) \Psi(\vec{r}, t) \\ &= -i\hbar \int d^3\vec{r} \Psi^*(\vec{r}, t) \vec{\nabla} \Psi(\vec{r}, t) \end{aligned}$$

Important: you have to respect the order of the definition of mean value (because there appear differential operators)

$$\begin{aligned} \langle H \rangle &= \int d^3\vec{r} \Psi^*(\vec{r}, t) \left(i\hbar \frac{\partial}{\partial t} \right) \Psi(\vec{r}, t) \\ &= i\hbar \int d^3\vec{r} \Psi^*(\vec{r}, t) \frac{\partial \Psi(\vec{r}, t)}{\partial t} \\ \langle H \rangle &= \int d^3\vec{r} \Psi^*(\vec{r}, t) \left(\frac{-\hbar^2}{2m} \nabla^2 + V(\vec{r}, t) \right) \Psi(\vec{r}, t) \end{aligned}$$

14 / 26

The wave function has information of expected values of any dynamic quantity

$$\langle f(\vec{r}, \vec{p}, t) \rangle = \int d^3\vec{r} \Psi^*(\vec{r}, t) f(\vec{r}, -i\hbar\vec{\nabla}, t) \Psi(\vec{r}, t)$$

15 / 26

Probability current

16 / 26

We study the propagation of the probability density over time for a particle moving under the influence of a potential V . We assume that $\Psi(\vec{r}, t)$ is normalized for all t . We have

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{r}, t) \right) \Psi(\vec{r}, t) = i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} \quad (1)$$

(Schrödinger eq.)

We take complex conjugates in (1). We assume

$$\begin{aligned} V &= V^* \text{ (real)} \\ V &= V(\vec{r}, t) \end{aligned}$$

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{r}, t) \right) \Psi^*(\vec{r}, t) = -i\hbar \frac{\partial \Psi^*(\vec{r}, t)}{\partial t}$$

17 / 26

The time variation of $|\Psi(\vec{r}, t)|^2$

$$\begin{aligned}\frac{\partial |\Psi|^2}{\partial t} &= \frac{\partial (\Psi^* \Psi)}{\partial t} = \frac{\partial \Psi^*}{\partial t} \Psi + \Psi^* \frac{\partial \Psi}{\partial t} \\ &= \frac{i\hbar}{2m} (\Psi^* \Delta \Psi - \Psi \Delta \Psi^*) \\ &\quad - \frac{1}{i\hbar} (V \Psi^*) \Psi + \Psi^* \frac{1}{i\hbar} V \Psi \\ \Delta &= \nabla^2\end{aligned}$$

As $V = V(\vec{r}, t) \rightarrow (V \Psi^*) \Psi = \Psi^* V \Psi$ (it is a product, V does not involve derivatives)

$$\frac{\partial |\Psi|^2}{\partial t} = \frac{i\hbar}{2m} (\Psi^* \Delta \Psi - \Psi \Delta \Psi^*)$$

18 / 26

We define **probability current (vector)**

$$\begin{aligned}\vec{J}(\vec{r}, t) &= \frac{i\hbar}{2m} (\Psi(\nabla \Psi^*) - \Psi^*(\nabla \Psi)) \\ &= \frac{\hbar}{m} \text{Re} \left(\frac{1}{i} \Psi^* (\nabla \Psi) \right) \\ \vec{\nabla} \cdot \vec{J} &= \frac{i\hbar}{2m} (\Psi \Delta \Psi^* - \Psi^* \Delta \Psi)\end{aligned}$$

a We used

$$\begin{aligned}\vec{\nabla} \cdot (\phi \vec{A}) &= \vec{\nabla} \phi \cdot \vec{A} + \phi \vec{\nabla} \cdot \vec{A} \\ \text{and} \\ a - a^* &= \frac{2}{i} \text{Re} \left(\frac{a^*}{i} \right)\end{aligned}$$

19 / 26

Therefore (**continuity equation**)

$$\frac{\partial |\Psi|^2}{\partial t} + \vec{\nabla} \cdot \vec{J} = 0$$

Analogy with fluids $\frac{\partial \rho}{\partial t} = - \vec{\nabla} \cdot \vec{j}$

$$|\Psi|^2 \rightarrow \rho$$

$$\vec{J} \rightarrow \vec{j}$$

In one dimension $J \rightarrow$ **probability current**

$$J = \frac{\hbar}{2mi} \left[\phi^* \frac{\partial \phi}{\partial x} - \phi \frac{\partial \phi^*}{\partial x} \right]$$

20 / 26

If we integrate over all space the continuity equation

$$\int_{all \ space} \frac{\partial |\Psi|^2}{\partial t} d\tau = - \int_{all \ space} \vec{\nabla} \cdot \vec{J} d\tau$$

By the theorem of Gauss

$$\frac{d}{dt} \int_{all \ space} |\Psi|^2 d\tau = - \oint_S \vec{J} \cdot d\vec{S}$$

But $\oint_S \vec{J} \cdot d\vec{S} = 0$ as the flow of \vec{J} through the surface surrounding the entire space is zero,

$$\frac{d}{dt} \int_{all \ space} |\Psi|^2 d\tau = 0$$

Therefore, the normalization of the wave function is the same at all times

21 / 26

Variation of probability in a volume V (over time) \rightarrow due to flow of \vec{J} through its surface (conservation of probability)

In 1D

$$\frac{d}{dt} \int_a^b dx P(x, t) = - \int_a^b dx \frac{\partial J(x, t)}{\partial x} = J(a, t) - J(b, t)$$

A change of prob. in a region is compensated by a net change of flow in the same region

The fact that $\int |\Psi|^2 d\tau$ extended to the entire space remains constant at all times does not mean that $|\Psi|^2$ should be independent of t at each point

22 / 26

The situation is equivalent to that given in **electromagnetism**. If, in an isolated system, there is charge distributed in the space with volume density $\rho(\vec{r}, t)$, the total charge (the integral of $\rho(\vec{r}, t)$ extended to all space) is conserved in the time. However, in the system, the spatial distribution of this charge can vary, resulting in charge currents.

If the charge contained within a fixed volume V varies over time, closed surface S which surround V must be traversed by a electric current. The variation dQ in time dt of the charge

contained in V is equal to $-I dt$, where I is the intensity of the current through S , ie, the flow of current density vector $J(\vec{r}, t)$ that leaves S

23 / 26

ANALOGY	$ \Psi ^2$	$\rho(\vec{r}, t)$
	probability	charge

1 dim.: $[\phi] \rightarrow L^{-\frac{1}{2}}$ $[J] \rightarrow T^{-1}$

3 dim.: $[\Psi] \rightarrow L^{-\frac{3}{2}}$ $[\vec{J}] \rightarrow L^{-2}T^{-1}$

24 / 26

Bibliography

25 / 26

- [1] D.J. Griffiths, "Introduction to Quantum Mechanics", ed. Pearson Education Inc., 2005
- [2] J.L.Basdevant and J. Dalibard, "Quantum mechanics", ed. Springer, 2002
- [3] A.P.French and E.F. Taylor, Introducción a la Física Cuántica, 1982
- [4] D. Park, "Introduction to the quantum theory", ed. McGraw-Hill, 1992
- [5] C. Sánchez del Río, Física cuántica, 2003

26 / 26