

Tema 9: The harmonic oscillator

Clara E. Alonso Alonso

March 19, 2012

Analytic method	2
.....	3
.....	4
.....	5
.....	6
.....	7
.....	8
.....	9
.....	10
.....	11
.....	12
.....	13
Algebraic solution of the harmonic oscillator. Introduction to the second quantization	14
.....	15
.....	16
.....	17
.....	18
.....	19
.....	20
.....	21
.....	22
.....	23
.....	24
.....	25
Bibliography	26
.....	27



$$V(x) = \frac{1}{2}kx^2 ; k = m\omega^2$$

$$-\frac{\hbar^2}{2m} \frac{d^2\phi(x)}{dx^2} + \frac{1}{2}m\omega^2 x^2 \phi(x) = E\phi(x) \quad \text{T.I.S.E.}$$

■ change of variables

$$y = \sqrt{\frac{m\omega}{\hbar}} x = \alpha x$$

$$\frac{d}{dx} = \alpha \frac{d}{dy} ; \quad \frac{d^2}{dx^2} = \alpha^2 \frac{d^2}{dy^2} ; \quad \alpha = \sqrt{\frac{m\omega}{\hbar}}$$

$$\left[\frac{d^2}{dy^2} + (\beta - y^2) \right] \Phi(y) = 0 \quad (1)$$

$$\beta = \frac{2E}{\hbar\omega} ; \quad \Phi(y) = \phi(x) \quad \text{where } y \text{ and } \beta \text{ are dimensionless}$$

■ asymptotically for $y \rightarrow \infty$

$$\left[\frac{d^2}{dy^2} - y^2 \right] \Phi^{as}(y) = 0 \quad (2)$$

We try $\Phi^{as}(y) = y^n e^{-\frac{y^2}{2}}$; n integer \rightarrow satisfies the differential equation (2)

■ we propose as a general solution $(\forall y)$

$$\Phi(y) = H(y) e^{-\frac{y^2}{2}}$$

so that (1) is fulfilled

$$H''(y) - 2yH'(y) + (\beta - 1)H(y) = 0 \quad (3)$$

5 / 27

we propose

$$H(y) = \sum_{k=0}^{\infty} a_k y^k \quad \text{power series}$$

since $V(x) = V(-x)$ in the sum only appear odd or even k 's

$$H'(y) = \sum_{k=1}^{\infty} k a_k y^{k-1}$$

$$\begin{aligned} H''(y) &= \sum_{k=2}^{\infty} k(k-1) a_k y^{k-2} \\ &= \sum_{p=0}^{\infty} (p+2)(p+1) a_{p+2} y^p \end{aligned}$$

6 / 27

Bring H' and H'' to (3)

$$\sum_{p=0}^{\infty} (p+2)(p+1) a_{p+2} y^p - 2y \left(\sum_{k=0}^{\infty} k a_k y^{k-1} \right) + (\beta - 1) \sum_{k=0}^{\infty} a_k y^k = 0$$

$$\sum_{k=0}^{\infty} [(k+2)(k+1) a_{k+2} - (2k - \beta + 1) a_k] y^k = 0 \quad \forall y$$

$$\frac{a_{k+2}}{a_k} = \frac{2k+1-\beta}{(k+2)(k+1)} \quad \text{recurrence relation}$$

$$\frac{a_{k+2}}{a_k} \rightarrow \frac{2}{k} \quad \text{when } k \rightarrow \infty$$

$$e^{y^2} = \sum_{n=0}^{\infty} \frac{y^{2n}}{n!} = \sum_{k=0(\text{even})}^{\infty} \frac{y^k}{\left(\frac{k}{2}\right)!} = \sum_{k=0(\text{even})}^{\infty} b_k y^k$$

7 / 27

$$\frac{b_{k+2}}{b_k} = \left(\frac{\left[\frac{k+2}{2}\right]!}{\left[\frac{k}{2}\right]!} \right)^{-1} = \left(\frac{k}{2} + 1 \right)^{-1} \rightarrow \frac{2}{k} \quad \text{when } k \rightarrow \infty$$

$$e^{y^2} = \sum_{t=1(\text{odd})}^{\infty} \frac{y^{t-1}}{\left(\frac{t-1}{2}\right)!}$$

$$\rightarrow y e^{y^2} = \sum_{t=1(\text{odd})}^{\infty} \frac{y^t}{\left(\frac{t-1}{2}\right)!} = \sum_{t=1(\text{odd})}^{\infty} c_t y^t$$

$$\frac{c_{t+2}}{c_t} = \left(\frac{\left[\frac{t+1}{2}\right]!}{\left[\frac{t-1}{2}\right]!} \right)^{-1} = \left(\frac{t+1}{2} \right)^{-1} \rightarrow \frac{2}{t} \quad \text{when } t \rightarrow \infty$$

8 / 27

If $H(y)$ is an infinite series, asymptotically

even series $y \rightarrow \infty$; $\Phi(y) = e^{y^2} e^{-\frac{y^2}{2}} = e^{\frac{y^2}{2}} \rightarrow \infty$ when $y \rightarrow \infty$

$$a_n \neq 0 ; a_{n+2} = 0 \rightarrow H(y) \text{ polynomial}$$

$$2n + 1 - \beta = 0 \rightarrow \beta = 2n + 1 \quad (n = 0, 1, 2, \dots)$$

We need

■ a_0 for even eigenfunctions (generates a_2, a_4, \dots)

$$(n = 0, 2, 4, \dots) ; a_1 = 0$$


■ a_1 for odd eigenfunctions (generates a_3, a_5, \dots)

$$(n = 1, 3, 5, \dots) ; a_0 = 0$$

9 / 27

$$\beta = \frac{2E}{\hbar\omega} = 2n + 1 \rightarrow$$

$$E_n = \frac{1}{2}(2n + 1)\hbar\omega = \left(n + \frac{1}{2}\right)\hbar\omega ; n = 0, 1, 2, \dots$$



10 / 27

$$E_0 = \frac{1}{2} \hbar \omega \quad \text{zero point energy}$$

equidistant levels $E_{n+1} - E_n = \hbar \omega$

$H(y)$ are the **Hermite polynomials**

$$H_n(y) = (-1)^n e^{y^2} \frac{d^n e^{-y^2}}{dy^n}$$

We define $\alpha = \sqrt{\frac{m\omega}{\hbar}} \rightarrow y = \alpha x$

$$n = 0 \quad ; \quad \phi_0(x) = \left(\frac{\alpha}{\sqrt{\pi}} \right)^{\frac{1}{2}} e^{-\frac{\alpha^2 x^2}{2}} \quad ; \quad H_0(y) = 1$$

$$n = 1 \quad ; \quad \phi_1(x) = \left(\frac{\alpha}{2\sqrt{\pi}} \right)^{\frac{1}{2}} 2\alpha x e^{-\frac{\alpha^2 x^2}{2}} \quad ; \quad H_1(y) = 2y$$

11 / 27


$$n = 2 \quad ; \quad \phi_2(x) = \left(\frac{\alpha}{8\sqrt{\pi}} \right)^{\frac{1}{2}} (4\alpha^2 x^2 - 2) e^{-\frac{\alpha^2 x^2}{2}} \quad ; \quad H_2 = 4y^2 - 2$$

$$\phi_n(x) = \Phi_n(y) = \left(\frac{\alpha}{2^n n! \sqrt{\pi}} \right)^{\frac{1}{2}} H_n(\alpha x) e^{-\frac{\alpha^2 x^2}{2}} = \mathcal{N}_n H_n(y) e^{-\frac{y^2}{2}} \quad \forall x$$

Normalized wave functions

$$\int_{-\infty}^{\infty} H_n^2(y) e^{-y^2} dy = \sqrt{\pi} 2^n n!$$

12 / 27



13 / 27

Algebraic solution of the harmonic oscillator. Introduction to the second quantization

14 / 27

$$y = \alpha x \quad ; \quad \alpha = \sqrt{\frac{m\omega}{\hbar}} \quad ; \quad \eta = \frac{E}{\hbar\omega} = \frac{\beta}{2}$$

$$\hat{D} = \frac{d}{dy}$$

$$(1) \quad \left[\frac{d^2}{dy^2} + (\beta - y^2) \right] \Phi(y) = 0 \rightarrow$$

$$\frac{1}{2} \left[-\hat{D}^2 + y^2 \right] \Phi(y) = \eta \Phi(y)$$

$$\hat{D}^2 - y^2 = (\hat{D} - y)(\hat{D} + y) - 1$$

$$\hat{D}y \neq y\hat{D} \quad ; \quad [\hat{D}, y] = 1$$

$$\frac{1}{2} \left[1 - (\hat{D} - y)(\hat{D} + y) \right] \Phi = \eta \Phi$$

15 / 27

$$-\frac{1}{\sqrt{2}} \left(\hat{D} - y \right) \frac{1}{\sqrt{2}} \left(\hat{D} + y \right) \Phi = \left(\eta - \frac{1}{2} \right) \Phi \quad (4)$$

$$b^\dagger = \frac{1}{\sqrt{2}} \left(y - \frac{d}{dy} \right) \quad ; \quad b = \frac{1}{\sqrt{2}} \left(y + \frac{d}{dy} \right)$$

$$b^\dagger b \Phi = \left(\eta - \frac{1}{2} \right) \Phi \quad \text{Schrödinger eq.} \quad (5)$$

$$[b, b^\dagger] = 1$$

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}} [b + b^\dagger] = \frac{1}{\sqrt{2}\alpha} [b + b^\dagger]$$

$$\hat{p} = -i\hbar \frac{d}{dx} = -i\hbar\alpha \frac{d}{dy} = -\frac{i\hbar\alpha}{\sqrt{2}} [b - b^\dagger]$$

16 / 27

Number operator

$$\hat{N} = b^\dagger b \quad (6)$$

$$\hat{H} = \hbar\omega \left[\hat{N} + \frac{1}{2} \right] \rightarrow \text{eigenfunctions of } \hat{H} = \text{eigenfunctions of } \hat{N}$$

Eigenvalues of $\hat{H} \rightarrow E_n = \hbar\omega(n + \frac{1}{2})$ (n are eigenvalues of \hat{N})

$$(5) \text{ and } (6) \quad \hat{N}\Phi_n = \left(\eta - \frac{1}{2} \right) \Phi_n = n \Phi_n \rightarrow n = \eta - \frac{1}{2}$$

(n = quanta of excitation, **phonons**)

17 / 27

- (i) (5) multiplied by b on the left

$$bb^\dagger b \Phi_n = n b \Phi_n \quad (bb^\dagger = 1 + b^\dagger b)$$

$$(1 + b^\dagger b)b \Phi_n = n b \Phi_n$$

$$b^\dagger b(b \Phi_n) = (n - 1) (b \Phi_n)$$

Φ_n eigenfunction of \hat{N} associated to eigenvalue $n \rightarrow b \Phi_n$ eigenfunction of \hat{N} associated to eigenvalue $n - 1$

$b =$ **annihilation operator** (annihilates one quantum of excitation = its number decreases by one unit)

18 / 27

Let's call the ground state $\Phi_{g.s.}$

$$b \Phi_{g.s.} = 0 \rightarrow (\hat{D} + y)\Phi_{g.s.} = 0 \quad (7)$$

$$\frac{d \Phi_{g.s.}}{dy} + y \Phi_{g.s.} = 0 \rightarrow \frac{d \Phi_{g.s.}}{dy} = -y \Phi_{g.s.} \rightarrow$$

$$\ln \Phi_{g.s.} = -\frac{y^2}{2} + A \rightarrow \Phi_{g.s.} = B e^{-\frac{y^2}{2}}$$

from (7) $\Phi_{g.s.}$ is eigenfunction of b associated to 0

$$\rightarrow b^\dagger b \Phi_{g.s.} = 0 \rightarrow \Phi_{g.s.} \text{ is eigenfunction of } \hat{N} \text{ associated to } 0 \rightarrow \Phi_0$$

$$\rightarrow n_{g.s.} = 0 \rightarrow n = 0, 1, 2, \dots$$

19 / 27

- (ii) (5) multiplied by b^\dagger on the left

$$b^\dagger b^\dagger b \Phi_n = n b^\dagger \Phi_n$$

$$b^\dagger (bb^\dagger - 1) \Phi_n = n b^\dagger \Phi_n$$

$$b^\dagger b [b^\dagger \Phi_n] = (n + 1) [b^\dagger \Phi_n]$$

$b^\dagger \Phi_n$ is eigenfunction of \hat{N} with eigenvalue $n + 1$

b^\dagger **creation operator**

$$\Phi_1 \propto b^\dagger \Phi_0 \quad ; \quad \Phi_n \propto (b^\dagger)^n \Phi_0$$

(\propto because the normalization of $(b^\dagger)^n \Phi_0$ is not guaranteed)

$$n = 0 \quad ; \quad \Phi_0 \propto e^{-\frac{y^2}{2}}$$

20 / 27

$$n = 1 \quad ; \quad \Phi_1 \propto b^\dagger \Phi_0 \propto y e^{-\frac{y^2}{2}}$$

$$n = 2 \quad ; \quad \Phi_2 \propto b^\dagger \Phi_1 \propto (2y^2 - 1) e^{-\frac{y^2}{2}} \dots$$

we get the Hermite polynomials

21 / 27

■ (iii) In Dirac's notation

$|n\rangle$ normalized eigenstate of \hat{N} and \hat{H}

$$\langle n|n'\rangle = \delta_{nn'}$$

◆ $(b^\dagger)^\dagger = b$ demonstration

$$\blacksquare \langle n|b^\dagger|m\rangle = \langle m|(b^\dagger)^\dagger|n\rangle^* \quad \forall m, n$$

$$\text{but } b^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle \quad \text{and } b|n\rangle = \sqrt{n}|n-1\rangle$$

$$\sqrt{m+1} \delta_{m+1,n} = \langle m|(b^\dagger)^\dagger|n\rangle^*$$

22 / 27

■

◆

■

$$\begin{aligned} \langle m|b|n\rangle^* &= [\sqrt{n} \delta_{m,n-1}]^* \\ &= \sqrt{m+1} \delta_{m+1,n} \\ &\quad \forall m, n \end{aligned}$$

$$\langle m|(b^\dagger)^\dagger|n\rangle^* = \langle m|b|n\rangle^* \rightarrow$$

$$(b^\dagger)^\dagger = b \rightarrow b^\dagger = (b)^\dagger$$

23 / 27

■

◆ $|n\rangle = A b^\dagger |n-1\rangle$

$$\begin{aligned}\langle n|n\rangle &= |A|^2 \langle n-1|bb^\dagger|n-1\rangle \\ &= |A|^2 \langle n-1|1 + b^\dagger b|n-1\rangle \\ &= |A|^2(1 + n - 1) = n |A|^2\end{aligned}$$

$$\rightarrow |A| = \frac{1}{\sqrt{n}}$$

Take real A $b^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle$

Analogously $b |n\rangle = \sqrt{n} |n-1\rangle$

24 / 27

■

◆

$$\begin{aligned}|n\rangle &= \frac{(b^\dagger)^n |0\rangle}{C} \\ &= \frac{(b^\dagger)^{n-1} b^\dagger |0\rangle}{C} \\ &= \frac{(b^\dagger)^{n-1} |1\rangle}{C} = \frac{(b^\dagger)^{n-2} \sqrt{2} |2\rangle}{C} \\ &= \frac{\sqrt{n} \cdots \sqrt{2} \sqrt{1} |n\rangle}{C} \rightarrow C = \sqrt{n!}\end{aligned}$$

$$|n\rangle = \frac{(b^\dagger)^n}{\sqrt{n!}} |0\rangle$$

25 / 27

- [1] D.J. Griffiths, "Introduction to Quantum Mechanics", ed. Pearson Education Inc., 2005
- [2] A.P.French and E.F. Taylor, Introducción a la Física Cuántica, ed. Reverté, 1982
- [3] Alastair I. M. Rae, Quantum mechanics, IOP, 2001
- [4] J. S. Townsend, "A Modern Approach to Quantum Mechanics", Univ. Science Books, 2000