

Lesson 2: Wave-particle duality

Clara E. Alonso Alonso

October 11, 2011

Photoelectric effect	2
.....	3
.....	4
.....	5
.....	6
Compton effect	7
.....	8
.....	9
.....	10
.....	11
.....	12
.....	13
.....	14
De Broglie hypothesis	15
.....	16
Electron diffraction: Davisson and Germer experiment. Thomson experiment	17
.....	18
.....	19
.....	20
.....	21
.....	22
.....	23
.....	24
.....	25
.....	26
.....	27
.....	28
.....	29
Wave properties of matter. Group velocity	30
.....	31
.....	32
.....	33
Wave packets and Schrödinger equation. Superposition principle	34
.....	35

.....	36
.....	37
.....	38
.....	39
.....	40
Bibliography	41
References	42

In 1887 Hertz observed

1. radiating **metals** with visible and U.V. radiation ($\nu \approx 10^{14} - 10^{17}$ Hz) \rightarrow emission of electrons
2. there is emission if $\nu > \nu_{threshold}$ or ν_{cutoff} ($\nu_{threshold}$ depends on the radiated metal)
3. with $\nu > \nu_{threshold}$ the e^- current is proportional to the intensity of the e.m. radiation
4. the maximum kinetic energy of the emitted electrons (photoelectrons)
 - doesn't depend on the e.m. radiation intensity
 - varies linearly with ν

3 / 42

Classically

- it is understood that there is e^- emission (1)
- it is not understood that there is $\nu_{threshold}$ (2)
- the energy of the incident wave (doesn't depends on ν) \propto source intensity (4)
- the delay time between the radiation arrival and the electron emission (larger when the intensity is smaller) \rightarrow is not observed experimentally (even whith $I \ll$)

Einstein \rightarrow radiation \rightarrow **energy quantum** $h\nu \rightarrow$ absorbed by an individual e^-
 - time for absorbing a quantum is smaller or similar to 10^{-9} s

4 / 42

$$h\nu = (E_c)_{max} + W$$

W **work function** (for e^- near the surface), depends on the metal

the e^- must surmount a potential energy step at the surface of the metal, they are confined
It explains

- $\exists \quad \nu_{threshold} = \frac{W}{h}$
- $(E_c)_{max}$ linear with ν
- proporcionalidad between e^- current and source intensity I

$I \gg \rightarrow$ > number of **photons** \rightarrow > number of **photoelectrons**


“particle” behaviour of light

5 / 42

- information on metals $W = h\nu_{threshold}$
($W \approx eV$)
 V_f stopping potential (the polarity of the voltage source is reversed)

$$(E_c)_{max} = e V_f = h\nu - W$$

$$V_f = \frac{h}{e}\nu - \frac{W}{e}$$




6 / 42

Scattering of radiation of $\lambda \simeq \text{\AA}$ (X rays) by metal foils

Classically $\rightarrow I(\theta) \propto 1 + \cos^2 \theta$ independent of λ_{inc}

Compton observes at scattering angle θ λ_{inc} and $\lambda_{inc} + \Delta\lambda(\theta)$



Compton incident radiation \rightarrow photons of energy $h\nu$

Compton effect: **elastic** scattering of e^- by photons


photon momentum?

Relativistic

$$E = \sqrt{(m_0 c^2)^2 + p^2 c^2}$$

$$v = \frac{dE}{dp} = \frac{pc^2}{E} = \frac{pc^2}{\sqrt{(m_0 c^2)^2 + p^2 c^2}}$$

for the photon $v = c \rightarrow m_0 = 0 \rightarrow p = \frac{h\nu}{c}$



10 / 42

■ Conservation of momentum

$$\vec{p} = \vec{p}' + \vec{p}_e$$

$$\vec{p}_e^2 = (\vec{p} - \vec{p}')^2 = p^2 + p'^2 - 2\vec{p} \cdot \vec{p}' \quad (1)$$

■ Conservation of energy

$$h\nu + mc^2 = h\nu' + (m^2c^4 + p_e^2c^2)^{\frac{1}{2}}$$

$$\begin{aligned} m^2c^4 + p_e^2c^2 &= (h\nu - h\nu' + mc^2)^2 \\ &= (h\nu - h\nu')^2 + 2mc^2(h\nu - h\nu') + m^2c^4 \end{aligned} \quad (2)$$

11 / 42

From (1) $p_e^2 = \left(\frac{h\nu}{c}\right)^2 + \left(\frac{h\nu'}{c}\right)^2 - 2\frac{h\nu}{c}\frac{h\nu'}{c}\cos\theta$

$$p = \frac{h\nu}{c} ; p' = \frac{h\nu'}{c}$$

$$p_e^2 c^2 = (h\nu - h\nu')^2 + 2(h\nu)(h\nu')(1 - \cos\theta) \quad (3)$$

From (2) and (3)

$$(h\nu)(1 - \cos\theta) = \frac{mc^2}{\nu'}(\nu - \nu')$$

$$\Delta\lambda = \lambda' - \lambda = \frac{h}{mc}(1 - \cos\theta)$$

$$\Delta\lambda \geq 0$$


12 / 42

λ_{inc} observed in expt. \rightarrow scattering by atom $m \rightarrow m_{at} \rightarrow \Delta\lambda \rightarrow 0$

$$\lambda_c = \frac{h}{m_e c} = 0.024 \text{ \AA} \quad \text{Compton wavelength}$$

- Experimentally \rightarrow agreement in e^- recoil
- Simultaneity between recoil e^- and outgoing photon
- Scattering interpreted as the one of billiard balls \rightarrow light could be regarded as classical particles in these experiments
- But radiation also has wave properties (**interference, diffraction**)

13 / 42



Photon self-identity issues

14 / 42

De Broglie hypothesis

15 / 42

1923 De Broglie suggests double nature **wave-particle** of matter

De Broglie relation

$$\lambda = \frac{h}{p}$$

plus Einstein relation

$$E = \hbar\omega$$

$$\hbar = \frac{h}{2\pi} \quad ; \quad \hbar c = 1970 \text{ eV}\text{\AA}$$

He proposes to observe e^- diffraction ($E_c \sim \text{tens of eV}$)

⇓


16 / 42

Electron diffraction: Davisson and Germer experiment. Thomson experiment


17 / 42

DAVISSON-GERMER EXPERIMENT


e^- scattering by Ni crystal (shows preferential directions)



18 / 42



19 / 42




$$\lambda_{DB} = \frac{h}{\sqrt{2m_e V e}} \rightarrow \lambda_{DB}(\text{\AA}) = \left(\frac{150}{V}\right)^{\frac{1}{2}} \quad (V \text{ in volts})$$

(no relativistic; $m_e = 0.511 \frac{\text{MeV}}{c^2}$)


20 / 42

- For $V = 54 \text{ V} \rightarrow \lambda_{DB} = 1.66 \text{ \AA}$ (for other V less intense phenomenon)

Fig. 2-6 Polar plots of scattered intensity showing dependence of diffraction pattern on accelerating voltage. (Based on data of Davisson and Germer.)




21 / 42



If the effect depended only on the rows of atoms in the surface \rightarrow any voltage $V \rightarrow$ strong diffraction peak at $\phi = \sin^{-1} \frac{\lambda}{D}$ Electrons of E very different from $54 \text{ eV} \rightarrow$ much reduced diffraction intensities

22 / 42

Observations explained by \rightarrow **reflections from atomic planes**




$$d = D \sin \alpha \quad \text{and} \quad D = 2.15 \text{ \AA}$$


$$\theta = \frac{\pi}{2} - \alpha \quad ; \quad \phi = 2\alpha \quad ;$$

$$\alpha = 25^\circ \rightarrow \phi = 50^\circ \quad ; \quad d = 0.91 \text{ \AA}$$

23 / 42



24 / 42



$$l = d \sin \theta$$

path difference

$$\delta = 2l = 2d \sin \theta = 2 \times 0.91 \text{ \AA} \times \sin 65^\circ = 1.65 \text{ \AA}$$

path difference $= n\lambda$ **Bragg's law** (condition of reinforcement)

$$n = 1 \rightarrow \lambda = 2d \sin \theta = D \sin \phi = 1.65 \text{ \AA} \text{ (first order diffraction maximum)}$$

25 / 42

Expressions necessary to obtain the last relation of previous page

$$d = D \sin \alpha = D \cos \theta$$


$$2d \sin \theta = 2D \cos \theta \sin \theta = D \sin 2\theta = D \sin \phi$$

26 / 42

Thomson experiment (1927-28)

- e^- wave properties are observed
 - Differs from Davisson-Germer experiment:
 - ◆ Energy of $e^- \rightarrow 10 - 40 \text{ KeV} \rightarrow \lambda_{DB} \approx 0.1 \text{ \AA}$ (D.G. $\rightarrow 30 - 600 \text{ eV} \rightarrow \lambda_{DB} \approx 1 \text{ \AA}$) $\rightarrow e^-$ much more penetrating, pass through thin films ($\approx 1000 \text{ \AA}$ thickness) \rightarrow TRANSMISSION
 - ◆ Many hundred of atomic planes contribute to the diffracted wave
 - ◆ Polycrystal (aggregate of very small crystals oriented at random)
 - ◆ Independently confirms the De Broglie relation
- Diffraction pattern are concentric circles


27 / 42



Electron diffraction by gold crystals

28 / 42

- Diffracted beams are collected on photographic plates located far from the diffracting film
- Results similar to those of X-rays (classically is radiation)
- This experiment verifies the applicability of the De Broglie relation to higher energies




29 / 42

De Broglie assigns λ to particles

Which is the responsible wave?

Ej. Representation of $\Psi(x, t)$ versus x at an instant t



We expect something like this plot for particles moving in the x -axis

- Modulate amplitude
- Different from zero in finite region of x -axis

31 / 42

Plane wave $\rightarrow e^{i(\vec{k} \cdot \vec{r} - \omega t)}$ **is not localized** in space (it has plane wavefronts, \vec{k} vector of module $\frac{2\pi}{\lambda}$ and direction the one of the wave propagation)

The need arises for wavepackets (superposition of plane waves with different amplitudes)

$$\Psi(\vec{r}, t) = \frac{1}{(2\pi\hbar)^{\frac{3}{2}}} \int d^3\vec{p} A(\vec{p}) e^{\frac{i(\vec{r} \cdot \vec{p} - E(\vec{p})t)}{\hbar}}$$

$$A(\vec{p}) = \frac{1}{(2\pi\hbar)^{\frac{3}{2}}} \int d^3\vec{r} \Psi(\vec{r}, 0) e^{-\frac{i\vec{r} \cdot \vec{p}}{\hbar}}$$

where $\vec{p} = \hbar\vec{k}$ (De Broglie) and $E = \hbar\omega = \frac{p^2}{2m}$

(superimposed plane waves have different λ and ν)

32 / 42

Group velocity $v_g = \frac{d\omega(k)}{dk}|_{\bar{k}} = 2\pi \frac{d\nu}{dk}|_{\bar{k}}$ ($\neq v_\Phi = \frac{\omega}{k}$ monochromatic waves, phase velocity)

$$\omega = \frac{\hbar k^2}{2m} \rightarrow v_g = \frac{\hbar \bar{k}}{m} = \frac{\bar{p}}{m} = \bar{v}$$

$$v_p = \frac{dE}{dp} = \frac{\hbar k_p}{m} \quad (\text{particle})$$

we want $v_g = v_p$

it is achieved if the wave packet is chosen with $\bar{k} = \frac{mv_p}{\hbar}$

33 / 42

Wave packets and Schrödinger equation. Superposition principle 34 / 42

Achievements with the wavepacket

- location in space
- $v_g = v$ of the particle

we seek the wave equation whose solution is the wavepacket

Conditions

- Linear equation (so that the **superposition principle** applies) \rightarrow it contains the first power of $\Psi, \frac{\partial \psi}{\partial x_i}, \frac{\partial^2 \psi}{\partial x_i^2}, \frac{\partial \psi}{\partial t}, \frac{\partial^2 \psi}{\partial t^2} \dots$

Linear \rightarrow If Ψ_1 and Ψ_2 they are solutions of the equation $\Psi = c_1 \Psi_1 + c_2 \Psi_2$ it is also solution
 $c_1, c_2 \in \mathcal{C}$

35 / 42

Wavepacket = linear combination of plane waves

↓

the plane wave must be solution of the wave equation sought

We postulate the Schrödinger equation

$$i\hbar \frac{\partial \Psi}{\partial t} = \left(\frac{-\hbar^2}{2m} \nabla^2 + V \right) \Psi$$

$$\nabla^2 = \Delta = \sum_i \frac{\partial^2}{\partial x_i^2} \quad (\text{laplacian operator})$$

36 / 42

The Schrödinger equation is valid for nonrelativistic particle of mass m subjected to potential V

Free particle $V = 0 \rightarrow \Psi = e^{i(\vec{k} \cdot \vec{r} - \omega t)}$ is solution of Schrödinger eq. with

$$\omega = \frac{E}{\hbar} \quad ; \quad E = \frac{p^2}{2m} = \frac{\hbar^2 k^2}{2m}$$

Matter waves are superimposed linearly as the classical \rightarrow there should result interference phenomena
(eg, Davisson-Germer, double slit)

37 / 42

If Ψ_1 and Ψ_2 describe physical situations

↓

also $\Psi = c_1\Psi_1 + c_2\Psi_2$; $c_1, c_2 \in \mathcal{C}$

Waves are added, not intensities

$$I \propto |\Psi|^2$$

$$\Psi_1 \rightarrow I_1 \propto |\Psi_1|^2 \quad ; \quad \Psi_2 \rightarrow I_2 \propto |\Psi_2|^2$$

$$\Psi_1, \Psi_2 \rightarrow I_t \propto |\Psi_1 + \Psi_2|^2$$

interference

$$I_t \neq I_1 + I_2$$

38 / 42

The superposition principle explains interference and diffraction → we adopt it for quantum mechanics

- If we have a plane wave associated with a free particle moving in one dimension in the direction of increasing x

$$\Psi_1(x, t) = A e^{i(kx - \omega t)}$$

- idem moving in one dimension in the direction of decreasing x

$$\Psi_2(x, t) = A e^{i(-kx - \omega t)}$$

39 / 42

If in $t = 0$

$$\Psi(x, 0) = \Psi_1(x, 0) + \Psi_2(x, 0) = 2A \cos kx = A \left(e^{ikx} + e^{-ikx} \right)$$

by the principle of superposition of waves they evolve independently in time

$$\Psi(x, t) = A \left(e^{i(kx - \omega t)} + e^{-i(kx + \omega t)} \right)$$

Ψ is complex in general. Intensities $\propto |\Psi|^2$ are real

40 / 42

Bibliography

41 / 42

References

- [1] S. Gasiorowicz, "Quantum Physics", ed. John Wiley, 2003
- [2] D.J. Griffiths, "Introduction to Quantum Mechanics", ed. Pearson Education Inc., 2005
- [3] D. Park, "Introduction to the quantum theory", ed. McGraw-Hill, 1992
- [4] A.P. French and E.F. Taylor, "Introducción a la Física Cuántica", ed. Reverté, 1982
- [5] R. Eisberg and R. Resnick, "Física cuántica", ed. Limusa, 2004
- [6] J.L. Basdevant and J. Dalibard, "Quantum Mechanics", ed. Springer, 2002

42 / 42