



从回旋加速器到费米梦想：
探幽入微之路

1、回旋加速器

密封在真空中的两个金属盒（ D_1 和 D_2 ）放在电磁铁两极间的强磁场中，两盒间接有交流电源，它在缝隙里的交变电场用以加速带电粒子。



(a)




(b)

交变电场的周期恰好为回旋周期时 — 粒子绕过半圈恰好电场反向，粒子又被加速。
因为回旋周期与半径无关，
所以粒子可被反复加速。

回旋频率

$$f = \frac{1}{T} = \frac{qB}{2\pi m}$$


动画演示



$$T = 2\pi \frac{m}{qB}$$

当粒子到达半圆边缘时，
粒子的最大速率为
(R_0 为最大半径)

$$v = \frac{qBR_0}{m} \quad R = \frac{mv_0}{qB}$$




粒子最大动能

$$E_k = \frac{1}{2}mv^2 = \frac{1}{2}m\left(\frac{BqR_0}{m}\right)^2 = \frac{q^2B^2R_0^2}{2m}$$





1932年,E . O . Lawrence
建造的第一台回旋加速器



1932年劳伦斯研制第一台回旋加速器的D型室.


此加速器可将质子和氘核加速到0.8MeV的能量

美国物理学家劳伦斯于1934年研制成功第一台实用加速器，于1939年获诺贝尔物理学奖。



回旋加速器一般只能将质子加速到25MeV左右。

考虑到狭义相对论效应发展的同步回旋加速器可以将提高到1GeV。



2、LHC



欧洲核子研究中心**CERN**，大环是直径27km的大型强子对撞机**LHC**，中环是质子同步加速器。加速能量为**14TeV**。

欧洲大型强子对撞机工作原理示意图

模拟宇宙大爆炸发生时的状态，有助人类进一步探索宇宙起源之谜

项目位于瑞士和法国边境地区


已耗时：将近20年

已耗资：数十亿美元

参与相关实验项目




80多个国家和地区的



7000多名科学家




THE PRINCIPAL MACHINE COMPONENTS OF AN ACCELERATOR


粒子加速器的主要构成







Francois Englert和Peter W. Higgs的发现获得 2013年诺贝尔物理学奖



标准粒子模型理论中共预言了62种基本粒子的存在，希格斯玻色子正是该理论依赖的基石。作为物质的质量之源, Higgs粒子被叫做“上帝粒子”

The Standard Model

Binding of nuclei and radioactivity require two additional short-range forces:

- **Strong Interactions:** Keep nucleus bound.
- **Weak interactions:** Allow beta decay of nuclei

The diagram illustrates the Standard Model of particle physics, organized by generation and type of particle.

Three generations of matter (fermions)

	I	II	III
Quarks	mass → 2.4 MeV/c ² charge → 2/3 spin → 1/2 u up	mass → 1.27 GeV/c ² charge → 2/3 spin → 1/2 c charm	mass → 171.2 GeV/c ² charge → 2/3 spin → 1/2 t top
	mass → 4.8 MeV/c ² charge → -1/3 spin → 1/2 d down	mass → 104 MeV/c ² charge → -1/3 spin → 1/2 s strange	mass → 4.2 GeV/c ² charge → -1/3 spin → 1/2 b bottom
Leptons	mass → <2.2 eV/c ² charge → 0 spin → 1/2 v_e electron neutrino	mass → <0.17 MeV/c ² charge → 0 spin → 1/2 v_μ muon neutrino	mass → <15.5 MeV/c ² charge → 0 spin → 1/2 v_τ tau neutrino
	mass → 0.511 MeV/c ² charge → -1 spin → 1/2 e electron	mass → 105.7 MeV/c ² charge → -1 spin → 1/2 μ muon	mass → 1.777 GeV/c ² charge → -1 spin → 1/2 τ tau

Gauge bosons

	Mass	Charge	Spin	Symbol	Name
Higgs	125 GeV	0	0	H	Higgs
photon	0	0	1	γ	photon
gluon	0	0	1	g	gluon
Z boson	91.2 GeV/c ²	0	1	Z⁰	Z boson
W boson	80.4 GeV/c ²	±1	1	W[±]	W boson

3 families of matter (chiral fermions)

Quarks : $Q_L \equiv \begin{pmatrix} u_L \\ d_L \end{pmatrix}, d_R, u_R$

Leptons : $L_L \equiv \begin{pmatrix} \nu_{eL} \\ e_L \end{pmatrix}$, e_R , ν_R ?

3 types of gauge bosons (spin 1)




- Strong (8 massless gluons, g)
- Electromag. (1 massless photon γ)
- Weak (3 massive Z , W^+ , W^-)

1 Higgs boson (spin 0) needed for SSB

$$\begin{aligned}
\mathcal{L} = & -\frac{1}{4} \sum_{C=1}^8 G_{\mu\nu}^C G^{C\mu\nu} - \frac{1}{4} \sum_{a=1}^3 W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} \\
& + D^\mu H^\dagger D_\mu H - V(H^\dagger H) \\
& + \sum_{n,i,\alpha} i Q_{ni\alpha}^\dagger \bar{\sigma}^\mu D_\mu Q_n^{i\alpha} + \sum_{n,i} i U_n^{\dagger i} \bar{\sigma}^\mu D_\mu U_{ni} + \sum_{n,i} i D_n^{\dagger i} \bar{\sigma}^\mu D_\mu D_{ni} \\
& + \sum_{n,\alpha} i L_{n\alpha}^\dagger \bar{\sigma}^\mu D_\mu L_n^\alpha + \sum_n i E_n \bar{\sigma}^\mu D_\mu E_n \\
& + \mathcal{L}_{\text{Yukawa}} + \mathcal{L}_{LLHH} .
\end{aligned}$$

Here $G_{\mu\nu}^C$, $W_{\mu\nu}^a$, and $B_{\mu\nu}$ are canonically normalized tension fields for the $SU(3)$, $SU(2)$, and $U(1)$ gauge symmetries,

$$\begin{aligned}
G_{\mu\nu}^C &= \partial_\mu G_\nu^C - \partial_\nu G_\mu^C - g_3 f^{CDE} G_\mu^D G_\nu^E , \\
W_{\mu\nu}^a &= \partial_\mu W_\nu^a - \partial_\nu W_\mu^a - g_2 F \epsilon^{abc} W_\mu^b W_\nu^c , \\
B_{\mu\nu} &= \partial_\mu B_\nu - \partial_\nu B_\mu ,
\end{aligned} \tag{9}$$



Intro: 750 GeV diphoton excess awakens!!


- [1] K. Harigaya and Y. Nomura, "Composite Models for the 750 GeV Diphoton Excess," [arXiv:1512.04850 \[hep-ph\]](#).
- [2] Y. Mambrini, G. Arcadi, and A. Djouadi, "The LHC diphoton resonance and dark matter," [arXiv:1512.04913 \[hep-ph\]](#).
- [3] M. Bachowicz, A. Mariotti, and D. Redigolo, "Di-photon excess illuminates Dark Matter," [arXiv:1512.04917 \[hep-ph\]](#).
- [4] A. Angelescu, A. Djouadi, and G. Moreau, "Scenarios for interpretations of the LHC diphoton excess: two Higgs doublets and vector-like quarks and leptons," [arXiv:1512.04921 \[hep-ph\]](#).
- [5] Y. Nakai, R. Sato, and K. Tobioka, "Footprints of New Strong Dynamics via Anomaly," [arXiv:1512.04924 \[hep-ph\]](#).
- [6] S. Knapen, T. Melia, M. Papucci, and K. Zurek, "Rays of light from the LHC," [arXiv:1512.04928 \[hep-ph\]](#).
- [7] D. Buttazzo, A. Greljo, and D. Marzocca, "Knocking on New Physics' door with a Scalar Resonance," [arXiv:1512.04929 \[hep-ph\]](#).
- [8] A. Pilaftsis, "Diphoton Signatures from Heavy Axion Decays at LHC," [arXiv:1512.04931 \[hep-ph\]](#).
- [9] R. Franceschini, G. F. Giudice, J. F. Kamenik, M. McCullough, A. Pomarol, R. Rattazzi, M. Rodi, F. Riva, A. Strumia, and R. Torre, "What is the gamma gamma resonance at 750 GeV?," [arXiv:1512.04933 \[hep-ph\]](#).
- [10] S. Di Chiara, L. Marzola, and M. Raidal, "First interpretation of the 750 GeV di-photon resonance at the LHC," [arXiv:1512.04939 \[hep-ph\]](#).
- [11] T. Higaki, K. S. Jeong, N. Kitajima, and F. Takahashi, "The QCD Axion from Aligned Axions and Diphoton Excess," [arXiv:1512.05295 \[hep-ph\]](#).
- [12] S. D. McDermott, P. Meade, and H. Ramani, "Singlet Scalar Resonances and the Diphoton Excess," [arXiv:1512.05326 \[hep-ph\]](#).
- [13] J. Ellis, S. A. R. Ellis, J. Quevillon, V. Sanz, and T. You, "On the Interpretation of a Possible ~ 750 GeV Particle Decaying into $\gamma\gamma$," [arXiv:1512.05327 \[hep-ph\]](#).
- [14] M. Low, A. Tesi, and L.-T. Wang, "A pseudoscalar decaying to photon pairs in the early LHC run 2 data," [arXiv:1512.05328 \[hep-ph\]](#).
- [15] B. Bellazzini, R. Franceschini, F. Sala, and J. Serra, "Goldstones in Diphotons," [arXiv:1512.05330 \[hep-ph\]](#).
- [16] R. S. Gupta, S. Jager, Y. Kats, G. Perez, and E. Stamou, "Interpreting a 750 GeV Diphoton Resonance," [arXiv:1512.05332 \[hep-ph\]](#).
- [17] C. Petersson and R. Torre, "The 750 GeV diphoton excess from the goldstino superpartner," [arXiv:1512.05333 \[hep-ph\]](#).
- [18] E. Molinaro, F. Sannino, and N. Vignaroli, "Strong dynamics or axion origin of the diphoton excess," [arXiv:1512.05334 \[hep-ph\]](#).
- [19] A. Falkowski, O. Slone, and T. Volansky, "Phenomenology of a 750 GeV Singlet," [arXiv:1512.05777 \[hep-ph\]](#).
- [20] B. Dutta, Y. Gao, T. Ghosh, I. Gogoladze, and T. Li, "Interpretation of the diphoton excess at CMS and ATLAS," [arXiv:1512.05439 \[hep-ph\]](#).
- [21] Q.-H. Cao, Y. Liu, K.-P. Xie, B. Yan, and D.-M. Zhang, "A Boost Test of Anomalous Diphoton Resonance at the LHC," [arXiv:1512.05542 \[hep-ph\]](#).
- [22] S. Matsuzaki and K. Yamawaki, "750 GeV Diphoton Signal from One-Family Walking Technipion," [arXiv:1512.05564 \[hep-ph\]](#).
- [23] A. Kobakhidze, F. Wang, L. Wu, J. M. Yang, and M. Zhang, "LHC diphoton excess
- [33] P. Agrawal, J. Fan, B. Heidenreich, M. Reece, and M. Strassler, "Experimental Considerations Motivated by the Diphoton Excess at the LHC," [arXiv:1512.05775 \[hep-ph\]](#).
- [36] C. Csaki, J. Hubisz, and J. Terning, "The Minimal Model of a Diphoton Resonance: Production without Gluon Couplings," [arXiv:1512.05776 \[hep-ph\]](#).
- [37] D. Aloni, K. Blum, A. Dery, A. Efrati, and Y. Nir, "On a possible large width 750 GeV diphoton resonance at ATLAS and CMS," [arXiv:1512.05778 \[hep-ph\]](#).
- [38] Y. Bai, J. Berger, and R. Lu, "A 750 GeV Dark Pion: Cousin of a Dark G-parity-odd WIMP," [arXiv:1512.05779 \[hep-ph\]](#).
- [39] E. Gabrielli, K. Kannike, B. Mele, M. Raidal, C. Spethmann, and H. Veermäe, "A SUSY Inspired Simplified Model for the 750 GeV Diphoton Excess," [arXiv:1512.05961 \[hep-ph\]](#).
- [40] R. Benbrik, C.-H. Chen, and T. Nomura, "Higgs singlet as a diphoton resonance in a vector-like quark model," [arXiv:1512.06028 \[hep-ph\]](#).
- [41] J. S. Kim, J. Reuter, K. Roliecki, and R. R. de Austri, "A resonance without resonance: scrutinizing the diphoton excess at 750 GeV," [arXiv:1512.06083 \[hep-ph\]](#).
- [42] A. Alves, A. G. Dias, and K. Sinha, "The 750 GeV S-cion: Where else should we look for it?," [arXiv:1512.06091 \[hep-ph\]](#).
- [43] E. Megias, O. Pujolas, and M. Quiros, "On dilatons and the LHC diphoton excess," [arXiv:1512.06106 \[hep-ph\]](#).
- [44] L. M. Carpenter, R. Colburn, and J. Goodman, "Supersoft SUSY Models and the 750 GeV Diphoton Excess, Beyond Effective Operators," [arXiv:1512.06107 \[hep-ph\]](#).
- [45] J. Bernon and C. Smith, "Could the width of the diphoton anomaly signal a three-body decay?," [arXiv:1512.06113 \[hep-ph\]](#).
- [46] W. Chao, "Symmetries Behind the 750 GeV Diphoton Excess," [arXiv:1512.06297 \[hep-ph\]](#).
- [47] M. T. Arun and P. Saha, "Gravitons in multiply warped scenarios - at 750 GeV and beyond," [arXiv:1512.06335 \[hep-ph\]](#).
- [48] C. Han, H. M. Lee, M. Park, and V. Sanz, "The diphoton resonance as a gravity mediator of dark matter," [arXiv:1512.06376 \[hep-ph\]](#).
- [49] S. Chang, "A Simple U(1) Gauge Theory Explanation of the Diphoton Excess," [arXiv:1512.06426 \[hep-ph\]](#).
- [50] I. Chakraborty and A. Kundu, "Diphoton excess at 750 GeV: Singlet scalars confront naturalness," [arXiv:1512.06508 \[hep-ph\]](#).
- [51] R. Ding, L. Huang, T. Li, and B. Zhu, "Interpreting 750 GeV Diphoton Excess with R-parity Violation Supersymmetry," [arXiv:1512.06560 \[hep-ph\]](#).
- [52] H. Pan, S. Wang, and S. Zheng, "Scalar Dark Matter Explanation of Diphoton Excess at LHC," [arXiv:1512.06562 \[hep-ph\]](#).
- [53] X.-F. Han and L. Wang, "Implication of the 750 GeV diphoton resonance on two-Higgs-doublet model and its extensions with Higgs field," [arXiv:1512.06587 \[hep-ph\]](#).
- [54] M.-x. Luo, K. Wang, T. Xu, L. Zhang, and G. Zhu, "Squarkonium/Diquarkonium and the Diphoton Excess," [arXiv:1512.06670 \[hep-ph\]](#).
- [55] J. Chang, K. Cheung, and C.-T. Lu, "Interpreting the 750 GeV Di-photon Resonance using photon-jets in Hidden-Valley-like models," [arXiv:1512.06671 \[hep-ph\]](#).
- [56] D. Bardhan, D. Bhatia, A. Chakraborty, U. Maitra, S. Raychaudhuri, and T. Samui, "Radion Candidate for the LHC Diphoton Resonance," [arXiv:1512.06674 \[hep-ph\]](#).
- [57] T.-F. Feng, X.-Q. Li, H.-B. Zhang, and S.-M. Zhao, "The LHC 750 GeV diphoton excess supersymmetry with gauged baryon and lepton numbers," [arXiv:1512.06696 \[hep-ph\]](#).
- [58] O. Antipin, M. Mojaza, and F. Sannino, "A natural Coleman-Weinberg theory explains the diphoton excess," [arXiv:1512.06708 \[hep-ph\]](#).
- [59] F. Wang, L. Wu, J. M. Yang, and M. Zhang, "750 GeV Diphoton Resonance, 125 GeV Higgs and Muon g-2 Anomaly in Deflected Anomaly Mediation SUSY Breaking Scenario," [arXiv:1512.06715 \[hep-ph\]](#).
- [60] J. Cao, C. Han, L. Shang, W. Su, J. M. Yang, and Y. Zhang, "Interpreting the 750 GeV diphoton excess by the singlet extension of the Manohar-Wise Model," [arXiv:1512.06722 \[hep-ph\]](#).
- [61] F. P. Huang, C. S. Li, Z. L. Liu, and Y. Wang, "750 GeV Diphoton Excess from Cascade Decay," [arXiv:1512.06732 \[hep-ph\]](#).
- [62] W. Liao and H.-q. Zheng, "Scalar resonance at 750 GeV as composite of heavy vector-like fermions," [arXiv:1512.06741 \[hep-ph\]](#).
- [63] J. J. Heckman, "750 GeV Diphotons from a D3-brane," [arXiv:1512.06773 \[hep-ph\]](#).
- [64] M. Dhuria and G. Goswami, "Perturbativity, vacuum stability and inflation in the light of 750 GeV diphoton excess," [arXiv:1512.06782 \[hep-ph\]](#).
- [65] X.-J. Bi, Q.-F. Xiang, P.-F. Yin, and Z.-H. Yu, "The 750 GeV diphoton excess at the LHC and dark matter constraints," [arXiv:1512.06787 \[hep-ph\]](#).
- [66] J. S. Kim, K. Roliecki, and R. R. de Austri, "Model-independent combination of diphoton constraints at 750 GeV," [arXiv:1512.06797 \[hep-ph\]](#).
- [67] L. Berthier, J. M. Cline, W. Shepherd, and M. Trost, "Effective interpretations of a diphoton excess," [arXiv:1512.06799 \[hep-ph\]](#).
- [68] W. S. Cho, D. Kim, K. Kong, S. H. Lim, K. T. Matchev, J.-C. Park, and M. Park, "The 750 GeV Diphoton Excess May Not Imply a 750 GeV Resonance," [arXiv:1512.06824 \[hep-ph\]](#).
- [69] J. M. Cline and Z. Liu, "LHC diphotons from electroweakly pair-produced composite pseudoscalars," [arXiv:1512.06827 \[hep-ph\]](#).
- [70] M. Bauer and M. Neubert, "Flavor Anomalies, the Diphoton Excess and a Dark Matter Candidate," [arXiv:1512.06828 \[hep-ph\]](#).
- [71] M. Chala, M. Duerr, F. Kahlhoefer, and K. Schmidt-Hoberg, "Tricking Landau-Yang: How to obtain the diphoton excess from a vector resonance," [arXiv:1512.06833 \[hep-ph\]](#).
- [72] D. Barducci, A. Goudelis, S. Kulkarni, and D. Sengupta, "One jet to rule them all: monojet constraints and invisible decays of a 750 GeV diphoton resonance," [arXiv:1512.06842 \[hep-ph\]](#).
- [73] G. M. Pelaggi, A. Strumia, and E. Vigiani, "Trifunctional can explain the di-photon and di-boson LHC anomalies," [arXiv:1512.07225 \[hep-ph\]](#).
- [74] S. M. Boucenna, S. Morisi, and A. Vicente, "The LHC diphoton resonance from gauge symmetry," [arXiv:1512.06878 \[hep-ph\]](#).
- [75] C. W. Murphy, "Vector Leptoquarks and the 750 GeV Diphoton Resonance at the LHC," [arXiv:1512.06976 \[hep-ph\]](#).
- [76] A. E. C. Hernández and I. Nisandzic, "LHC diphoton 750 GeV resonance as an indication of $SU(3)_c \times SU(3)_L \times U(1)_1$ gauge symmetry," [arXiv:1512.07165 \[hep-ph\]](#).

探索基础物理学的未解之谜

- 粒子是否有相对应的超对称（SUSY）粒子存在？
- 有更高维度的空间(Kaluza-Klein theory, extra-dimensions) 存在吗？
- 是否可以探测到和弦论预言有关的物理现象？
- 为何物质与反物质是不对称的？
- 宇宙有96%的质量是目前天文学上无法观测到的（暗物质），这些到底是什么？
- 为何万有引力比起其他三个基本作用力（电磁力，强作用力，弱作用力）差了这么多个数量级？
- 微型黑洞是否存在？
- 是否存在超出标准模型的物理？
- 模拟早期宇宙的物理性质
-

3、中国的加速器

1958年6月30号中国第一个回旋加速器建成



北京正负电子对撞机: 长度0.24公里, 加速能量1-1.15GeV, 改造后能量最大为4Gev

- | | | |
|----------------|-----------------|----------|
| 1. 2. 第一对撞点实验厅 | 3. 储存环电源厅、中央控制室 | |
| 4. 高频站 | 5. 第二对撞点实验厅 | 6. 储存环隧道 |
| 7. 输运线隧道 | 8. 直线加速器隧道 | 9. 速调管走廊 |
| 10. 核物理实验厅 | 11. 输运线、电源厅 | |
| 12. 同步辐射实验东厅 | 13. 同步辐射实验西厅 | 14. 计算中心 |



- | | |
|---|---------------------------------|
| 1. 2. 1st. I.R. Experi. hall | |
| 3. Power Station of ring mag. and computer center | |
| 4. RF Station | 5. 2nd I.R. Experi. hall |
| 6. Tunnel of storage ring | 7. Tunnel of Trans. line |
| 8. Tunnel of Linac | 9. Klystron gallery |
| 10. Nuclear phy. Experi. hall | 11. Power sta. of trans. line |
| 12. East hall for S. R. experi. | 13. West hall for S. R. experi. |
| 14. Computer center | |








2013年中国电子对撞机 发现4夸克物质



a) pion



b) proton



c) $Z_c(3900)$



四夸克物质 $Z_c(3900)$ 的共振结构



Study of $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and Observation of a Charged Charmoniumlike State at Belle

Z. Q. Liu,¹⁷ C. P. Shen,^{36,*} C. Z. Yuan,¹⁷ I. Adachi,¹³ H. Aihara,⁵⁹ D. M. Asner,⁴⁶ V. Aulchenko,³ T. Aushev,²¹ T. Aziz,⁵⁴ A. M. Bakich,⁵³ A. Bala,⁴⁷ K. Belous,¹⁹ V. Bhardwaj,³⁸ B. Bhuyan,¹⁵ M. Bischofberger,³⁸ A. Bondar,³ G. Bonvicini,⁶⁵ A. Bozek,⁴² M. Bračko,^{31,22} J. Brodzicka,⁴² T. E. Browder,¹² P. Chang,⁴¹ V. Chekelian,³² A. Chen,³⁹ P. Chen,⁴¹ B. G. Cheon,¹¹ R. Chistov,²¹ K. Cho,²⁶ V. Chobanova,³² S.-K. Choi,¹⁰ Y. Choi,⁵² D. Cinabro,⁶⁵ J. Dalseno,^{32,55} M. Danilov,^{21,34} Z. Doležal,⁴ Z. Drásal,⁴ A. Drutskoy,^{21,34} D. Dutta,¹⁵ S. Eidelman,³ D. Epifanov,⁵⁹ H. Farhat,⁶⁵ J. E. Fast,⁴⁶ M. Feindt,²⁴ T. Ferber,⁶ A. Frey,⁹ V. Gaur,⁵⁴ N. Gabyshev,³ S. Ganguly,⁶⁵ R. Gillard,⁶⁵ Y. M. Goh,¹¹ B. Golob,^{30,22} J. Haba,¹³ K. Hayasaka,³⁷ H. Hayashii,³⁸ Y. Hori,³⁷ Y. Hoshi,³⁷ W.-S. Hou,⁴¹ Y. B. Hsiung,⁴¹ H. J. Hyun,²⁸ T. Iijima,^{37,36} K. Inami,³⁶ A. Ishikawa,⁵⁸ R. Itoh,¹³ Y. Iwasaki,¹³ D. Joffe,²⁵ T. Julius,³³ D. H. Kah,²⁸ J. H. Kang,⁶⁷ T. Kawasaki,⁴⁴ C. Kiesling,³² H. J. Kim,²⁸ J. B. Kim,²⁷ M. J. Kim,²⁶ K. T. Kim,²⁷ M. J. Kim,²⁶ Y. J. Kim,²⁶ K. Kinoshita,⁵ J. Klucar,²² B. R. Ko,²⁷ P. Kodyš,⁴ S. Korpar,^{31,22} P. Krizan,^{30,22} P. Krokovny,³ T. Kuhr,²⁴ Y.-J. Kwon,⁶⁷ J. S. Lange,⁷ S.-H. Lee,²⁷ J. Li,⁵¹ Y. Li,⁶⁴ J. Libby,¹⁶ C. Liu,⁵⁰ P. Lukin,³ D. Matvienko,³ K. Miyabayashi,³⁸ H. Miyata,⁴⁴ R. Mizuk,^{21,34} G. B. Mohanty,⁵⁴ A. Moll,^{32,55} R. Mussa,²⁰ E. Nakano,⁴⁵ M. Nakao,¹³ H. Nakazawa,³⁹ Z. Natkaniec,⁴² M. Nayak,¹⁶ E. Nedelkovska,³² N. K. Nisar,⁵⁴ S. Nishida,¹³ O. Nitoh,⁶² S. Ogawa,⁵⁶ S. Okuno,²³ S. L. Olsen,⁵¹ Y. Onuki,⁵⁹ W. Ostrowicz,⁴² C. Oswald,² P. Pakhlov,^{21,34} G. Pakhlova,²¹ H. Park,²⁸ H. K. Park,²⁸ T. K. Pedlar,⁶⁸ R. Pestotnik,²² M. Petrič,²² L. E. Piihonien,⁶⁴ M. Ritter,³² M. Röhrken,²⁴ A. Rostomyan,⁶ H. Sahoo,¹² T. Saito,⁵⁸ Y. Sakai,¹³ S. Sandilya,⁵⁴ D. Santel,⁵ T. Sanuki,⁵⁸ Y. Sato,⁵⁸ V. Savinov,⁴⁸ O. Schneider,²⁹ G. Schnell,^{1,14} C. Schwanda,¹⁸ R. Seidl,⁴⁹ D. Semmler,⁷ K. Senyo,⁶⁶ O. Seon,³⁶ M. E. Sevir,³³ M. Shapkin,¹⁹ T.-A. Shibata,⁶⁰ J.-G. Shiu,⁴¹ B. Shwartz,³ A. Sibidanov,⁵³ F. Simon,^{32,55} P. Smerkol,²² Y.-S. Sohn,⁶⁷ A. Sokolov,¹⁹ E. Solovieva,²¹ M. Starič,²² M. Steder,⁶ M. Sumihama,⁸ T. Sumiyoshi,⁶¹ U. Tamponi,^{20,63} K. Tanida,⁵¹ G. Tatishvili,⁴⁵ Y. Teramoto,⁴⁵ K. Trabelsi,¹³ T. Tsuboyama,¹³ M. Uchida,⁶⁰ S. Uehara,¹³ T. Uglav,^{21,35} Y. Uno,¹¹ S. Uno,¹³ S. E. Vahsen,¹² C. Van Hulse,¹ P. Vanhoefer,³² G. Vamer,¹² K. E. Varvell,⁵³ V. Vorobyev,³ M. N. Wagner,⁷ C. H. Wang,⁴⁰ M.-Z. Wang,⁴¹ P. Wang,¹⁷ X. L. Wang,⁶⁴ M. Watanabe,⁴⁴ Y. Watanabe,²³ E. Won,²⁷ B. D. Yabsley,⁵³ J. Yamaoka,¹² Y. Yamashita,⁴³ S. Yashchenko,⁶ Y. Yook,⁶⁷ Y. Yusa,⁴⁴ C. C. Zhang,¹⁷ Z. P. Zhang,⁵⁰ V. Zhilich,³ and A. Zupanc²⁴

(Belle Collaboration)

¹University of the Basque Country UPV/EHU, 48080 Bilbao

²University of Bonn, 53115 Bonn

³Budker Institute of Nuclear Physics SB RAS and Novosibirsk State University, Novosibirsk 630090

⁴Faculty of Mathematics and Physics, Charles University, 121 16 Prague

⁵University of Cincinnati, Cincinnati, Ohio 45221

⁶Deutsches Elektronen-Synchrotron, 22607 Hamburg

⁷Justus-Liebig-Universität Gießen, 35392 Gießen

⁸Gifu University, Gifu 501-1193

⁹II. Physikalisches Institut, Georg-August-Universität Göttingen, 37073 Göttingen

¹⁰Gyeongsang National University, Chinju 660-701

¹¹Hanyang University, Seoul 133-791

¹²University of Hawaii, Honolulu, Hawaii 96822

¹³High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801

¹⁴Lkerbasque, 48011 Bilbao

¹⁵Indian Institute of Technology Guwahati, Assam 781039

¹⁶Indian Institute of Technology Madras, Chennai 600036

¹⁷Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049

¹⁸Institute of High Energy Physics, Vienna 1050

¹⁹Institute for High Energy Physics, Protvino 142281

²⁰INFN-Sezione di Torino, 10125 Torino

²¹Institute for Theoretical and Experimental Physics, Moscow 117218

²²J. Stefan Institute, 1000 Ljubljana

²³Kanagawa University, Yokohama 221-8686

²⁴Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, 76131 Karlsruhe

²⁵Kennesaw State University, Kennesaw, Georgia 30144

²⁶Korea Institute of Science and Technology Information, Daejeon 305-806

²⁷Korea University, Seoul 136-713

²⁸Kyungpook National University, Daegu 702-701

²⁹École Polytechnique Fédérale de Lausanne (EPFL), Lausanne 1015

³⁰Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana

³¹University of Maribor, 2000 Maribor

³²Max-Planck-Institut für Physik, 80805 München

³³School of Physics, University of Melbourne, Victoria 3010

³⁴Moscow Physical Engineering Institute, Moscow 115409

³⁵Moscow Institute of Physics and Technology, Moscow Region 141700

³⁶Graduate School of Science, Nagoya University, Nagoya 464-8602

³⁷Kobayashi-Maskawa Institute, Nagoya University, Nagoya 464-8602

³⁸Nara Women's University, Nara 630-8506

³⁹National Central University, Chung-li 32054

⁴⁰National United University, Miao Li 36003

⁴¹Department of Physics, National Taiwan University, Taipei 10617

⁴²H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342

⁴³Nippon Dental University, Niigata 951-8580

⁴⁴Niigata University, Niigata 950-2181

⁴⁵Osaka City University, Osaka 558-8585

⁴⁶Pacific Northwest National Laboratory, Richland, Washington 99352

⁴⁷Panjab University, Chandigarh 160014

⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260

⁴⁹RIKEN BNL Research Center, Upton, New York 11973

⁵⁰University of Science and Technology of China, Hefei 230026

⁵¹Seoul National University, Seoul 151-742

⁵²Sungkyunkwan University, Suwon 440-746

⁵³School of Physics, University of Sydney, New South Wales 2006

⁵⁴Tata Institute of Fundamental Research, Mumbai 400005

⁵⁵Excellence Cluster Universe, Technische Universität München, 85748 Garching

⁵⁶Toho University, Funabashi 274-8510

⁵⁷Tohoku Gakuin University, Tagajo 985-8537

⁵⁸Tohoku University, Sendai 980-8578

⁵⁹Department of Physics, University of Tokyo, Tokyo 113-0033

⁶⁰Tokyo Institute of Technology, Tokyo 152-8550

⁶¹Tokyo Metropolitan University, Tokyo 192-0397

⁶²Tokyo University of Agriculture and Technology, Tokyo 184-8588

⁶³University of Torino, 10124 Torino

⁶⁴CNP, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

⁶⁵Wayne State University, Detroit, Michigan 48202

⁶⁶Yamagata University, Yamagata 990-8560

⁶⁷Yonsei University, Seoul 120-749

⁶⁸Luther College, Decorah, Iowa 52101

(Received 30 March 2013; published 17 June 2013; publisher error corrected 19 June 2013)

The cross section for $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ between 3.8 and 5.5 GeV is measured with a 967 fb⁻¹ data sample collected by the Belle detector at or near the $Y(nS)$ ($n = 1, 2, \dots, 5$) resonances. The $Y(4260)$ state is observed, and its resonance parameters are determined. In addition, an excess of $\pi^+\pi^- J/\psi$ production around 4 GeV is observed. This feature can be described by a Breit-Wigner parametrization with properties that are consistent with the $Y(4008)$ state that was previously reported by Belle. In a study of $Y(4260) \rightarrow \pi^+\pi^- J/\psi$ decays, a structure is observed in the $M(\pi^+\pi^- J/\psi)$ mass spectrum with 5.2 σ significance, with mass $M = (3894.5 \pm 6.6 \pm 4.5)$ MeV/ c^2 and width $\Gamma = (63 \pm 24 \pm 26)$ MeV/ c^2 , where the errors are statistical and systematic, respectively. This structure can be interpreted as a new charged charmoniumlike state.

DOI: 10.1103/PhysRevLett.110.252002

PACS numbers: 14.40.Rt, 13.25.Gv, 13.66.Bc, 14.40.Pq

$Y(4260)$ state was first observed by the BABAR experiment in the initial-state-radiation (ISR) process $\gamma_{\text{ISR}} \pi^+ \pi^- J/\psi$ [1] and then confirmed by the [2] and Belle experiments [3] using the same

technique. Subsequently, a charged $Z(4430)^\pm$ charmoniumlike state was reported in the $\pi^\pm \psi(2S)$ invariant mass spectrum of $B \rightarrow K \pi^\pm \psi(2S)$ [4] and two Z^\pm states were observed in the $\pi^\pm \chi_{c1}$ invariant mass distribution of


中国的“大型对撞机”大辩论



在规划中，CEPC-SPPC项目将在50-100公里长的地下环形通道内，利用相同的隧道，建造两座超级对撞机：正负电子对撞机和质子对撞机。

该质子对撞机和LHC相似，都在环形通道内使用质子进行对撞。但该对撞机的通道长度将是LHC的2-4倍，对撞能级可达70-100TeV或100-140TeV，远超LHC的14TeV。中科院高能所建议的超大对撞机预算将高于200亿美元（1335亿人民币）。


环形正负电子对撞机 (25TeV) 超级质子对撞机 (50-70TeV)



费米梦想

费米曾在1954年提出环绕地球建一台加速器的设想，称为费米的梦。其能量可达数千TeV。

Fermi's Dream Accelerator (ca. 1954)



Magnetic field	2T
Radius	8000km
Cost	$\$170 \times 10^9$
beam energy	5000TeV
center of mass energy	3TeV
Time to construct	40 years

谢谢！

