



26°C vs 10°C




142 peaks (20.7%) significantly gain H3K27me3 as temperature rises from 10°C to 26°C according to a fold-change of 2 and a q-value of 0.05


H3K27me3 levels in peaks responding to temperature





**H3K27me3 levels in peaks
responding to temperature**



**Expression levels in genes
marked by H3K27me3 peaks
responding to temperature**




Ribosome biogenesis, protein translation





ostta01g04120 a homolog of the yeast MDN gene, which encodes a non-ribosomal protein involved in the maturation and assembly of the 60S ribosomal subunit

Ribosome biogenesis, protein translation





ostta05g04180 RNA helicase family

Ribosome biogenesis, protein translation



ostta10g00010 Encodes a nucleolar protein that is a ribosome biogenesis co-factor.



Ribosome biogenesis, protein translation



AT3G08730 - Encodes a protein-serine kinase that phosphorylates ribosomal protein in vitro. Involved in translational up-regulation of ribosomal proteins.

Cell cycle, Cellular organization



ostta02g01020



ostta02g01020 – AT5G42020 HSP70 Luminal binding protein (BiP2) involved in polar nuclei fusion during proliferation of endosperm nuclei.



<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2824368/>

Cell cycle, Cellular organization





AT3G57060 - Similar to mammalian condensin. Mutants have reduced fertility.

Cell cycle, Cellular organization





AT4G33270 - Encodes a CDC20 protein that interacts with APC subunits

Cell cycle, Cellular organization





AT5G57390 AINTEGUMENTA-LIKE 5 EMBRYOMAKER Encodes a member of the AP2 family of transcriptional regulators. May be involved in germination and seedling growth. Essential for the developmental transition between the embryonic and vegetative phases in plants. Overexpression results in the formation of somatic embryos on cotyledons.

Cell cycle, Cellular organization





AT3G18524 Encodes a DNA mismatch repair homolog of human MutS gene, MSH6. MSH2 is involved in maintaining genome stability and repressing recombination of mismatched heteroduplexes

Cell cycle, Cellular organization





ostta10g02060 AT5G49150 Encodes a transmembrane domain containing protein expressed in sperm cells. Mutants are defective in gamete fusion.

Cell cycle, Cellular organization




AT3G23890 - Encodes a topoisomerase II that is highly expressed in young seedlings. The protein is localized in the nucleus and gene expression levels are increased in proliferative tissues.

Cell cycle, Cellular organization






AT5G40440 - Encodes a mitogen-activated protein kinase kinase. Activates MPK8 and is a target of MPKKK20. Mutant root growth is sensitive oryzalin and suggestive of a role in signaling during microtubule organization.





AT5G19680


PP1 Regulatory Subunit3.








AT2G43770 - Transducin/WD40 repeat-like superfamily protein









ostta07g00900






AT3G06270 Protein phosphatase 2C family protein








AT3G49660 WD40 repeat Encodes a structural core component of a COMPASS-like H3K4 histone methylation complex.





ostta09g01615





AT3G55100 ABC-2 type transporter family protein





AT2G20030 ARABIDOPSIS TOXICOS EN LEVADURA 12, ATL12







AT2G21860 - Violaxanthin de-epoxidase located in chloroplast.





AT1G21840 - Encodes a urease accessory protein which is essential for the activation of plant urease.












AT4G21090 MITOCHONDRIAL FERREDOXIN 2




ostta17g00300




AT3G45770 Encodes a mitochondrial enoyl-ACP reductase, a component of the fatty acid synthase complex.




ostta18g00970






ostta20g00030





AT2G30200 - Malonyl-ACP expressed in developing seeds

ostta01g02760



ostta01g02760 Carbohydrate binding module family 20

osta02g00050



osta02g00050 Ribonuclease H-like domain