

Project1__

September 30, 2023

1 Project #1 - PHYS 7127

1.1 Part A: Due Friday, September 29, 11:59pm

1.2 A. (40 points; due Sept 29) Generating CMB power spectra and matter power spectra

1.2.1 (a) What is the wavenumber of the first acoustic peak? How does this compare with the angular sizes that was calculated in Question 2(c)?

```
[ ]: import camb
import numpy as np
import matplotlib.pyplot as plt
```

```
[ ]: H0=67.66; ombh2=0.02242; omch2=0.11933
h = H0/100
omb = ombh2/h**2
omc = omch2/h**2
omm = omb+omc
```

```
[ ]: pars = camb.read_ini("./inifiles/params.ini")
pars.set_cosmology(H0=H0, ombh2=ombh2, omch2=omch2)
```

```
[ ]: class: <CAMBparams>
      WantCls = True
      WantTransfer = True
      WantScalars = True
      WantTensors = False
      WantVectors = False
      WantDerivedParameters = True
      Want_cl_2D_array = True
      Want_CMB = True
      Want_CMB_lensing = True
      DoLensing = True
      NonLinear = NonLinear_none
      Transfer: <TransferParams>
        high_precision = False
        accurate_massive_neutrinos = False
        kmax = 1.4
```

```

k_per_logint = 0
PK_num_redshifts = 1
PK_redshifts = [0.0]
want_zstar = False
want_zdrag = False
min_l = 2
max_l = 2200
max_l_tensor = 600
max_eta_k = 4400.0
max_eta_k_tensor = 1200.0
ombh2 = 0.02242
omch2 = 0.11933
omk = 0.0
omnuh2 = 0.000644866570625114
H0 = 67.66
TCMB = 2.7255
YHe = 0.24587684703799
num_nu_massless = 2.0293333333333337
num_nu_massive = 1
nu_mass_eigenstates = 1
share_delta_neff = True
nu_mass_degeneracies = [1.0146666666666666]
nu_mass_fractions = [1.0]
nu_mass_numbers = [1]
InitPower: <InitialPowerLaw>
    tensor_parameterization = tensor_param_indeptilt
    ns = 0.96
    nrun = 0.0
    nrunrun = 0.0
    nt = 0.0
    ntrun = 0.0
    r = 0.0
    pivot_scalar = 0.05
    pivot_tensor = 0.05
    As = 2.1e-09
    At = 0.0
Recomb: <Recfast>
    min_a_evolve_Tm = 0.0011098779505118728
    RECFAST_fudge = 1.125
    RECFAST_fudge_He = 0.86
    RECFAST_Heswitch = 6
    RECFAST_Hswitch = True
    AGauss1 = -0.14
    AGauss2 = 0.079
    zGauss1 = 7.28
    zGauss2 = 6.73
    wGauss1 = 0.18

```

```

wGauss2 = 0.33
Reion: <TanhReionization>
  Reionization = True
  use_optical_depth = True
  redshift = 10.0
  optical_depth = 0.09
  delta_redshift = 1.5
  fraction = -1.0
  include_helium_fullreion = True
  helium_redshift = 3.5
  helium_delta_redshift = 0.4
  helium_redshiftstart = 5.5
  tau_solve_accuracy_boost = 1.0
  timestep_boost = 1.0
  max_redshift = 50.0
DarkEnergy: <DarkEnergyFluid>
  w = -1.0
  wa = 0.0
  cs2 = 1.0
  use_tabulated_w = False
NonLinearModel: <Halofit>
  Min_kh_nonlinear = 0.005
  halofit_version = mead2020
  HMCode_A_baryon = 3.13
  HMCode_eta_baryon = 0.603
  HMCode_logT_AGN = 7.8
Accuracy: <AccuracyParams>
  AccuracyBoost = 1.0
  lSampleBoost = 1.0
  lAccuracyBoost = 1.0
  AccuratePolarization = True
  AccurateBB = False
  AccurateReionization = True
  TimeStepBoost = 1.0
  BackgroundTimeStepBoost = 1.0
  IntTolBoost = 1.0
  SourcekAccuracyBoost = 1.0
  IntkAccuracyBoost = 1.0
  TransferkBoost = 1.0
  NonFlatIntAccuracyBoost = 1.0
  BessIntBoost = 1.0
  LensingBoost = 1.0
  NonlinSourceBoost = 1.0
  BesselBoost = 1.0
  LimberBoost = 1.0
  SourceLimmerBoost = 1.0
  KmaxBoost = 1.0

```

```

    neutrino_q_boost = 1.0
SourceTerms: <SourceTermParams>
    limber_windows = True
    limber_phi_lmin = 100
    counts_density = True
    counts_redshift = True
    counts_lensing = False
    counts_velocity = True
    counts_radial = False
    counts_timedelay = True
    counts_ISW = True
    counts_potential = True
    counts_evolve = False
    line_phot_dipole = False
    line_phot_quadrupole = False
    line_basic = True
    line_distortions = True
    line_extra = False
    line_reionization = False
    use_21cm_mK = True
z_outputs = []
scalar_initial_condition = initial_adiabatic
InitialConditionVector = []
OutputNormalization = 1
Alens = 1.0
MassiveNuMethod = Nu_trunc
DoLateRadTruncation = True
Evolve_baryon_cs = False
Evolve_delta_xe = False
Evolve_delta_Ts = False
Do21cm = False
transfer_21cm_cl = False
Log_lvalues = False
use_cl_spline_template = True
min_l_logl_sampling = 5000
SourceWindows = []
CustomSources: <CustomSources>
    num_custom_sources = 0
    c_source_func = None
    custom_source_ell_scales = []

```


```

[ ]: results = camb.get_results(pars)
powers = results.get_cmb_power_spectra(pars, CMB_unit='muK')
totalCL = powers['total']
C_TT = totalCL[:,0]
ls = np.arange(totalCL.shape[0])

```

```
kh, z, pk = results.get_matter_power_spectrum()
```

```
[ ]: plt.plot(ls, C_TT)
plt.xlim(xmin=2)
plt.xlabel("$\ell$")
plt.ylabel("$C_{TT} \text{ } [\mu K^2]$")
plt.title("temperature angular power spectrum vs. wavenumber")
plt.show()
```



```
[ ]: l_1max = np.argmax(C_TT)
theta = np.pi/l_1max
```

```
[ ]: print(f"The wavenumber l of the first acoustic peak is {l_1max}.")
print(f"The corresponding angular size is {np.rad2deg(np.pi/l_1max):.2f}°")
```

The wavenumber l of the first acoustic peak is 220.

The corresponding angular size is 0.82 degree.

Since $H_0 = 67.66$, and $\Omega_m \approx 1$ at the last scattering surface, from ProblemSet1 2(c), it can be inferred that the corresponding angular radius is around 0.8 degree. Therefore, the angular size

corresponding to wavenumber 220 is pretty close to the angular sizes that was calculated in Question 2(c) of ProblemSet1.

1.2.2 (b) Use CAMB to compute the CMB power spectra for the following cosmological models. Plot the C_{TT} power spectra on a single plot. For each model, describe the qualitative differences between it and the Planck 2018 model and the physical reasons behind the differences.

```
[ ]: low_baryon_fraction = camb.read_ini("./inifiles/params.ini")
low_omb = 0.02
low_baryon_fraction.set_cosmology(H0=H0, ombh2=low_omb*h**2,
    ↪omch2=(omm-low_omb)*h**2)
low_results = camb.get_results(low_baryon_fraction)
low_powers = low_results.get_cmb_power_spectra(low_baryon_fraction,
    ↪CMB_unit='muK')
low_totalCL = low_powers['total']
low_C_TT = low_totalCL[:,0]
low_ls = np.arange(low_C_TT.shape[0])
low_kh, low_z, low_pk = low_results.get_matter_power_spectrum()
```

```
[ ]: high_baryon_fraction = camb.read_ini("./inifiles/params.ini")
high_omb = 0.1
high_baryon_fraction.set_cosmology(H0=H0, ombh2=high_omb*h**2,
    ↪omch2=(omm-high_omb)*h**2)
high_results = camb.get_results(high_baryon_fraction)
high_powers = high_results.get_cmb_power_spectra(high_baryon_fraction,
    ↪CMB_unit='muK')
high_totalCL = high_powers['total']
high_C_TT = high_totalCL[:,0]
high_ls = np.arange(high_C_TT.shape[0])
high_kh, high_z, high_pk = high_results.get_matter_power_spectrum()
```

```
[ ]: open_universe_no_dark_energy = camb.read_ini("./inifiles/params.ini")
open_omlambda = 0
open_universe_no_dark_energy.set_cosmology(H0=H0, ombh2=ombh2, omch2=omch2,
    ↪omk=(1-omb-omc-open_omlambda))
open_results = camb.get_results(open_universe_no_dark_energy)
open_powers = open_results.get_cmb_power_spectra(open_universe_no_dark_energy,
    ↪CMB_unit='muK')
open_totalCL = open_powers['total']
open_C_TT = open_totalCL[:,0]
open_ls = np.arange(open_C_TT.shape[0])
open_kh, open_z, open_pk = open_results.get_matter_power_spectrum()
```

```
[ ]: small_matter_fraction = camb.read_ini("./inifiles/params.ini")
small_omc = 0.1
small_omk = 0
```

```

small_matter_fraction.set_cosmology(H0=H0, ombh2=ombh2, omch2=small_omc*h**2,
    ↪omk=small_omk)
small_results = camb.get_results(small_matter_fraction)
small_powers = small_results.get_cmb_power_spectra(small_matter_fraction,
    ↪CMB_unit='muK')
small_totalCL = small_powers['total']
small_C_TT = small_totalCL[:,0]
small_ls = np.arange(small_C_TT.shape[0])
small_kh, small_z, small_pk = small_results.get_matter_power_spectrum()

```

```

[ ]: very_early_reionization = camb.read_ini("./inifiles/params.ini")
early_z = 30
very_early_reionization.set_cosmology(H0=H0, ombh2=ombh2, omch2=ombh2,
    ↪zrei=early_z, tau=None)
early_results = camb.get_results(very_early_reionization)
early_powers = early_results.get_cmb_power_spectra(very_early_reionization,
    ↪CMB_unit='muK')
early_totalCL = early_powers['total']
early_C_TT = early_totalCL[:,0]
early_ls = np.arange(early_C_TT.shape[0])
early_kh, early_z, early_pk = early_results.get_matter_power_spectrum()


```

WARNING: You seem to have set the optical depth, but use_optical_depth = F
 WARNING: You seem to have set the optical depth, but use_optical_depth = F

```

[ ]: plt.plot(ls, C_TT, label="Planck 2018 model", c='k')
plt.plot(low_ls, low_C_TT, label="Low baryon fraction")
plt.plot(high_ls, high_C_TT, label="High baryon fraction")
plt.plot(open_ls, open_C_TT, label='Open universe, no dark energy')
plt.plot(small_ls, small_C_TT, label='Small matter fraction, flat universe')
plt.plot(early_ls, early_C_TT, label="Very early reionization")
plt.legend()
plt.xlim(xmin=2, xmax=2000)
plt.xlabel("$\ell$")
plt.ylabel("$C_{TT} \ [\mu K^2]$")
plt.title("temperature angular power spectrum vs. wavenumber")
plt.show()

```



Low baryon fraction:

In the low baryon fraction universe, dark matter becomes more dominant in the early universe's dynamics. Dark matter does not experience the same oscillations as baryonic matter, and its presence suppresses the amplitude of the acoustic peaks.

Changing the baryon fraction also affects the sound horizon at the time of recombination. The sound horizon determines the scale of the acoustic oscillations, which in turn affects the position (wavenumber) of the peaks. When you lower the baryon fraction, you effectively decrease the sound speed of the fluid and the distance it can travel before recombination. This leads to a slight shift in the wavenumber of the first peak to smaller values.

High baryon fraction:

As opposite to the low baryon fraction case, high baryon fraction will increase the sound speed and lead to a smaller wavenumber. Besides, since there are more baryons to oscillate, high baryon fraction increase the amplitude of the acoustic peaks.

Open universe, no dark energy:

The absence of dark energy in an open universe can lead to differences in the damping of acoustic oscillations, causing a suppression of the amplitude of the CMB peaks at small angular scales (1). This is because the absence of dark energy can result in a faster expansion rate and a different angular diameter distance to the last scattering surface.

The positions (wavenumbers) of the CMB peaks can also be shifted due to differences in the angular diameter distance and the evolution of perturbations. In an open universe, the positions of the peaks may be shifted to larger l values compared to a flat universe with dark energy.

Small matter fraction, flat universe:

With lower dark matter density, the baryonic matter has a more dominant role in shaping the CMB power spectrum. Baryonic matter has a more pronounced effect on the acoustic oscillations, and in this scenario, the peaks could be enhanced due to the increased influence of baryons.

The shift in the wavenumber of the first peak likely reflects differences in the sound horizon and the dynamics of acoustic oscillations. In the universe with lower dark matter density, the altered balance between dark matter and baryons could result in a different sound horizon and lead to a slightly shifted first peak.

Very early reionization:

Early reionization means that the universe became ionized at a higher redshift (z), indicating that the intergalactic medium was significantly heated and ionized much earlier in cosmic history. This process suppresses the growth of small-scale density perturbations. As a result, the first acoustic peak and small-scale fluctuations, such as those responsible for high- l CMB anisotropies, are damped or reduced.

The positions (wavenumbers) of the CMB peaks can also be affected by the damping of fluctuations. The damping process can lead to a shift in the peak positions to larger l values.


1.2.3 (c) Go crazy and create an alternative universe, e.g. massive neutrinos, no ionization, nearly dominated by dark energy, or a closed universe. If CAMB crashes, don't panic and determine which parameter(s) are too extreme and retry the experiment. Compare your alternative universe with the standard cosmological model (part a) by plotting their TT power spectra in one plot.

```
[ ]: no_ionization_universe = camb.read_ini("./inifiles/params.ini")
no_ioni_z = 0
no_ionization_universe.set_cosmology(H0=H0, ombh2=ombh2, omch2=ombh2,
    ↪zrei=no_ioni_z, tau=None, )
no_ioni_results = camb.get_results(no_ionization_universe)
no_ioni_powers = no_ioni_results.get_cmb_power_spectra(no_ionization_universe,
    ↪CMB_unit='muK')
no_ioni_totalCL = no_ioni_powers['total']
no_ioni_C_TT = no_ioni_totalCL[:,0]
no_ioni_ls = np.arange(no_ioni_C_TT.shape[0])
no_ioni_kh, no_ioni_z, no_ioni_pk = no_ioni_results.get_matter_power_spectrum()
```

```
WARNING: You seem to have set the optical depth, but use_optical_depth = F
WARNING: You seem to have set the optical depth, but use_optical_depth = F
```

```
[ ]: plt.plot(ls, C_TT, label="Planck 2018 model", c='k')
plt.plot(no_ioni_ls, no_ioni_C_TT, label="no ionization universe")
plt.legend()
```

```
plt.xlim(xmin=2, xmax=2000)
plt.xlabel("$\ell$")
plt.ylabel("$C_{\ell} [\mu K^2]$")
plt.title("temperature angular power spectrum vs. wavenumber")
plt.show()
```




Without ionization, the baryonic acoustic oscillations (BAO) in the early universe, responsible for the acoustic peaks, will be less damped. This results in sharper and more pronounced peaks in the CMB power spectrum. The first acoustic peak, in particular, will be well-defined and have a higher contrast.

The position (wavenumber) of the first acoustic peak is closely related to the sound horizon at the time of recombination. In a universe with no ionization, the altered expansion history can lead to a different sound horizon. A larger sound horizon would result in a higher peak position in the CMB power spectrum.

1.2.4 (d) CAMB also calculates the matter power spectrum (if get transfer = T), among many other quantities. As discussed in class, this describes how clustered matter is at recombination. In one plot, show the matter spectra that were calculated for each of the cosmological models in parts (b) and (c). You will need these results for the next part.

```
[ ]: plt.loglog(kh, pk[0,:], label="Planck 2018 model")
plt.loglog(low_kh, low_pk[0,:], label="Low baryon fraction")
plt.loglog(high_kh, high_pk[0,:], label="High baryon fraction")
plt.loglog(open_kh, open_pk[0,:], label="Open universe, no dark energy")
plt.loglog(small_kh, small_pk[0,:], label="Small matter fraction, flat_
    universe")
plt.loglog(early_kh, early_pk[0,:], label="Very early reionization")
plt.loglog(no_ioni_kh, no_ioni_pk[0,:], label="no ionization universe")
plt.legend(fontsize="9")
plt.xlabel("k/h [Mpc]")
plt.ylabel("Matter power")
plt.ylim(ymin=1)
plt.show()
```



```
[ ]:
```