2 A student observes the orbits of some of the moons around the planet Saturn, as shown in Fig. 2.1.




Fig. 2.1 (not to scale)

For the moon Pandora, the period of the orbit and the mean distance from the centre of Saturn are determined.

The measurements of period T and mean distance r are repeated for other moons.

It is suggested that T and r are related by the equation

$$T = \frac{2\pi r^n}{k}$$

where n and k are constants.

(a) A graph is plotted of lg T on the y-axis against lg r on the x-axis.

Determine expressions for the gradient and y-intercept.

gradient = ...

y-intercept = ...

[1]

DO NOT WRITE IN THIS MARGIN

(b) Values of r and T are given for different moons in Table 2.1.

Table 2.1

moon	$r/10^8\,\text{m}$	$T/10^3\,\text{s}$	$\lg(r/10^8\,\text{m})$	$\lg(T/10^3\,\text{s})$
Pandora	1.42	52 ± 5		
Mimas	1.86	81 ± 5		
Enceladus	2.38	120 ± 10		
Tethys	2.95	170 ± 10		
Dione	3.77	240 ± 20		
Rhea	5.28	390 ± 30		

Calculate and record values of $\lg(r/10^8\,\text{m})$ and $\lg(T/10^3\,\text{s})$ in Table 2.1. Include the absolute uncertainties in $\lg(T/10^3\,\text{s})$. [2]


(c) **(i)** Plot a graph of $\lg(T/10^3\,\text{s})$ against $\lg(r/10^8\,\text{m})$. Include error bars for $\lg(T/10^3\,\text{s})$. [2]

(ii) Draw the straight line of best fit and a worst acceptable straight line on your graph. Label both lines. [2]

(iii) Determine the gradient of the line of best fit. Include the absolute uncertainty in your answer.

gradient = ... [2]

DO NOT WRITE IN THIS MARGIN



DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(iv) Determine the *y*-intercept of the line of best fit. Include the absolute uncertainty in your answer.

y-intercept = ... [2]

(d) Using your answers to **(a)**, **(c)(iii)** and **(c)(iv)**, determine the values of *n* and *k*. Include the absolute uncertainties in *n* and *k*. You need not be concerned with units.

n = ...

k = ...
[3]

(e) Titan is another moon of Saturn. The orbit of Titan has a period of 1.38×10^6 s.

Determine the value of *r* for Titan.

r = ... m [1]

[Total: 15]

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN