3 **(a)** State the reason why two objects that are at the same temperature are described as being in thermal equilibrium.

.....

..... [1]

(b) Fig. 3.1 shows the variations with temperature of the densities of mercury and of water between 0 °C and 100 °C.




Fig. 3.1

Temperature may be measured using the variation with temperature of the density of a liquid.

Suggest why, for measuring temperature over this temperature range:

(i) mercury is a suitable liquid

.....

..... [1]

(ii) water is not a suitable liquid.

.....

.....

..... [2]

(c) A beaker contains a liquid of mass 120 g. The liquid is supplied with thermal energy at a rate of 810 W. The beaker has a mass of 42 g and a specific heat capacity of $0.84\,J\,g^{-1}\,K^{-1}$. The beaker and the liquid are in thermal equilibrium with each other at all times and are insulated from the surroundings.

Fig. 3.2 shows the variation with time *t* of the temperature of the liquid.

© UCLES 2023




Fig. 3.2

(i) State the boiling temperature, in °C, of the liquid.

temperature = ... °C [1]

(ii) Determine the specific heat capacity, in $J\,g^{-1}\,K^{-1}$, of the liquid.

specific heat capacity = ... $J\,g^{-1}\,K^{-1}$ [4]

(d) The experiment in **(c)** is repeated using water instead of the liquid in **(c)**. The mass of liquid used, the power supplied, and the initial temperature are all unchanged.
The specific heat capacity of water is approximately twice that of the liquid in **(c)**.
The boiling temperature of water is 100 °C.

On Fig. 3.2, sketch the variation with time t of the temperature of the water between $t = 0$ and $t = 60$ s. Numerical calculations are not required. [2]

[Total: 11]