

Inclusion-Exclusion Principle: Proof by Mathematical Induction For Dummies

Vita Smid*

December 2, 2009

Definition (Discrete Interval). $[n] := \{1, 2, 3, \dots, n\}$

Theorem (Inclusion-Exclusion Principle). *Let A_1, A_2, \dots, A_n be finite sets. Then*

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{\substack{J \subseteq [n] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right|$$

Proof (induction on n). The theorem holds for $n = 1$:

$$\left| \bigcup_{i=1}^1 A_i \right| = |A_1| \tag{1}$$

$$\sum_{\substack{J \subseteq [1] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| = (-1)^0 \left| \bigcap_{i \in \{1\}} A_i \right| = |A_1| \tag{2}$$

For the induction step, let us suppose the theorem holds for $n - 1$.

$$\left| \bigcup_{i=1}^n A_i \right| = \left| \left(\bigcup_{i=1}^{n-1} A_i \right) \cup A_n \right| = \tag{3}$$

We can use the formula $|X \cup Y| = |X| + |Y| - |X \cap Y|$.

$$= \left| \bigcup_{i=1}^{n-1} A_i \right| + |A_n| - \left| \left(\bigcup_{i=1}^{n-1} A_i \right) \cap A_n \right| = \tag{4}$$

Intersection of unions can be rewritten as a union of intersections.

$$= \left| \bigcup_{i=1}^{n-1} A_i \right| + |A_n| - \left| \bigcup_{i=1}^{n-1} (A_i \cap A_n) \right| = \tag{5}$$

Let us define the substitution $B_i := A_i \cap A_n$.

$$= \left| \bigcup_{i=1}^{n-1} A_i \right| + |A_n| - \left| \bigcup_{i=1}^{n-1} B_i \right| = \tag{6}$$

We can now use the induction hypothesis on both $\bigcup A_i$ and $\bigcup B_i$.

$$= \sum_{\substack{J \subseteq [n-1] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + |A_n| - \sum_{\substack{K \subseteq [n-1] \\ K \neq \emptyset}} (-1)^{|K|-1} \left| \bigcap_{i \in K} B_i \right| = \tag{7}$$

*<http://ze.phyr.us>

The expression $(-1) \sum (-1)^{|K|-1} |\bigcap B_i|$ is equivalent to $\sum (-1)(-1)^{|K|-1} |\bigcap B_i| = \sum (-1)^{|K|} |\bigcap B_i|$.

$$= \sum_{\substack{J \subseteq [n-1] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + |A_n| + \sum_{\substack{K \subseteq [n-1] \\ K \neq \emptyset}} (-1)^{|K|} \left| \bigcap_{i \in K} B_i \right| \quad (8)$$

Since $K \neq \emptyset$, we can revert the substitution:

$$\bigcap_{i \in K} B_i = \bigcap_{i \in K} (A_i \cap A_n) = \left(\bigcap_{i \in K} A_i \right) \cap A_n = \bigcap_{i \in K \cup \{n\}} A_i$$

(8) now becomes

$$\sum_{\substack{J \subseteq [n-1] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + |A_n| + \sum_{\substack{K \subseteq [n-1] \\ K \neq \emptyset}} (-1)^{|K|} \left| \bigcap_{i \in K \cup \{n\}} A_i \right| \quad (9)$$

Let us substitute $K \cup \{n\}$ with J . The expression $|K|$ thus becomes $|J| - 1$ (with K defined as a subset of $[n-1]$, K cannot contain n and thus $|K \cup \{n\}| = |K| + 1$).

To replace the expression $i \in K \cup \{n\}$ with $i \in J$, we must impose several conditions on J as the summation index:

- $J \neq \emptyset$ (the same condition that was imposed on K),
- $J \subseteq [n] \wedge n \in J$ (n must be contained in every J , since J replaces $K \cup \{n\}$),
- $J \neq \{n\}$ ($K \cup \{n\} \neq \{n\}$, since $K \neq \emptyset$ and $n \notin K \subseteq [n-1]$).

(9) now becomes

$$\sum_{\substack{J \subseteq [n-1] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + |A_n| + \sum_{\substack{J \subseteq [n] \\ n \in J \\ J \neq \emptyset, \{n\}}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| \quad (10)$$

The Inclusion-Exclusion Principle can be used on A_n alone (we have already shown that the theorem holds for one set):

$$\sum_{\substack{J \subseteq \{n\} \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| = (-1)^{|\{n\}|-1} \left| \bigcap_{i \in \{n\}} A_i \right| = |A_n|$$

(10) now becomes

$$\sum_{\substack{J \subseteq [n-1] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + \sum_{\substack{J \subseteq \{n\} \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + \sum_{\substack{J \subseteq [n] \\ n \in J \\ J \neq \emptyset, \{n\}}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| = \quad (11)$$

For better readability, let us define

$$\begin{aligned} P_1 &:= \mathcal{P}([n-1]) \setminus \{\emptyset\} \\ P_2 &:= \mathcal{P}(\{n\}) \setminus \{\emptyset\} = \{\{n\}\} \\ P_3 &:= \mathcal{P}([n]) \setminus \mathcal{P}([n-1]) \setminus \{\emptyset, \{n\}\} \end{aligned}$$

and rewrite (11) in this way:

$$= \sum_{J \in P_1} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + \sum_{J \in P_2} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| + \sum_{J \in P_3} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| = \quad (12)$$

Since the three sums consist of the same terms, we can combine them into one. As the sets P_1, P_2, P_3 are disjoint, the summation condition now becomes

$$\begin{aligned} J \in (P_1 \cup P_2 \cup P_3) &= (\mathcal{P}([n-1]) \setminus \{\emptyset\}) \cup \{\{n\}\} \cup (\mathcal{P}([n]) \setminus \mathcal{P}([n-1]) \setminus \{\emptyset, \{n\}\}) = \\ &= (\mathcal{P}([n-1]) \cup \{\{n\}\}) \cup (\mathcal{P}([n]) \setminus \mathcal{P}([n-1]) \setminus \{\{n\}\}) \setminus \{\emptyset\} = \mathcal{P}([n]) \setminus \{\emptyset\} \end{aligned}$$

Finally, we can replace the logical condition $J \in \mathcal{P}([n]) \setminus \{\emptyset\}$ by the equivalent statement $J \subseteq [n], J \neq \emptyset$. The resulting formula is an instance of the Inclusion-Exclusion Theorem for n sets:

$$= \sum_{\substack{J \subseteq [n] \\ J \neq \emptyset}} (-1)^{|J|-1} \left| \bigcap_{i \in J} A_i \right| \quad (13)$$

□

Remark. It can be easily seen that every possible value of J is covered *exactly once* by the new summation condition ($J \subseteq [n], J \neq \emptyset$):

$$\forall J \subseteq [n], J \neq \emptyset \begin{cases} n \notin J & (\Leftrightarrow J \subseteq [n-1]) \\ n \in J & \begin{cases} J = \{n\} & (\Leftrightarrow J \subseteq \{n\}) \\ J \neq \{n\} & (\Leftrightarrow J \subseteq [n], n \in J, J \neq \{n\}) \end{cases} \end{cases}$$