

The cut property of Minimum Spanning Trees

Vassos Hadzilacos

In this handout we use “graph” to mean “undirected, connected graph”, and “tree” to mean “free tree” (i.e., an undirected, connected, acyclic graph).

You already know that a tree with n nodes has $n - 1$ edges. Using this, you can easily prove the following fact: Adding an edge to a tree creates a unique cycle; removing any edge from that cycle yields a tree again. More precisely:

Fact 1 *Let $T = (V, E)$ be a tree and e be an edge not in E . Then the graph $T^+ = (V, E \cup \{e\})$ has a unique cycle. Furthermore, if e' is any edge on that cycle, the graph $T' = (V, (E \cup \{e\}) - e')$ is a tree.*

A **cut** of a graph $G = (V, E)$ is a partition of the set of nodes V into two sets of nodes, $S \subseteq V$ and $V - S$; we denote this cut as the pair $(S, V - S)$. An edge e of G **crosses** the cut $(S, V - S)$ if one of the endpoints of e is in S and the other is in $V - S$.

Theorem 2 (Cut property of MSTs) *Let $G = (V, E)$ be a graph, $(S, V - S)$ be a cut of G , F be a set of edges of G , and e be an edge of G such that*

- (a) *F is contained in a MST of G ,*
- (b) *no edge in F crosses the cut $(S, V - S)$, and*
- (c) *e is a minimum weight edge that crosses the cut $(S, V - S)$.*

Then $F \cup \{e\}$ is also contained in a MST of G .

Intuitively, this theorem says that we can extend a partial MST (i.e., a subset of the edges of an MST of a graph) to a better partial MST (i.e., an even bigger subset of the edges of an MST of the graph) as follows: Find **any** cut that no edge in F crosses, and pick a minimum weight edge e that crosses the cut; then $F \cup \{e\}$ is the expanded partial MST. It is clear how such a property leads to greedy MST algorithms: We start with the trivial partial solution (an empty set of edges), and then we use the cut property to greedily expand this partial solution, until our “partial” solution contains $n - 1$ edges — i.e., is a full MST. The correctness of Prim’s and Kruskal’s MST algorithms is based on the cut property. (Seeing how is left as an exercise.) We now prove Theorem 2.

PROOF. Let T be a MST that contains the edges in F . (Such a MST exists by hypothesis (a) of the theorem.) If T contains e , we are done. So, suppose T does not contain e .

Let T^+ be the graph that results when we add e to T . By Fact 1, T^+ has a unique cycle; let u and v be the endpoints of e . By hypothesis (c), e crosses the cut $(S, V - S)$, so without loss of generality, suppose that $u \in S$ and $v \in V - S$. The unique cycle of T^+ consists of e , and a path of edges in T that connects u to v . This path contains an edge, say e' , that has one endpoint in S and the other in $V - S$; this is because $u \in S$ and $v \in V - S$. So, e' also crosses the cut $(S, V - S)$.

Since both e and e' cross the cut $(S, V - S)$, and e is a minimum weight edge that crosses this cut (see hypothesis (c)), $\text{weight}(e) \leq \text{weight}(e')$. By Fact 1, removing e' from T^+ results in a spanning tree T' of G . Recall that T' is constructed by adding e to T and then removing e' from the resulting graph. So,

$$\text{weight}(T') = \text{weight}(T) + \text{weight}(e) - \text{weight}(e') \leq \text{weight}(T)$$

(where the last inequality follows from the fact that $\mathbf{weight}(e) \leq \mathbf{weight}(e')$). Since (i) T' is a spanning tree of G , (ii) T is a MST of G , and (iii) $\mathbf{weight}(T') \leq \mathbf{weight}(T)$, it follows that

$$T' \text{ is also a MST of } G. \tag{1}$$

Next we argue that T' contains all the edges in F . To see this recall that T contains all the edges in F , and e' , the only edge of T that T' does not contain, is not in F . (This is because e' crosses $(S, V - S)$ and no edge in F crosses $(S, V - S)$, by hypothesis (b).) So, the edges of T excluding e' contain the edges in F . T' contains all the edges of T except e' , and so

$$T' \text{ contains the edges in } F. \tag{2}$$

By its definition,

$$T' \text{ contains the edge } e. \tag{3}$$

By (1), (2), and (3), T' is a MST of G that contains the edges in $F \cup \{e\}$, as wanted. \square