
LABEL: PYRUKYND- mitapivat tablet, film coated PYRUKYND- mitapivat kit

VIEW PACKAGE PHOTOS





VIEW MORE

NDC Code(s): 71334-205-05, 71334-205-07, 71334-205-14, 71334-210-07, view more
Packager: Agios Pharmaceuticals, Inc.

Category: HUMAN PRESCRIPTION DRUG LABEL
DEA Schedule: None
Marketing Status: New Drug Application

DRUG LABEL INFORMATION

Updated June 30, 2026

If you are a consumer or patient please visit this version.

VIEW ALL SECTIONS

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PYRUKYND safely and effectively. See full prescribing information for PYRUKYND. PYRUKYND - ®(mitapivat) tablets, for oral ...

TABLE OF CONTENTS

Table of Contents

1 INDICATIONS AND USAGE

PYRUKYND is indicated for the treatment of hemolytic anemia in adults with pyruvate kinase (PK) deficiency.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose - PYRUKYND is taken with or without food and swallowed whole. Do not split, crush, chew, or dissolve the tablets. The starting dosage for PYRUKYND is 5 mg orally twice daily ...

3 DOSAGE FORMS AND STRENGTHS

5 mg tablets: round, blue, film-coated tablets with "M5" printed on one side. 20 mg tablets: round, blue, film-coated tablets with "M20" printed on one side. 50 mg tablets: oblong, blue ...

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Acute Hemolysis with Abrupt Treatment Interruption - Acute hemolysis with subsequent anemia has been observed following abrupt interruption or discontinuation of PYRUKYND in a dose-ranging ...

6 ADVERSE REACTIONS

The following clinically significant adverse reaction is described elsewhere in labeling: Acute Hemolysis with Abrupt Treatment Discontinuation - [see - Warnings and Precautions ...

7 DRUG INTERACTIONS

7.1 Effect of Other Drugs on PYRUKYND - Strong CYP3A Inhibitors - Clinical Impact - Co-administration of PYRUKYND with strong CYP3A inhibitors increased mitapivat plasma concentrations ...

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy - Risk Summary - Available data from clinical trials of PYRUKYND are insufficient to evaluate for a drug- associated risk of major birth defects, miscarriage or other adverse ...

11 DESCRIPTION

The active ingredient of PYRUKYND is mitapivat, a pyruvate kinase activator, present as mitapivat sulfate. The chemical name of mitapivat sulfate is 8-quinolinesulfonamide, N-[4-[[4 ...

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action - Mitapivat is a pyruvate kinase activator that acts by allosterically binding to the pyruvate kinase tetramer and increasing pyruvate kinase (PK) activity. The red blood ...

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility - Carcinogenesis - Mitapivat was not carcinogenic in transgenic rasH2 mice up to the highest doses tested at 500 mg/kg/day in males and ...

14 CLINICAL STUDIES

Patients with PK Deficiency - Patients Not Regularly Transfused - The efficacy of PYRUKYND was evaluated in ACTIVATE, a multinational, randomized, double-blind, placebo-controlled clinical study ...

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied - PYRUKYND 28-Day Packs - Tablet Strength - Description - Imprint - NDC - 5 mg Round, blue, film-coated tablets "M5" printed on one ...

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information). Acute Hemolysis with Abrupt Treatment Interruption - Inform patients of the risk of developing acute hemolysis ...

PATIENT PACKAGE INSERT

PATIENT INFORMATION - PYRUKYND [pye roo' kind] (mitapivat) tablets, for oral use - What is PYRUKYND? PYRUKYND is a prescription medicine used to treat low ...

PRINCIPAL DISPLAY PANEL - 5 MG TABLET BLISTER PACK CARTON

NDC 71334-205-05 - Do not use if seal - is broken or damaged - Rx Only - pyrukynd ® (mitapivat) tablets - 5 mg per tablet - 56 tablets - Contains 4-week supply of PYRUKYND ...

PRINCIPAL DISPLAY PANEL - 20 MG TABLET BLISTER PACK CARTON

NDC 71334-210-20 - Do not use if seal - is broken or damaged - Rx Only - pyrukynd ® (mitapivat) tablets - 20 mg per tablet - 56 tablets - Contains 4-week supply of PYRUKYND ...

PRINCIPAL DISPLAY PANEL - 50 MG TABLET BLISTER PACK CARTON

NDC 71334-215-50 - Do not use if seal - is broken or damaged - Rx Only - pyrukynd ® (mitapivat) tablets - 50 mg per tablet - 56 tablets - Contains 4-week supply of PYRUKYND ...

PRINCIPAL DISPLAY PANEL - 5 MG TABLET BLISTER PACK CARTON - 220-11

NDC 71334-220-11 - Do not use if seal is broken or damaged - Rx Only - pyrukynd ® (mitapivat) tablets - Week 1 - 5 mg per tablet - 5 mg TAPER PACK - 7 tablets - Swallow tablets whole ...

PRINCIPAL DISPLAY PANEL - 20 MG/5 MG TABLET BLISTER PACK CARTON - 225-12

NDC 71334-225-12 - Do not use if seal is broken or damaged - Rx Only - pyrukynd ® (mitapivat) tablets - Week 1 - 20 mg per tablet - Week 2 - 5 mg per tablet - 20 mg and 5 mg TAPER ...

PRINCIPAL DISPLAY PANEL - 50 MG/20 MG TABLET BLISTER PACK CARTON - 230-13

NDC 71334-230-13 - Do not use if seal is broken or damaged - Rx Only - pyrukynd ® (mitapivat) tablets - Week 1 - 50 mg per tablet - Week 2 - 20 mg per tablet - 50 mg and 20 mg ...

INGREDIENTS AND APPEARANCE

Product Information

VIEW ALL SECTIONS

FIND ADDITIONAL RESOURCES (also available in the left menu)

SAFETY

Report Adverse Events, FDA Safety Recalls, Presence in Breast Milk

RELATED RESOURCES

Medline Plus, Clinical Trials, PubMed, Biochemical Data Summary

MORE INFO ON THIS DRUG

View Labeling Archives, RxNorm, Get Label RSS Feed, View NDC Code(s)NEW!

NIH | NLM | Health & Human Services