


This article was downloaded by: [134.117.10.200]

On: 29 November 2014, At: 12:14

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Applicable Analysis: An International Journal

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gapa20>

An inverse source problem for a one-dimensional space-time fractional diffusion equation

Salih Tatar^a & Süleyman Ulusoy^a

^a Faculty of Education, Department of Mathematics, Zirve University, Sahinbey, Gaziantep, 27260 Turkey.

Published online: 11 Nov 2014.

To cite this article: Salih Tatar & Süleyman Ulusoy (2014): An inverse source problem for a one-dimensional space-time fractional diffusion equation, *Applicable Analysis: An International Journal*, DOI: [10.1080/00036811.2014.979808](https://doi.org/10.1080/00036811.2014.979808)

To link to this article: <http://dx.doi.org/10.1080/00036811.2014.979808>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

An inverse source problem for a one-dimensional space–time fractional diffusion equation

Salih Tatar* and Süleyman Ulusoy

Faculty of Education, Department of Mathematics, Zirve University, Sahinbey, Gaziantep 27260, Turkey

Communicated by G. N'Guérékata

(Received 30 June 2014; accepted 17 October 2014)

Fractional(nonlocal) diffusion equations replace the integer-order derivatives in space and time by their fractional-order analogues and they are used to model anomalous diffusion, especially in physics. This paper deals with a nonlocal inverse source problem for a one-dimensional space–time fractional diffusion equation $\partial_t^\beta u = -r^\beta(-\Delta)^{\alpha/2}u(t, x) + f(x)h(t, x)$ where $(t, x) \in \Omega_T := (0, T) \times \Omega$ and $\Omega = (-1, 1)$. At first we define and analyze the direct problem for the space–time fractional diffusion equation. Later we define the inverse source problem. Furthermore, we set up an operator equation $A_r f(x) + \Theta(x) = f(x)$ and derive the relation between the solutions of the operator equation and the inverse source problem. We also prove some important properties of the operator A_r . By using these properties and analytic Fredholm theorem, we prove that the inverse source problem is well posed, i.e. $f(t, x)$ can be determined uniquely and depends continuously on additional data $u(T, x)$, $x \in \Omega$.

Keywords: fractional derivative; fractional Laplacian; inverse source problem; Mittag-Leffler function

AMS Subject Classifications: 34A12; 35R11; 35R30; 65F22

1. Introduction

In this paper, we consider an initial-boundary value problem for a space–time fractional equation

$$\begin{cases} \frac{\partial^\beta}{\partial t^\beta} u(t, x) = -r^\beta(-\Delta)^{\alpha/2}u(t, x) + f(x)h(t, x), & (t, x) \in \Omega_T, \\ u(t, -1) = u(t, 1) = 0, & 0 < t < T, \\ u(0, x) = 0, & x \in \Omega, \end{cases} \quad (1.1)$$

where $\Omega_T := (0, T) \times \Omega$, $\Omega = (-1, 1)$, $r > 0$ is a parameter, $f(x) \in L_2(\Omega)$, $h(t, x) \in C^1([0, T]; L^\infty(\Omega))$ are given functions, $\beta \in (0, 1)$, $\alpha \in (1, 2)$ are fractional order of the time and the space derivatives, respectively, and $T > 0$ is a final time.

*Corresponding author. Email: salih.tatar@zirve.edu.tr

The fractional-time derivative considered here is the Caputo fractional derivative of order $0 < \beta < 1$ and is defined by

$$\frac{\partial^\beta f(t)}{\partial t^\beta} := \frac{1}{\Gamma(1-\beta)} \int_0^t \frac{\partial f(r)}{\partial r} \frac{dr}{(t-r)^\beta}, \quad (1.2)$$

where Γ is the Gamma function. This was intended to properly handle initial values,[1–3] since its Laplace transform(LT) $s^\beta \tilde{f}(s) - s^{\beta-1} f(0)$ incorporates the initial value in the same way as the first derivative. Here, $\tilde{f}(s)$ is the usual Laplace transform. It is well known that the Caputo derivative has a continuous spectrum,[2] with eigenfunctions given in terms of the Mittag-Leffler function

$$E_\beta(z) := \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(1+\beta k)}.$$

In fact, it is easy to see that, $f(t) = E_\beta(-\lambda t^\beta)$ solves the eigenvalue equation

$$\frac{\partial^\beta f(t)}{\partial t^\beta} = -\lambda f(t), \quad f(0) = 1$$

for any $\lambda > 0$. This is easily verified by differentiating term by term and using the fact that t^p has Caputo derivative $t^{p-\beta} \frac{\Gamma(p+1)}{\Gamma(p+1-\beta)}$ for $p > 0$ and $0 < \beta \leq 1$. $0 < \beta < 1$ is taken for slow diffusion, and is related to the parameter specifying the large-time behavior of the waiting-time distribution function, see [4] and some of the references cited therein.

For $0 < \alpha < 2$, $(-\Delta)^{\alpha/2} u$ denotes the fractional Laplacian of u . It turns out that it is easier to define it by using the spectral decomposition of the Laplace operator: We take $\{\bar{\lambda}_k, \psi_k\}$ the eigenvalues and corresponding eigenvectors of the Laplacian operator in Ω with Dirichlet boundary conditions on $\partial\Omega$:

$$\begin{cases} -\Delta \psi_k = \bar{\lambda}_k \psi_k, & \text{in } \Omega, \\ \psi_k = 0, & \text{on } \partial\Omega. \end{cases}$$

We then define the operator $(-\Delta)^{\alpha/2}$ by

$$(-\Delta)^{\alpha/2} u := \sum_{k=0}^{\infty} c_k \psi_k(x) \mapsto - \sum_{k=0}^{\infty} c_k \bar{\lambda}_k^{\alpha/2} \psi_k(x),$$

which maps $H_0^\alpha(\Omega)$ onto $L_2(\Omega)$.

If f is C^1 -function on $[0, \infty)$ satisfying $|f'(t)| \leq C t^{\gamma-1}$ for some $\gamma > 0$, then by (1.2), the Caputo derivative $\frac{\partial^\beta f(t)}{\partial t^\beta}$ of f exists for all $t > 0$ and the derivative is continuous in $t > 0$. Kilbas et al. [5] and Podlubny [4] can be referred for further properties of the Caputo derivative.

In (1.1), the term $f(x)h(t, x)$ models a source term, and it is important to determine $f(x)$ for realizing observation data. That is to say, our main goal in this paper is: Let $r > 0$ be fixed. Determine $u(t, x) = u(r, f)(t, x)$ and $f(x)$ for $t \in (0, T)$ and $x \in \Omega$ satisfying (1.1) and

$$u(T, x) = \varphi(x), \quad x \in \bar{\Omega}. \quad (1.3)$$

This inverse problem is by final overdetermining data. We prove that the inverse problem is well-posed in the sense of Hadamard except for a finite set of $r > 0$. Our main tools

are the representation formula of the solution to (1.1), the theory of analytic perturbation of linear operators (see e.g. [6]), and the uniqueness in the inverse problem (1.1)–(1.3). For similar studies, one can refer to [7–9] and some of the references cited therein. The main difference from these studies is that the current study includes both space and time fractional derivatives.

This paper is organized as follows: In the next section, formulation of the direct and inverse problems along with their analyses is presented. Section 3 includes some properties of the operator A_r and the final section of the paper includes both the statement and the proof of the main result of the paper.

2. Formulations and analyses of the direct and inverse problems

In this section, we define the direct and the inverse source problems. For given inputs β , α , $f(x)$, and $h(t, x)$, the problem (1.1) is called the direct (forward) problem. Throughout this paper, we assume the following condition on $h(t, x)$

$$|h(T, x)| \geq \delta > 0, \quad x \in \overline{\Omega}. \quad (2.1)$$

Like most direct problems of the mathematical physics, the problem (1.1) is well-posed, see the very recent interesting paper [10] for details.

The formal solution to the direct problem (1.1) is given in the form (see [11] for details)

$$u(t, x) = \sum_{n=1}^{\infty} \left(\int_0^t \tau^{\beta-1} E_{\beta, \beta}(-\lambda_n r^{\beta} \tau^{\beta}) \left\langle f(x) h(t - \tau, x), \psi_n(x) \right\rangle d\tau \right) \psi_n(x), \quad (2.2)$$

where $\lambda_n = (\bar{\lambda}_n)^{\alpha/2}$, $\bar{\lambda}_n$ and $\{\psi_n\}_{n \geq 1}$ are eigenvalues and eigenvectors of the classical Laplace operator $-\Delta$, respectively, i.e. $-\Delta \psi_n = \bar{\lambda}_n \psi_n$. A simple calculation yields $\bar{\lambda}_n = \frac{n^2 \pi^2}{4}$ hence $\lambda_n = (\frac{n\pi}{2})^{\alpha}$ with $\psi_n(x) = \sin(\frac{n\pi x}{2})$ when n is even and $\psi_n(x) = \cos(\frac{n\pi x}{2})$ when n is odd. $\langle \cdot, \cdot \rangle$ denotes the standard inner product on $L^2(\Omega)$ and $E_{\alpha, \beta}(z)$ is the generalized Mittag-Leffler function defined as follows

$$E_{\alpha, \beta}(z) := \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}, \quad z \in \mathbb{C},$$

where $\alpha > 0$ and $\beta \in \mathbb{R}$ are arbitrary constants, Γ is the Gamma function. We note that $\{\bar{\lambda}_n\}_{n \geq 1}$ is a sequence of positive numbers $0 < \bar{\lambda}_1 \leq \bar{\lambda}_2 \leq \dots$, $\{\psi_n\}_{n \geq 1}$ is an orthonormal basis for $L_2(\Omega)$. It is proved in [11] that (2.2) is the generalized solution to the problem (1.1) that can be interpreted as the solution in the classical sense under certain additional conditions. For practical use in latter sections, if we use the substitution $\tau^* = t - \tau$ (later replace τ^* by τ) in (2.2), we get the following useful formula for the solution of (1.1)

$$u(t, x) = \sum_{n=1}^{\infty} \left(\int_0^t (t - \tau)^{\beta-1} E_{\beta, \beta}(-\lambda_n r^{\beta} (t - \tau)^{\beta}) \times \left\langle f(x) h(\tau, x), \psi_n(x) \right\rangle d\tau \right) \psi_n(x). \quad (2.3)$$

The following theorems indicate some important properties of the Mittag-Leffler function, (see the formula (1.83) and Theorem 1.6 in [4] respectively) which provide technical convenience in ensuing theorems ahead.

LEMMA 2.1 *If $\alpha < 2$, β is arbitrary real number, μ is such that $\frac{\pi\alpha}{2} < \mu < \min\{\pi\alpha, \pi\}$, $\mu \leq |\arg(z)| \leq \pi$, and C_0 is a real constant, then*

$$|E_{\alpha,\beta}(z)| \leq \frac{C_0}{1 + |z|}.$$

LEMMA 2.2 *The following equality holds for $\lambda > 0$, $\alpha > 0$ and $m \in \mathbb{N}$*

$$\frac{d^m}{dt^m} E_{\alpha,1}(-\lambda t^\alpha) = -\lambda t^{\alpha-m} E_{\alpha,\alpha-m+1}(-\lambda t^\alpha), \quad t > 0.$$

Now we show that the series on the right-hand side of (2.3) is convergent uniformly in $x \in \Omega$ and $t \in (0, T]$. By using Theorem 2.1, Cauchy–Schwarz inequality, the analytical formula of $\psi_n(x)$, the assumption $f \in L_2$, boundedness of the function $h(t, x)$, and the substitution $\tau = t(1 - s)$ in the integral appearing at the third step we get

$$\begin{aligned} & \left| \sum_{n=1}^{\infty} \left(\int_0^t (t-\tau)^{\beta-1} E_{\beta,\beta}(-\lambda_n r^\beta (t-\tau)^\beta) \langle f(x)h(\tau, x), \psi_n(x) \rangle d\tau \right) \psi_n(x) \right| \\ & \leq C_0 \sum_{n=1}^{\infty} \left| \int_0^t \frac{(t-\tau)^{\beta-1}}{1 + \lambda_n r^\beta (t-\tau)^\beta} \|fh\| d\tau \right| \\ & \leq C_1 \sum_{n=1}^{\infty} \left| \int_0^t \frac{(t-\tau)^{\beta-1}}{1 + \lambda_n r^\beta (t-\tau)^\beta} d\tau \right| \\ & = C_1 \sum_{n=1}^{\infty} \left| \int_0^1 \frac{s^{\beta-1} t^\beta}{1 + \lambda_n r^\beta s^\beta t^\beta} ds \right|. \end{aligned} \quad (2.4)$$

Next we apply Young's inequality $ab \leq \frac{a^p}{p} + \frac{b^q}{q}$ to the numerator $1 + \lambda_n r^\beta s^\beta t^\beta$ with $a = (1 - \mu)^{-(1-\mu)}$, $b = (\lambda_n r^\beta s^\beta t^\beta)^\mu \mu^{-\mu}$, $p = \frac{1}{1-\mu}$, $q = \frac{1}{\mu}$ and $\mu \in (0, 1)$ to arrive

$$C(\mu)(\lambda_n r^\beta s^\beta t^\beta)^\mu \leq 1 + \lambda_n r^\beta s^\beta, \quad (2.5)$$

where $C(\mu) = (1 - \mu)^{-(1-\mu)} \mu^{-\mu}$. Therefore, we have

$$\sum_{n=1}^{\infty} \left| \int_0^1 \frac{s^{\beta-1} t^\beta}{1 + \lambda_n r^\beta s^\beta t^\beta} ds \right| \leq \sum_{n=1}^{\infty} \left| \frac{t^\beta}{C(\mu)(\lambda_n r^\beta t^\beta)^\mu} \int_0^1 \frac{s^{\beta-1}}{s^{\beta\mu}} ds \right|. \quad (2.6)$$

Since $\mu \in (0, 1)$ and $\beta > 0$, the integral appearing at the right-hand side of (2.6) is always convergent. Thus, there exist a constant $C_2 > 0$ so that the following holds

$$\sum_{n=1}^{\infty} \left| \int_0^1 \frac{s^{\beta-1} t^{\beta}}{1 + \lambda_n r^{\beta} s^{\beta} t^{\beta}} ds \right| \leq C_2 \frac{t^{\beta}}{C(\mu)(rt)^{\beta\mu}} \sum_{n=1}^{\infty} \frac{1}{(\lambda_n)^{\mu}}. \quad (2.7)$$

By $\lambda_n = \left(\frac{n\pi}{2}\right)^{\alpha}$ and $\mu \in (0, 1)$ for the series in (2.7) to be convergent α must satisfy $\alpha\mu > 1$ so that we must have $\alpha > 1$ (we take $\alpha \in (1, 2)$). By (2.4), we conclude that the series on the right-hand side of (2.3) is convergent uniformly in $x \in \Omega$ and $t \in (0, T]$.

In the proof of Lemma 2.4, we employ Young's inequality for convolutions. For the sake of the completeness, we recall it here in the form we need.

LEMMA 2.3 [12] Suppose that f is in $L_p(0, \infty)$, g is in $L_q(0, \infty)$, $\frac{1}{p} + \frac{1}{q} = \frac{1}{r} + 1$ with $1 \leq p$ and $q, r \leq \infty$. Then

$$\|f * g\|_r \leq \|f\|_p \|g\|_q,$$

where $f * g$ is convolution of f and g , i.e. $(f * g)(t) = \int_0^t f(\tau)g(t - \tau) d\tau$.

Now we prove the following theorem which will be useful for the proof of the main result of the paper.

THEOREM 2.1 Let $f(x) \in L_2(\Omega)$, $h(t, x) \in C^1([0, T]; L^{\infty}(\Omega))$. Then there exists a unique weak solution of the problem (1.1) such that $u \in L_2(0, T; H_0^{\alpha}(\Omega))$ and $\frac{\partial^{\beta}}{\partial t^{\beta}} u \in L_2((0, T) \times \Omega)$. Moreover, there exists a constant C such that the following inequality holds

$$\|u\|_{L_2(0, T; H_0^{\alpha}(\Omega))} + \|\partial_t^{\beta} u\|_{L_2((0, T) \times \Omega)} \leq C \|f\|_{L_2(\Omega)}. \quad (2.8)$$

Before the proof of this theorem, we recall the fractional Sobolev space

$$H_0^{\alpha}(\Omega) := \left\{ u = \sum_{n=1}^{\infty} a_n \psi_n : \|u\|_{H_0^{\alpha}}^2 = \sum_{n=1}^{\infty} a_n^2 \bar{\lambda}_n^{\alpha} < +\infty \right\}, \quad (2.9)$$

with the following equivalence

$$\|u\|_{H_0^{\alpha}(\Omega)} = \|(-\Delta)^{\frac{\alpha}{2}} u\|_{L_2(\Omega)}. \quad (2.10)$$

Proof The existence and uniqueness of the weak solution can be established following the arguments in [2, 7, 11]. So we only prove (2.8).

By using Theorem 2.1, Lemma 2.2, and $E_{\beta,\beta}(-\eta) \geq 0$, $0 < \beta < 1$, we have

$$\begin{aligned}
 \int_0^\eta |t^{\beta-1} E_{\beta,\beta}(-\lambda_n t^\beta)| dt &= \int_0^\eta t^{\beta-1} E_{\beta,\beta}(-\lambda_n t^\beta) dt \\
 &= -\frac{1}{\lambda_n} \int_0^\eta \frac{d}{dt} E_{\beta,1}(-\lambda_n t^\beta) dt \\
 &= \frac{1}{\lambda_n} (1 - E_{\beta,1}(-\lambda_n \eta^\beta)).
 \end{aligned} \tag{2.11}$$

Following [4, p140], by means of the Laplace transform we see that

$$\begin{aligned}
 \frac{\partial^\beta}{\partial t^\beta} \int_0^t \left\langle f(x)h(\tau, x), \psi_n(x) \right\rangle (t-\tau)^{\beta-1} E_{\beta,\beta}(-\lambda_n r^\beta (t-\tau)^\beta) d\tau \\
 = -\lambda_n \int_0^t \left\langle f(x)h(\tau, x), \psi_n(x) \right\rangle (t-\tau)^{\beta-1} E_{\beta,\beta}(-\lambda_n r^\beta (t-\tau)^\beta) d\tau \\
 + \left\langle f(x)h(t, x), \psi_n(x) \right\rangle.
 \end{aligned} \tag{2.12}$$

From Lemma 2.3, (2.11) and (2.12) we have

$$\begin{aligned}
 &\left\| \frac{\partial^\beta}{\partial t^\beta} \int_0^t \left\langle f(x)h(\tau, x), \psi_n(x) \right\rangle (t-\tau)^{\beta-1} E_{\beta,\beta}(-\lambda_n r^\beta (t-\tau)^\beta) d\tau \right\|_{L_2(0,T)} \\
 &\leq C_3 \left\| \left\langle f(x)h(t, x), \psi_n(x) \right\rangle \right\|_{L_2(0,T)} + C_4 \left\| \left\langle f(x)h(t, x), \psi_n(x) \right\rangle \right\|_{L_2(0,T)} \\
 &\quad \times \left\| \lambda_n t^{\beta-1} E_{\beta,\beta}(-\lambda_n r^\beta t^\beta) \right\|_{L_1(0,T)} \leq C_5 \left\| \left\langle f(x)h(t, x), \psi_n(x) \right\rangle \right\|_{L_2(0,T)}.
 \end{aligned} \tag{2.13}$$

By using (2.13) and the assumption $h(t, x) \in C^1([0, T]; L^\infty(\Omega))$, we get

$$\begin{aligned}
 \left\| \frac{\partial^\beta}{\partial t^\beta} u \right\|_{L_2((0,T) \times \Omega)}^2 &= \sum_{n=1}^\infty \int_0^T \left| \frac{\partial^\beta}{\partial t^\beta} \left(\int_0^t \left\langle f(x)h(\tau, x), \psi_n(x) \right\rangle (t-\tau)^{\beta-1} \right. \right. \\
 &\quad \times \left. \left. E_{\beta,\beta}(-\lambda_n r^\beta (t-\tau)^\beta) d\tau \right) \right|^2 dt \leq C_5 \sum_{n=1}^\infty \int_0^T \left| \left\langle f(x), \psi_n(x) \right\rangle \right|^2 dt \\
 &\leq C_6 \|f\|_{L_2(\Omega)}^2.
 \end{aligned} \tag{2.14}$$

By (2.10), we see also $\|u\|_{L_2(0,T;H_0^\alpha(\Omega))} \leq C_7 \|f\|_{L_2(\Omega)}$. This completes the proof. \square

Next, we define the inverse problem. The inverse problem here consists of determining the pair of the unknown functions $\{u(t, x), f(x)\}$ in the space-time fractional diffusion problem (1.1) from the final overdetermining data (also called additional condition or

measured output data) (1.3), for the given function $h(t, x)$, the orders of time and space derivatives β , α , and a fixed $r > 0$, that is, the problem (1.1), (1.3) is called the inverse problem.

Now we reformulate the inverse problem. For this purpose, we define the following operator equation

$$A_r f(x) + \Theta(x) = f(x), \quad (2.15)$$

where $A_r(f) : L_2(\Omega) \longrightarrow L^2(\Omega)$, $A_r(f)$ and $\Theta(x)$ defined by

$$A_r f(x) = \frac{\frac{\partial^\beta}{\partial t^\beta} u(r, f)(T, x)}{h(T, x)}, \quad \Theta(x) = -\frac{r^\beta (\Delta^{\alpha/2} \varphi)(x)}{h(T, x)}, \quad (2.16)$$

respectively. Here we denote $u = u(r, f)$ to emphasize the dependence of the solution $u(t, x)$ of (1.1) to both $r > 0$ and $f(x)$. The following lemma indicates the relationship between the operator Equation (2.15) and the inverse problem (1.1), (1.3).

LEMMA 2.4 *Let $I \subset (0, \infty)$ and $r \in I$ be fixed. Then the operator Equation (2.15) has a solution (a unique solution) $f \in L_2(\Omega)$ if and only if the inverse problem (1.1), (1.3) has a solution (a unique solution) $\{u(r, f), f\} \in L_2(0, T; H_0^\alpha(\Omega)) \times L_2(\Omega)$.*

Proof First, suppose that the operator Equation (2.15) has a solution $f \in L_2(\Omega)$. Then if we substitute this $f(x)$ into the one dimensional space–time fractional diffusion equation, we have a unique solution $u(r, f)$ to (1.1), see [7]. Now we show that the function $u(r, f)$ satisfies the additional condition (1.3). For this purpose, let's assume that

$$u(T, x) = \varphi_1(x), \quad x \in \Omega.$$

Since $u \in L_2(0, T; H_0^\alpha(\Omega))$, we have $\varphi_1(x) \in H_0^\alpha(\Omega)$. By (2.15), we conclude that $-r^\beta (-\Delta)^{\alpha/2} (\varphi_1 - \varphi) = 0$. Since the Fractional Laplacian operator is invertible, $r \neq 0$ and $\varphi_1(x) - \varphi(x) = 0$, $x \in \Omega$ we get $\varphi(x) = \varphi_1(x)$, $x \in \Omega$. The converse of the assertion is directly seen in rewriting the space–time fractional diffusion equation in (2.15). The uniqueness part of the Lemma 2.4 is a direct result of the unique solvability of the problem (1.1) and the first part of the Lemma 2.4. \square

The next section is devoted to show some important properties of the operator A_r , defined by (2.16). These properties with Lemma 2.4 will be used to prove the main result of the paper.

3. Some properties of the operator A_r

This section is devoted to state and prove some properties of the operator A_r which are useful tools for a certain answer to the question of the unique solvability of the inverse problem (1.1), (1.3).

THEOREM 3.1 *For $r \in I$, the operator $A_r : L_2(\Omega) \rightarrow L_2(\Omega)$ is a compact operator.*

Proof By (2.2) we have

$$\begin{aligned} \frac{\partial^\beta}{\partial t^\beta} u(t, x) &= f(x)h(t, x) \\ &- \sum_{n=1}^{\infty} \left(\int_0^t \tau^{\beta-1} E_{\beta, \beta}(-\lambda_n r^\beta \tau^\beta) \left\langle f(x)h(t-\tau, x), \psi_n(x) \right\rangle d\tau \right) r^\beta \lambda_n \psi_n(x). \end{aligned} \quad (3.1)$$

By Lemma 2.2 and integration by parts, we have

$$\begin{aligned} &\int_0^t \tau^{\beta-1} E_{\beta, \beta}(-\lambda_n r^\beta \tau^\beta) \left\langle f(x)h(t-\tau, x), \psi_n(x) \right\rangle d\tau \\ &= \int_0^t \left\langle f(x)h(t-\tau, x), \psi_n(x) \right\rangle \frac{d}{d\tau} \left(-\frac{1}{\lambda_n r^\beta} E_\beta(-\lambda_n r^\beta \tau^\beta) \right) d\tau \\ &= \frac{1}{\lambda_n r^\beta} \left(\left\langle f(x)h(t, x), \psi_n(x) \right\rangle - \left\langle f(x)h(0, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta t^\beta) \right) \\ &\quad + \frac{1}{\lambda_n r^\beta} \int_0^t \left\langle f(x) \frac{\partial}{\partial \tau} h(t-\tau, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta \tau^\beta) d\tau. \end{aligned} \quad (3.2)$$

If we substitute (3.2) in (3.1), we get

$$\begin{aligned} \frac{\partial^\beta}{\partial t^\beta} u(t, x) &= \sum_{n=1}^{\infty} \left(\left\langle f(x)h(0, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta t^\beta) \right) \psi_n(x) \\ &\quad - \sum_{n=1}^{\infty} \left(\int_0^t \left\langle f(x) \frac{\partial}{\partial \tau} h(t-\tau, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta \tau^\beta) d\tau \right) \psi_n(x). \end{aligned} \quad (3.3)$$

Consequently, by using (2.9), (2.1), Cauchy–Schwarz inequality and Parseval’s identity, we obtain

$$\begin{aligned} &\left\| \frac{\partial^\beta}{\partial t^\beta} u(T, x) \right\|_{H_0^\alpha(\Omega)}^2 \\ &= \sum_{n=1}^{\infty} \bar{\lambda}_n^\alpha \left(\left\langle f(x)h(0, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta T^\beta) \right. \\ &\quad \left. - \int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T-\tau, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta \tau^\beta) d\tau \right)^2 \\ &\leq 2 \sum_{n=1}^{\infty} \bar{\lambda}_n^\alpha \left\{ \left\langle f(x)h(0, x), \psi_n(x) \right\rangle^2 \left(E_\beta(-\lambda_n r^\beta T^\beta) \right)^2 \right. \\ &\quad \left. + \left(\int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T-\tau, x), \psi_n(x) \right\rangle E_\beta(-\lambda_n r^\beta \tau^\beta) d\tau \right)^2 \right\} \end{aligned}$$

$$\begin{aligned}
 &\leq 2 \sum_{n=1}^{\infty} \bar{\lambda}_n^{\alpha} \left\{ \left(\frac{C_8}{1 + \lambda_n r^{\beta} T^{\beta}} \right)^2 \left\langle f(x) h(0, x), \psi_n(x) \right\rangle^2 \right. \\
 &\quad \left. + \int_0^T \left(\frac{C_8}{1 + \lambda_n r^{\beta} (T - \tau)^{\beta}} \right)^2 d\tau \int_0^T \left(\left\langle f(x) \frac{\partial}{\partial \tau} h(\tau, x), \psi_n(x) \right\rangle \right)^2 d\tau \right\} \\
 &\leq 2 \sum_{n=1}^{\infty} \bar{\lambda}_n^{\alpha} \left\{ \left(\frac{C_8}{\lambda_n r^{\beta} T^{\beta}} \right)^2 \left\langle f(x) h(0, x), \psi_n(x) \right\rangle^2 \right. \\
 &\quad \left. + \int_0^T \left(\frac{C_8}{1 + \lambda_n r^{\beta} (T - \tau)^{\beta}} \right)^2 d\tau \int_0^T \left(\left\langle f(x) \frac{\partial}{\partial \tau} h(\tau, x), \psi_n(x) \right\rangle \right)^2 d\tau \right\} \\
 &\leq C_9 \left\{ \|f(x) h(0, x)\|_{L_2(\Omega)}^2 + \frac{T^{1-\beta}}{r^{\beta}(1-\beta)} \left\| f(x) \frac{\partial}{\partial \tau} h(\tau, x) \right\|_{L_2(\Omega_T)}^2 \right\} \\
 &\leq C_{10} \|f(x)\|_{L_2(\Omega)}^2.
 \end{aligned} \tag{3.4}$$

Since $H_0^{\alpha}(\Omega)$ is compactly embedded in $L_2(\Omega)$, see [13] and [14], the operator $h(T, x)A_r f$ is compact. Finally by using (2.1), we conclude that the operator $A_r : L_2(\Omega) \rightarrow L_2(\Omega)$ is compact. This completes the proof. \square

THEOREM 3.2 $A_r f : I \rightarrow L_2(\Omega)$, defined by (2.16), is real analytic in $r \in I$ for arbitrarily fixed $f \in L_2(\Omega)$.

Proof We use the last theorem on page 65 of [15]. To show the real analyticity in $r \in I$ for $u(r, f) := u(r)$, it is enough to prove the following

- (a) $\frac{\partial^{\beta}}{\partial t^{\beta}} u(r) \in C^{\infty}(0, \infty; L_2(\Omega))$,
- (b) $\left\| \frac{d^m}{dr^m} \left(\frac{\partial^{\beta}}{\partial r^{\beta}} u(r)(T, x) \right) \right\|_{L_2(\Omega)} \leq M \eta^{-m} m!$, for $m \in \mathbb{N}$, $r \in J := [r_0, R_0] \subset I$,
 $M > 0$ and $\eta > 0$.

By (3.4) and the compact embedding, we have

$$\left\| \frac{\partial^{\beta}}{\partial t^{\beta}} u(r)(T, x) \right\|_{L_2(\Omega)} \leq M.$$

By (3.3) and $\frac{d^m}{dr^m} E_{\beta}(-\lambda r^{\beta} \tau^{\beta}) = -\lambda_n r^{\beta-m} \tau^{\beta} E_{\beta, \beta-m+1}(-\lambda r^{\beta} \tau^{\beta})$ we have

$$\begin{aligned}
 \frac{d^m}{dr^m} \left(\frac{\partial^{\beta}}{\partial t^{\beta}} u(T, x) \right) &= \sum_{n=1}^{\infty} \left(\left\langle f(x) h(0, x), \psi_n(x) \right\rangle (-\lambda_n) r^{\beta-m} T^{\beta} E_{\beta, \beta-m+1}(-\lambda_n r^{\beta} T^{\beta}) \right. \\
 &\quad \left. - \int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T - \tau, x), \psi_n(x) \right\rangle (-\lambda_n) r^{\beta-m} \tau^{\beta} E_{\beta, \beta-m+1}(-\lambda_n r^{\beta} \tau^{\beta}) d\tau \right) \psi_n(x).
 \end{aligned} \tag{3.5}$$

Therefore, we obtain

$$\begin{aligned}
& \left\| \frac{d^m}{dr^m} \left(\frac{\partial^\beta}{\partial t^\beta} u(T, x) \right) \right\|_{L_2(\Omega)}^2 \\
& \leq 2 \sum_{n=1}^{\infty} \left\{ \left\langle f(x)h(0, x), \psi_n(x) \right\rangle^2 \lambda_n^2 r^{2\beta-2m} T^\beta \left(\frac{C_{11}}{1 + \lambda_n r^\beta T^\beta} \right)^2 \right. \\
& \quad \left. + \left(\int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T - \tau, x), \psi_n(x) \right\rangle^2 \lambda_n^2 r^{2\beta-2m} d\tau \right) \left(\int_0^T \tau^{2\beta} \left(\frac{C_{11}}{1 + \lambda_n r^\beta T^\beta} \right)^2 d\tau \right) \right\} \\
& \leq \frac{2C_{11}^2}{r^{2m}} \left\{ \|f(x)h(0, x)\|_{L_2(\Omega)}^2 + T \|f \frac{\partial}{\partial \tau} h\|_{L_2((0,T) \times \Omega)}^2 \right\} \leq M^2 r^{-2m}, \tag{3.6}
\end{aligned}$$

where $M^2 = 2C_{11}^2 \left\{ \|f(x)h(0, x)\|_{L_2(\Omega)}^2 + T \|f \frac{\partial}{\partial \tau} h\|_{L_2((0,T) \times \Omega)}^2 \right\}$. Finally, by (2.1) and (2.16) we conclude that

$$\left\| \frac{d^m}{dr^m} A_r f \right\|_{L_2(\Omega)} \leq M \delta^{-1} r_0^{-m}.$$

The proof is completed. \square

THEOREM 3.3 *There exists a constant $0 < \mathcal{C}(r) < 1$ such that*

$$\|A_r f\|_{L_2(\Omega)} \leq \mathcal{C}(r) \|f\|_{L_2(\Omega)}, \tag{3.7}$$

where $R^* < r$ and $R^* > 0$ is a large number.

Proof By (2.1), Theorem 2.1 and (3.3) we have

$$\begin{aligned}
\|A_r(f)\|_{L_2(\Omega)}^2 & \leq \frac{2}{\delta^2} \sum_{n=1}^{\infty} \underbrace{\left(\left(\frac{C_{11}}{1 + \lambda_n r^\beta T^\beta} \right)^2 \left\langle f(x)h(0, x), \psi_n(x) \right\rangle^2 \right)}_{=: I_1} \\
& \quad + \underbrace{\left(\int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T - \tau, x), \psi_n(x) \right\rangle^2 E_\beta(-\lambda_n r^\beta \tau^\beta) d\tau \right)}_{=: I_2} \tag{3.8}
\end{aligned}$$

Now we estimate I_1 and I_2 . For I_1 we have

$$I_1 \leq \frac{C_{11}^2}{\lambda_1^2 r^{2\beta} T^{2\beta}} \left\langle f(x)h(0, x), \psi_n(x) \right\rangle^2. \tag{3.9}$$

For I_2 we have

$$\begin{aligned} I_2 &\leq \int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T - \tau, x), \psi_n(x) \right\rangle^2 d\tau \int_0^T \left(E_\beta(-\lambda_n r^\beta \tau^\beta) \right)^2 d\tau \\ &\leq \frac{C_{11}^2}{\lambda_1 r^\beta} \frac{T^{1-\beta}}{1-\beta} \int_0^T \left\langle f(x) \frac{\partial}{\partial \tau} h(T - \tau, x), \psi_n(x) \right\rangle^2 d\tau. \end{aligned} \quad (3.10)$$

If we substitute (3.9) and (3.10) in (3.8) and use Parseval's identity, we get

$$\begin{aligned} &\left\| A_r f \right\|_{L_2(\Omega)}^2 \\ &\leq \underbrace{\frac{C_{11}^2}{\delta^2} \left(\frac{1}{\lambda_1^2 r^{2\beta} T^{2\beta}} \|h(0, x)\|_{L_\infty(\Omega)}^2 + \frac{1}{\lambda_1 r^\beta} \frac{T^{1-\beta}}{1-\beta} \|\partial_t h\|_{L_\infty((0,T) \times \Omega)}^2 \right)}_{=: I_3} \|f\|_{L_2(\Omega)}^2. \end{aligned} \quad (3.11)$$

For large $r > 0$, $I_3 < 1$. This completes the proof. \square

The following corollary is obtained directly by Theorem 3.3.

COROLLARY 3.1 *I is not an eigenvalue of the operator A_r for large $r > 0$.*

4. Proof of the main result

In the proof of the main result of the paper, we will use the Analytic Fredholm Theorem. For the sake of the completeness we recall it here, (see page 266, Theorem 8.92 in [16]).

THEOREM 4.1 (Analytic Fredholm Theorem) *Let $G \subset \mathbb{C}$ be a domain and L be a bounded linear operator on G . If L is real analytic on G and the operator $L(\lambda)$ is compact for each $\lambda \in G$ then either $(I - L(\lambda))^{-1}$ does not exist for any $\lambda \in G$ or $(I - L(\lambda))^{-1}$ exists for every $\lambda \in G \setminus S$ where S is a discrete set in G .*

Now we are ready to state and prove the main result of the paper.

THEOREM 4.2 *There exists a finite set $S \subset I$ such that for $r \in I \setminus S$ and $\varphi \in H_0^\alpha(\Omega)$, the inverse problem (1.1), (1.3) has a unique solution. Moreover there exists a constant $C_{12} > 0$ such that*

$$\|f\|_{L^2(\Omega)} + \|u\|_{L_2(0,T;H_0^\alpha(\Omega))} + \|\partial_t^\beta u\|_{L_2(0,T;L_2(\Omega))} \leq C_{12} \|\varphi\|_{H_0^\alpha(\Omega)}. \quad (4.1)$$

Proof If we use Corollary 3.1 and apply Theorem 4.1 to the operator A_r , we see that the first alternative can not occur that is, $(I - A_r)^{-1}$ exists for every $\lambda \in I \setminus S$ where S is a discrete set in I . By Lemma 2.1, we conclude that the inverse problem (1.1), (1.3) is

uniquely solvable. By (2.10), (2.15), and Theorem 3.3, we also conclude

$$\|f\|_{L_2(\Omega)} \leq C_{12} \|\varphi\|_{H_0^g(\Omega)}. \quad (4.2)$$

(4.2) with (2.8) implies (4.1). The proof is completed. \square

Acknowledgements

The research has been supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) through the project Nr 113F373, also by the Zirve University Research Fund. The authors thank the referee for his/her very careful reading and for pointing out several mistakes as well as for the useful comments and suggestions.

References

- [1] Caputo M. Linear models of diffusion whose Q is almost frequency independent, part II. *Geophys. J. R. Astron. Soc.* 1967;13:529–539.
- [2] Chen QZ, Meerschaert MM, Nane E. Space-time fractional diffusion on bounded domains. *J. Math. Anal. Appl.* 2012;393:479–488.
- [3] Eidelman SD, Ivasyshen SD, Kochubei AN. *Analytic methods in the theory of differential and pseudo-differential equations of parabolic type.* Basel: Birkhäuser; 2004.
- [4] Podlubny I. *Fractional differential equations.* San Diego (CA): Academic Press; 1999.
- [5] Kilbas AA, Srivastava HM, Trujillo JJ. *Theory and applications of fractional differential equations.* Amsterdam: Elsevier; 2006.
- [6] Kato T. *Perturbation theory for linear operators.* Berlin: Springer-Verlag; 1976.
- [7] Sakamoto K, Yamamoto M. Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *J. Math. Anal. Appl.* 2011;382:426–447.
- [8] Sakamoto K, Yamamoto M. Inverse heat source problem from time distributing overdetermination. *Appl. Anal.* 2009;88:735–748.
- [9] Sakamoto K, Yamamoto M. Inverse source problem with a final overdetermination for a fractional diffusion equation. *Math. Control Relat. Fields.* 2011;1:509–518.
- [10] Kolokoltsov VN, Veretennikova MA. Well-posedness and regularity of the Cauchy problem for nonlinear fractional in time and space equations. *Fractional Differ. Calculus.* Forthcoming.
- [11] Luchko Y. Initial-boundary-value problems for the one-dimensional time-fractional diffusion equation. *Fractional Calculus Appl. Anal.* 2012;15:141–160.
- [12] Bogachev V. *Measure theory I.* Berlin: Springer-Verlag; 2007.
- [13] Adams RA. *Sobolev spaces.* Vol. 65, Pure and applied mathematics. New York (NY): Academic Press; 1975.
- [14] Stein EM. *Singular integrals and differentiability properties of functions.* Vol. 30, Princeton Mathematical Series. Princeton (NJ): Princeton University Press; 1970.
- [15] John F. *Partial differential equations.* Berlin: Springer-Verlag; 1982.
- [16] Renardy M, Rogers RC. *An introduction to partial differential equations.* New York (NY): Springer-Verlag; 2004.