

参考文献

- [1] Akao, S. and Kouchi, H. (1992). A supernodulating mutant isolated from soybean cultivar enrei. *Soil Science and Plant Nutrition*, 38(1):183–187.
- [2] Delves, A., Higgins, A., and Gresshoff, P. (1987). Shoot control of supernodulation in a number of mutant soybeans, glycine max (l.) merr. *Australian Journal of Plant Physiology*, 14(6):689–694.
- [3] Franche, C., Lindström, K., and Elmerich, C. (2009). Nitrogen-fixing bacteria associated with leguminous and non-leguminous plants. *Plant and Soil*, 321(1):35–59.
- [4] Gremaud, M. F. and Harper, J. E. (1989). Selection and initial characterization of partially nitrate tolerant nodulation mutants of soybean. *Plant Physiology*, 89(1):169–173.
- [5] Kenjo, T., Yamaya, H., and Arima, Y. (2010). Shoot-synthesized nodulation-restricting substances of wild-type soybean present in two different high-performance liquid chromatography peaks of the ethanol-soluble medium-polarity fraction. *Soil Science and Plant Nutrition*, 56(3):399–406.
- [6] Kosslak, R. M. and Bohlool, B. B. (1984). Suppression of nodule development of one side of a split-root system of soybeans caused by prior inoculation of the other side. *Plant Physiology*, 75(1):125–130.
- [7] Schmutz, J., Cannon, S. B., Schlueter, J., Ma, J., Mitros, T., Nelson, W., Hyten, D. L., Song, Q., Thelen, J. J., Cheng, J., Xu, D., Hellsten, U., May, G. D., Yu, Y., Sakurai, T., Umezawa, T., Bhattacharyya, M. K., Sandhu, D., Valliyodan, B., Lindquist, E., Peto, M., Grant, D., Shu, S., Goodstein, D., Barry, K., Futrell-Griggs, M., Abernathy, B., Du, J., Tian, Z., Zhu, L., Gill, N., Joshi, T., Libault, M., Sethuraman, A., Zhang, X.-C., Shinozaki, K., Nguyen, H. T., and Wing, R. A. (2010). Genome sequence of the palaeopolyploid soybean. *Nature*, 463(7278):178–183.
- [8] Yamaya, H. and Arima, Y. (2004). ダイズ小植物体葉身アポプラストへの溶液連続導入法の確立と根粒超多着生形質解明への応用. *日本土壌肥科学雑誌*, 75(6):685–691.