

qPCR 08 Oct · 384-well · cDNA 1000 ng / 20.00 µL,
diluted 1:5 to 10 ng/µL · 3× replicates · wells A1–I13



		Ev 1	Ev 2	Wt 1	Wt 2	Mut1	Mut 2	Ev ap 1	Ev ap 2	Wt ap1	Wt ap 2	Mut ap 1	Mut ap 2	NTC
		1	2	3	4	5	6	7	8	9	10	11	12	13
hprt	A	1a	2a	3a	4a	5a	6a	7a	8a	9a	10a	11a	12a	13a
hprt	B	1a	2a	3a	4a	5a	6a	7a	8a	9a	10a	11a	12a	13a
hprt	C	1a	2a	3a	4a	5a	6a	7a	8a	9a	10a	11a	12a	13a
f2rl1	D	1b	2b	3b	4b	5b	6b	7b	8b	9b	10b	11b	12b	13b
f2rl1	E	1b	2b	3b	4b	5b	6b	7b	8b	9b	10b	11b	12b	13b
f2rl1	F	1b	2b	3b	4b	5b	6b	7b	8b	9b	10b	11b	12b	13b
cox2	G	1c	2c	3c	4c	5c	6c	7c	8c	9c	10c	11c	12c	13c
cox2	H	1c	2c	3c	4c	5c	6c	7c	8c	9c	10c	11c	12c	13c
cox2	I	1c	2c	3c	4c	5c	6c	7c	8c	9c	10c	11c	12c	13c

- 1

 Ev 1
- 2

 Ev 2
- 3

 Wt 1
- 4

 Wt 2
- 5

 Mut1
- 6

 Mut 2
- 7

 Ev ap 1
- 8

 Ev ap 2
- 9

 Wt ap1
- 10

 Wt ap 2
- 11

 Mut ap 1
- 12

 Mut ap 2
- 13

 NTC
- a

 hprt
- b

 f2rl1
- c

 cox2

Master mix per gene · cDNA (10 ng/µL) 2 µL per reaction added separately

GENE	WELLS	*	MIX	SYBR ×2	F primer (50 µM)	R primer (50 µM)	nfw
a hprt	39	43	344 µL	215	4.3	4.3	120.4
b f2rl1	39	43	344 µL	215	4.3	4.3	120.4
c cox2	39	43	344 µL	215	4.3	4.3	120.4