

NCERT Mathematics Formula Book
Classes 11 & 12

Compiled from NCERT Textbooks

Contents

Mathematical Constants and Symbols	5
1 Class 11 Mathematics Formulas	7
1.1 Sets	7
1.2 Relations and Functions	7
1.3 Trigonometric Functions	8
1.4 Complex Numbers	9
1.5 Linear Inequalities	9
1.6 Permutations and Combinations	9
1.7 Binomial Theorem	9
1.8 Sequence and Series	10
1.9 Straight Lines	10
1.10 Conic Sections	11
1.11 Introduction to 3D Geometry	11
1.12 Limits and Derivatives	12
1.13 Statistics	13
1.14 Probability	13
2 Class 12 Mathematics Formulas	15
2.1 Relations and Functions	15
2.2 Inverse Trigonometric Functions	15
2.3 Matrices	16
2.4 Determinants	16
2.5 Continuity and Differentiability	16
2.6 Applications of Derivatives	17
2.7 Integrals	17
2.8 Applications of Integrals	18
2.9 Differential Equations	18
2.10 Vector Algebra	19
2.11 Three Dimensional Geometry	19
2.12 Probability	20
2.13 Linear Programming	20

Mathematical Constants and Symbols

Important Mathematical Constants

Constant	Symbol	Value
Pi	π	3.141592653589793
Euler's number	e	2.718281828459045
Golden ratio	ϕ	1.618033988749895
Imaginary unit	i	$\sqrt{-1}$

Chapter 1

Class 11 Mathematics Formulas

1.1 Sets

Union of Sets: $A \cup B = \{x : x \in A \text{ or } x \in B\}$

Definition: Set containing all elements from both sets.

Intersection of Sets: $A \cap B = \{x : x \in A \text{ and } x \in B\}$

Definition: Set containing only common elements.

Complement of Set: $A' = \{x : x \in U \text{ and } x \notin A\}$

Definition: Set of all elements not in A but in universal set.

Difference of Sets: $A - B = \{x : x \in A \text{ and } x \notin B\}$

Definition: Set of elements in A but not in B.

De Morgan's Laws:

$$(A \cup B)' = A' \cap B'$$

$$(A \cap B)' = A' \cup B'$$

Definition: Complement of union equals intersection of complements and vice versa.

1.2 Relations and Functions

Relation: $R \subseteq A \times B$

Definition: Subset of Cartesian product of two sets.

Function: $f : A \rightarrow B$

Definition: Relation where each element of A maps to exactly one element of B.

Domain: Set of all possible input values

Range: Set of all possible output values

One-one Function (Injective): $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$

Definition: Function where different inputs give different outputs.

Onto Function (Surjective): Range = Codomain

Definition: Function where every element of codomain is mapped.

Bijjective Function: Both one-one and onto

Definition: Function that is both injective and surjective.

1.3 Trigonometric Functions

Basic Identities:

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

Definition: Fundamental Pythagorean identities.

Sum and Difference Formulas:

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

Definition: Formulas for trigonometric functions of sum/difference of angles.

Double Angle Formulas:

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$$

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

Definition: Formulas for trigonometric functions of double angles.

Product to Sum Formulas:

$$\sin A \cos B = \frac{1}{2} [\sin(A + B) + \sin(A - B)]$$

$$\cos A \cos B = \frac{1}{2} [\cos(A + B) + \cos(A - B)]$$

$$\sin A \sin B = \frac{1}{2} [\cos(A - B) - \cos(A + B)]$$

Definition: Conversion of products to sums.

Law of Sines: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$

Definition: Ratio of side length to sine of opposite angle is constant.

Law of Cosines:

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Definition: Relates sides and angles of any triangle.

1.4 Complex Numbers

Complex Number: $z = a + ib$ where $i^2 = -1$

Definition: Number with real and imaginary parts.

Modulus: $|z| = \sqrt{a^2 + b^2}$

Definition: Distance from origin in complex plane.

Argument: $\arg(z) = \theta = \tan^{-1} \left(\frac{b}{a} \right)$

Definition: Angle with positive real axis.

Polar Form: $z = r(\cos \theta + i \sin \theta) = re^{i\theta}$

Definition: Complex number in polar coordinates.

De Moivre's Theorem: $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$

Definition: Powers of complex numbers in polar form.

1.5 Linear Inequalities

Linear Inequality: $ax + b > 0$ (or $<, \geq, \leq$)

Definition: Inequality of first degree.

Solution Set: Set of all values satisfying the inequality

Graphical Representation: Half-plane in coordinate system

1.6 Permutations and Combinations

Factorial: $n! = n \times (n-1) \times \cdots \times 2 \times 1$

Definition: Product of all positive integers up to n.

Permutation: ${}^nP_r = \frac{n!}{(n-r)!}$

Definition: Arrangements of r objects from n distinct objects.

Combination: ${}^nC_r = \frac{n!}{r!(n-r)!}$

Definition: Selections of r objects from n distinct objects.

Circular Permutation: $(n-1)!$

Definition: Arrangements around a circle.

1.7 Binomial Theorem

Binomial Theorem:

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

Definition: Expansion of binomial raised to positive integer power.

General Term: $T_{r+1} = \binom{n}{r} a^{n-r} b^r$

Definition: $(r+1)$ th term in binomial expansion.

Middle Term:

- When n is even: $\left(\frac{n}{2} + 1\right)$ th term
- When n is odd: $\left(\frac{n+1}{2}\right)$ th and $\left(\frac{n+3}{2}\right)$ th terms

1.8 Sequence and Series

Arithmetic Progression (AP):

n^{th} term: $a_n = a + (n - 1)d$

Sum: $S_n = \frac{n}{2}[2a + (n - 1)d] = \frac{n}{2}(a + l)$

Common difference: $d = a_n - a_{n-1}$

Definition: Sequence with constant difference between terms.

Geometric Progression (GP):

n^{th} term: $a_n = ar^{n-1}$

Sum: $S_n = a \left(\frac{1 - r^n}{1 - r} \right) \quad (r \neq 1)$

Sum to infinity: $S_\infty = \frac{a}{1 - r} \quad (|r| < 1)$

Common ratio: $r = \frac{a_n}{a_{n-1}}$

Definition: Sequence with constant ratio between terms.

Harmonic Progression (HP): Reciprocals form AP

Definition: Sequence where reciprocals of terms form AP.

Arithmetic Mean: $AM = \frac{a + b}{2}$

Geometric Mean: $GM = \sqrt{ab}$

Harmonic Mean: $HM = \frac{2ab}{a + b}$

Relation: $AM \geq GM \geq HM$

1.9 Straight Lines

Distance Formula: $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Definition: Distance between two points in Cartesian plane.

Section Formula:

$$\begin{aligned}\text{Internal: } P &= \left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n} \right) \\ \text{External: } P &= \left(\frac{mx_2 - nx_1}{m-n}, \frac{my_2 - ny_1}{m-n} \right)\end{aligned}$$

Definition: Coordinates of point dividing line segment in given ratio.

Slope of Line: $m = \frac{y_2 - y_1}{x_2 - x_1} = \tan \theta$

Definition: Tangent of angle made with positive x-axis.

Equation of Line:

Slope-intercept: $y = mx + c$

Two-point: $\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$

Intercept: $\frac{x}{a} + \frac{y}{b} = 1$

Normal: $x \cos \alpha + y \sin \alpha = p$

Definition: Different forms of straight line equation.

Distance from Point to Line: $d = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$

Definition: Perpendicular distance from point to line $ax + by + c = 0$.

Angle between Lines: $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$

Definition: Acute angle between two lines with slopes m_1 and m_2 .

1.10 Conic Sections

Circle: $(x - h)^2 + (y - k)^2 = r^2$

Definition: Locus of points equidistant from center.

Parabola: $y^2 = 4ax$ (standard form)

Definition: Locus of points equidistant from focus and directrix.

Ellipse: $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

Definition: Locus of points where sum of distances to two foci is constant.

Hyperbola: $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

Definition: Locus of points where difference of distances to two foci is constant.

1.11 Introduction to 3D Geometry

Distance Formula: $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

Definition: Distance between two points in 3D space.

Section Formula:

$$\begin{aligned}\text{Internal: } P &= \left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right) \\ \text{External: } P &= \left(\frac{mx_2 - nx_1}{m-n}, \frac{my_2 - ny_1}{m-n}, \frac{mz_2 - nz_1}{m-n} \right)\end{aligned}$$

Definition: Coordinates in 3D space.

Centroid of Triangle: $G = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$

1.12 Limits and Derivatives

Limit: $\lim_{x \rightarrow a} f(x) = L$

Definition: Value function approaches as input approaches some point.

Standard Limits:

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin x}{x} &= 1 \\ \lim_{x \rightarrow 0} \frac{1 - \cos x}{x} &= 0 \\ \lim_{x \rightarrow 0} \frac{e^x - 1}{x} &= 1 \\ \lim_{x \rightarrow 0} \frac{\log(1+x)}{x} &= 1\end{aligned}$$

Definition: Important limit results.

Derivative: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

Definition: Instantaneous rate of change of function.

Standard Derivatives:

$$\begin{aligned}\frac{d}{dx}(x^n) &= nx^{n-1} \\ \frac{d}{dx}(\sin x) &= \cos x \\ \frac{d}{dx}(\cos x) &= -\sin x \\ \frac{d}{dx}(\tan x) &= \sec^2 x \\ \frac{d}{dx}(\cot x) &= -\csc^2 x \\ \frac{d}{dx}(\sec x) &= \sec x \tan x \\ \frac{d}{dx}(\csc x) &= -\csc x \cot x \\ \frac{d}{dx}(e^x) &= e^x \\ \frac{d}{dx}(a^x) &= a^x \ln a \\ \frac{d}{dx}(\ln x) &= \frac{1}{x} \\ \frac{d}{dx}(\log_a x) &= \frac{1}{x \ln a}\end{aligned}$$

Definition: Derivatives of common functions.

1.13 Statistics

Mean: $\bar{x} = \frac{\sum x_i}{n}$

Definition: Average of data.

Median: Middle value of ordered data

Mode: Most frequent value

Variance: $\sigma^2 = \frac{\sum (x_i - \bar{x})^2}{n}$

Definition: Average of squared deviations from mean.

Standard Deviation: $\sigma = \sqrt{\text{Variance}}$

Definition: Measure of data dispersion.

1.14 Probability

Probability: $P(A) = \frac{\text{Number of favorable outcomes}}{\text{Total number of outcomes}}$

Definition: Measure of likelihood of event.

Addition Rule: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

Definition: Probability of union of two events.

Multiplication Rule: $P(A \cap B) = P(A) \cdot P(B|A)$

Definition: Probability of intersection of two events.

Chapter 2

Class 12 Mathematics Formulas

2.1 Relations and Functions

Types of Relations:

- Reflexive: $(a, a) \in R$ for all a
- Symmetric: $(a, b) \in R \Rightarrow (b, a) \in R$
- Transitive: $(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R$
- Equivalence: Reflexive, symmetric and transitive

Inverse Function: $f^{-1}(f(x)) = x$

Definition: Function that reverses the mapping.

Composite Function: $(g \circ f)(x) = g(f(x))$

Definition: Application of one function to result of another.

Binary Operations: $* : A \times A \rightarrow A$

Definition: Operation that combines two elements to form another.

2.2 Inverse Trigonometric Functions

Principal Values:

$$\begin{aligned}\sin^{-1} x &\in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \\ \cos^{-1} x &\in [0, \pi] \\ \tan^{-1} x &\in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)\end{aligned}$$

Definition: Ranges of inverse trigonometric functions.

Basic Formulas:

$$\begin{aligned}\sin^{-1} x + \cos^{-1} x &= \frac{\pi}{2} \\ \tan^{-1} x + \cot^{-1} x &= \frac{\pi}{2} \\ \sec^{-1} x + \csc^{-1} x &= \frac{\pi}{2}\end{aligned}$$

Definition: Complementary relationships.

2.3 Matrices

Matrix: Rectangular array of numbers

Order: $m \times n$ (m rows, n columns)

Matrix Addition: $[A + B]_{ij} = A_{ij} + B_{ij}$

Definition: Element-wise addition.

Matrix Multiplication: $[AB]_{ij} = \sum_{k=1}^n A_{ik}B_{kj}$

Definition: Dot product of rows and columns.

Transpose: $[A^T]_{ij} = A_{ji}$

Definition: Rows become columns and vice versa.

Determinant:

$$\begin{aligned} 2 \times 2: \quad & \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc \\ 3 \times 3: \quad & \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a(ei - fh) - b(di - fg) + c(dh - eg) \end{aligned}$$

Definition: Scalar value for square matrices.

Inverse: $A^{-1} = \frac{1}{|A|} \text{adj}(A)$

Definition: Matrix that when multiplied gives identity.

2.4 Determinants

Properties:

- $|AB| = |A||B|$
- $|A^T| = |A|$
- $|kA| = k^n|A|$ for $n \times n$ matrix
- If two rows/columns identical, $|A| = 0$

Cramer's Rule: Solution of system $AX = B$ is $x_i = \frac{|A_i|}{|A|}$

Definition: Method to solve linear equations using determinants.

2.5 Continuity and Differentiability

Continuity: $\lim_{x \rightarrow c} f(x) = f(c)$

Definition: Function is continuous at point if limit equals function value.

Product Rule: $\frac{d}{dx}(uv) = u\frac{dv}{dx} + v\frac{du}{dx}$

Definition: Derivative of product of two functions.

Quotient Rule: $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$

Definition: Derivative of quotient of two functions.

Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Definition: Derivative of composite function.

Implicit Differentiation: Differentiate both sides w.r.t x

Definition: Method when y is not explicitly expressed in terms of x.

Parametric Differentiation: $\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$

Definition: When x and y are expressed in terms of parameter t.

Second Order Derivative: $\frac{d^2y}{dx^2} = \frac{d}{dx}\left(\frac{dy}{dx}\right)$

Definition: Derivative of the derivative.

2.6 Applications of Derivatives

Rate of Change: $\frac{dy}{dx}$ represents rate of change

Definition: How one quantity changes with respect to another.

Tangent: $y - y_1 = m(x - x_1)$ where $m = f'(x_1)$

Definition: Line touching curve at one point.

Normal: $y - y_1 = -\frac{1}{m}(x - x_1)$

Definition: Line perpendicular to tangent at point of contact.

Increasing Function: $f'(x) > 0$

Decreasing Function: $f'(x) < 0$

Local Maxima/Minima: $f'(x) = 0$ and check sign change

Definition: Highest/lowest points in neighborhood.

2.7 Integrals

Indefinite Integral: $\int f(x)dx = F(x) + C$ where $F'(x) = f(x)$

Definition: Anti-derivative of function.

Definite Integral: $\int_a^b f(x)dx = F(b) - F(a)$

Definition: Area under curve between limits a and b.

Standard Integrals:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad (n \neq -1)$$

$$\int \frac{1}{x} dx = \ln |x| + C$$

$$\int e^x dx = e^x + C$$

$$\int a^x dx = \frac{a^x}{\ln a} + C$$

$$\int \sin x dx = -\cos x + C$$

$$\int \cos x dx = \sin x + C$$

$$\int \sec^2 x dx = \tan x + C$$

$$\int \csc^2 x dx = -\cot x + C$$

$$\int \sec x \tan x dx = \sec x + C$$

$$\int \csc x \cot x dx = -\csc x + C$$

Definition: Integrals of common functions.

Integration by Parts: $\int u dv = uv - \int v du$

Definition: Method to integrate product of functions.

Integration by Substitution: $\int f(g(x))g'(x)dx = \int f(u)du$

Definition: Method using change of variable.

Partial Fractions: Decompose rational function

Definition: Express as sum of simpler fractions.

2.8 Applications of Integrals

Area under Curve: $A = \int_a^b f(x)dx$

Definition: Area between curve and x-axis.

Area between Curves: $A = \int_a^b [f(x) - g(x)]dx$

Definition: Area between two curves.

Volume of Revolution: $V = \pi \int_a^b [f(x)]^2 dx$

Definition: Volume when curve rotated about x-axis.

2.9 Differential Equations

Order: Highest derivative present

Degree: Power of highest derivative

Variable Separable: $\frac{dy}{dx} = f(x)g(y)$

Solution: $\int \frac{dy}{g(y)} = \int f(x)dx$

Homogeneous: $\frac{dy}{dx} = f\left(\frac{y}{x}\right)$

Solution: Substitute $y = vx$

Linear: $\frac{dy}{dx} + Py = Q$

Solution: $y \cdot e^{\int P dx} = \int Q \cdot e^{\int P dx} dx + C$

2.10 Vector Algebra

Vector: Quantity with magnitude and direction

Scalar: Quantity with only magnitude

Dot Product: $\vec{a} \cdot \vec{b} = |a||b| \cos \theta$

Definition: Scalar product of two vectors.

Cross Product: $\vec{a} \times \vec{b} = |a||b| \sin \theta \hat{n}$

Definition: Vector product perpendicular to both vectors.

Scalar Triple Product: $[\vec{a} \vec{b} \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c})$

Definition: Volume of parallelepiped.

2.11 Three Dimensional Geometry

Direction Cosines: $l = \cos \alpha, m = \cos \beta, n = \cos \gamma$

Definition: Cosines of angles with coordinate axes.

Equation of Line:

Vector: $\vec{r} = \vec{a} + \lambda \vec{b}$

Cartesian: $\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$

Definition: Different forms of line equation in 3D.

Equation of Plane:

Vector: $\vec{r} \cdot \vec{n} = d$

Cartesian: $ax + by + cz = d$

Definition: Different forms of plane equation.

Distance from Point to Plane: $d = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$

Definition: Perpendicular distance in 3D.

2.12 Probability

Conditional Probability: $P(A|B) = \frac{P(A \cap B)}{P(B)}$

Definition: Probability of A given that B has occurred.

Multiplication Theorem: $P(A \cap B) = P(A) \cdot P(B|A)$

Definition: Probability of both events occurring.

Independent Events: $P(A \cap B) = P(A)P(B)$

Definition: Occurrence of one doesn't affect the other.

Bayes' Theorem: $P(A|B) = \frac{P(B|A)P(A)}{P(B)}$

Definition: Relates conditional probabilities.

Random Variable: Variable whose values are numerical outcomes

Probability Distribution: List of probabilities for each value

Binomial Distribution: $P(X = r) = \binom{n}{r} p^r (1 - p)^{n-r}$

Definition: Probability of r successes in n independent trials.

Mean: $E(X) = \sum x_i p_i$

Variance: $Var(X) = E(X^2) - [E(X)]^2$

2.13 Linear Programming

Objective Function: $Z = ax + by$

Definition: Function to be maximized or minimized.

Constraints: $a_1x + b_1y \leq c_1$, $a_2x + b_2y \leq c_2$, etc.

Definition: Limitations on variables.

Feasible Region: Area satisfying all constraints

Definition: Set of all possible solutions.

Corner Point Method: Evaluate objective function at vertices

Definition: Optimal solution occurs at corner points.