

Stochastic Modelling and Random Processes

Problem Sheet 3

Yiming MA

December 15, 2020

Contents

1 Geometric Brownian Motion

1

1 Geometric Brownian Motion

Let $(X_t : t \geq 0)$ be a Brownian motion with constant drift on \mathbb{R} with generator

$$(\mathcal{L}f)(x) = \mu f'(x) + \frac{1}{2}\sigma^2 f''(x), \mu \in \mathbb{R}, \sigma > 0,$$

and initial condition $X_0 = 0$. **Geometric Brownian motion** is defined as

$$(Y_t : t \geq 0) \quad \text{with} \quad Y_t = e^{X_t}.$$

- (a) Show that $(Y_t : t \geq 0)$ is a diffusion process on $[0, \infty)$ and compute its generator. Write down the associated SDE and Fokker-Planck equation.

Sol. Notice that

$$\begin{aligned} \mathbb{E}[(\mathcal{L}_Y f)(Y_t)] &= \frac{d}{dt} \mathbb{E}[f(Y_t)] \\ &= \frac{d}{dt} \mathbb{E}[f(e^{X_t})]. \end{aligned} \tag{1}$$

Let $F = f \circ \exp$, then (1) becomes

$$\begin{aligned} \mathbb{E}[(\mathcal{L}_Y f)(Y_t)] &= \frac{d}{dt} \mathbb{E}[f(e^{X_t})] \\ &= \frac{d}{dt} \mathbb{E}[F(X_t)] \\ &= \mathbb{E}[(\mathcal{L}_X F)(X_t)]. \end{aligned} \tag{2}$$

Since (2) holds for all $f \in C^1\mathbb{R}$, we know

$$\begin{aligned}
(\mathcal{L}_Y f)(Y_t) &= (\mathcal{L}_X F)(X_t) \\
&= \mu \frac{d}{dx} f(e^{X_t}) + \frac{1}{2} \sigma^2 \frac{d^2}{dx^2} f(e^{X_t}) \\
&= \mu f'(e^{X_t}) e^{X_t} + \frac{1}{2} \sigma^2 \frac{d}{dx} (f'(e^{X_t}) e^{X_t}) \\
&= \mu f'(e^{X_t}) e^{X_t} + \frac{1}{2} \sigma^2 (f''(e^{X_t}) e^{2X_t} + f'(e^{X_t}) e^{X_t}) \\
&= \mu f'(Y_t) Y_t + \frac{1}{2} \sigma^2 (f''(Y_t) Y_t^2 + f'(Y_t) Y_t) \\
&= (\mu + \frac{1}{2} \sigma^2) Y_t f'(Y_t) + \frac{1}{2} (\sigma Y_t)^2 f''(Y_t),
\end{aligned}$$

which shows $(Y_t : t \geq 0)$ is a diffusion process with the drift $(\mu + \frac{1}{2} \sigma^2)y$ and the diffusion σy .

$$\begin{aligned}
&\int_{\mathbb{R}} \frac{\partial}{\partial t} p_t(x, y) f(y) dy \\
&= \frac{\partial}{\partial t} \int_{\mathbb{R}} p_t(x, y) f(y) dy \\
&= \frac{\partial}{\partial t} \mathbb{E}[f(Y_t)] \\
&= \mathbb{E}[(\mathcal{L}_Y f)(Y_t)] \\
&= \mathbb{E} \left[(\mu + \frac{1}{2} \sigma^2) Y_t f'(Y_t) + \frac{1}{2} (\sigma Y_t)^2 f''(Y_t) \right] \\
&= \int_{\mathbb{R}} \left[(\mu + \frac{1}{2} \sigma^2) y f'(y) + \frac{1}{2} (\sigma y)^2 f''(y) \right] p_t(x, y) dy \\
&= (\mu + \frac{1}{2} \sigma^2) \int_{\mathbb{R}} y f'(y) p_t(x, y) dy + \frac{1}{2} \sigma^2 \int_{\mathbb{R}} y^2 f''(y) p_t(x, y) dy \\
&= (\mu + \frac{1}{2} \sigma^2) \int_{\mathbb{R}} y p_t(x, y) df(y) + \frac{1}{2} \sigma^2 \int_{\mathbb{R}} y^2 p_t(x, y) df'(y) \\
&= (\mu + \frac{1}{2} \sigma^2) \left\{ y p_t(x, y) f(y) \Big|_{y=-\infty}^{y=\infty} - \int_{\mathbb{R}} f(y) \left[p_t(x, y) + y \frac{\partial}{\partial y} p_t(x, y) \right] dy \right\} \\
&\quad + \frac{1}{2} \sigma^2 \left\{ y^2 p_t(x, y) f'(y) \Big|_{y=-\infty}^{y=\infty} - \int_{\mathbb{R}} f'(y) \left[2y p_t(x, y) + y^2 \frac{\partial}{\partial y} p_t(x, y) \right] dy \right\} \\
&= -(\mu + \frac{1}{2} \sigma^2) \int_{\mathbb{R}} f(y) \left[p_t(x, y) + y \frac{\partial}{\partial y} p_t(x, y) \right] dy - \frac{1}{2} \sigma^2 \int_{\mathbb{R}} f'(y) \left[2y p_t(x, y) + y^2 \frac{\partial}{\partial y} p_t(x, y) \right] dy \\
&= -(\mu + \frac{1}{2} \sigma^2) \int_{\mathbb{R}} f(y) \left[p_t(x, y) + y \frac{\partial}{\partial y} p_t(x, y) \right] dy - \frac{1}{2} \sigma^2 \int_{\mathbb{R}} 2y p_t(x, y) + y^2 \frac{\partial}{\partial y} p_t(x, y) df(y) \\
&= -(\mu + \frac{1}{2} \sigma^2) \int_{\mathbb{R}} f(y) \left[p_t(x, y) + y \frac{\partial}{\partial y} p_t(x, y) \right] dy - \frac{1}{2} \sigma^2 \left\{ [2y p_t(x, y) + y^2 \frac{\partial}{\partial y} p_t(x, y)] f(y) \Big|_{y=-\infty}^{y=\infty} \right. \\
&\quad \left. - \int_{\mathbb{R}} f(y) [2p_t(x, y) + 4y \frac{\partial}{\partial y} p_t(x, y) + y^2 \frac{\partial^2}{\partial y^2} p_t(x, y)] dy \right\} \\
&= \int_{\mathbb{R}} f(y) \left[(\frac{1}{2} \sigma^2 - \mu) p_t(x, y) + (\frac{3}{2} \sigma^2 - \mu) y \frac{\partial}{\partial y} p_t(x, y) + \frac{1}{2} \sigma^2 y^2 \frac{\partial^2}{\partial y^2} p_t(x, y) \right] dy,
\end{aligned}$$

so the Fokker-Planck equation is

$$\frac{\partial}{\partial t} p_t(x, y) = \left(\frac{1}{2}\sigma^2 - \mu\right)p_t(x, y) + \left(\frac{3}{2}\sigma^2 - \mu\right)y \frac{\partial}{\partial y} p_t(x, y) + \frac{1}{2}\sigma^2 y^2 \frac{\partial^2}{\partial y^2} p_t(x, y).$$

An easier way is to use the conclusion for a diffusion process directly:

$$\begin{aligned} \frac{\partial}{\partial t} p_t(x, y) &= -\frac{\partial}{\partial y} \left[\left(\mu + \frac{1}{2}\sigma^2\right) y p_t \right] + \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2 y^2 p_t) \\ &= -\left(\mu + \frac{1}{2}\sigma^2\right)(p_t + y \frac{\partial}{\partial y} p_t(x, y)) + \frac{1}{2}\sigma^2 \frac{\partial}{\partial y} \left(2y p_t + y^2 \frac{\partial}{\partial y} p_t(x, y) \right) \\ &= \left(\frac{1}{2}\sigma^2 - \mu\right)p_t(x, y) + \left(\frac{3}{2}\sigma^2 - \mu\right)y \frac{\partial}{\partial y} p_t(x, y) + \frac{1}{2}\sigma^2 y^2 \frac{\partial^2}{\partial y^2} p_t(x, y). \end{aligned}$$

The associated SDE is

$$dY_t = \left(\mu + \frac{1}{2}\sigma^2\right)Y_t dt + \sigma Y_t dB_t.$$

- (b) Use the evolution equation of expectation values of test functions $f : \mathbb{R} \rightarrow \mathbb{R}$

$$\frac{d}{dt} \mathbb{E}[f(Y_t)] = \mathbb{E}[\mathcal{L}f(Y_t)],$$

to derive ODEs for the mean $m(t) := \mathbb{E}[Y_t]$ and the second moment $m_2(t) := \mathbb{E}[Y_t^2]$. (No need to solve the ODEs.)

Sol.

- (c) Under which conditions on μ and σ^2 is $(Y_t : t \geq 0)$ a martingale?

What is the asymptotic behaviour of the variance $v(t) = m_2(t) - m(t)^2$ in that case?

Sol.

- (d) Show that δ_0 is the unique stationary distribution of the process on the state space $[0, \infty)$. Under which conditions on μ and σ^2 does the process with $Y_0 = 1$ converge to the stationary distribution?

Under which conditions on μ and σ^2 is the process ergodic? Justify your answer.

Sol.

- (e) For $\sigma^2 = 1$ choose $\mu = -1/2$ and two other values $\mu < -1/2$ and $\mu > 1/2$. Simulate and plot a sample path of the process with $Y_0 = 1$ up to time $t = 10$, by numerically integrating the corresponding SDE with time steps $\Delta t = 0.1$ and 0.01 .

Sol.