

# 作业 08

陈昱全 SA18234049

## 题 1

$A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ , find eigenvalue  $\lambda$ , eigenstates  $|\lambda\rangle$  of  $A$ . With  $\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ , show

$$\sigma_z \otimes A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & -1 \end{pmatrix},$$

find eigenvalue and eigenvectors of  $\sigma_z \otimes A$ , show the relation with  $\pm\lambda$ , and  $|0\rangle|\lambda\rangle$ ,  $|1\rangle|\lambda\rangle$ .

解. 已知  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ , 则本征值和本征态为

```
In[ ]:= A = {{1, 1}, {1, 1}};
```

```
In[ ]:= Grid[Insert[Transpose[{Eigenvalues[A], Normalize /@  
[格子] [插入] [转置] [特征值] [正规化]  
Eigenvalues[A]}], {"Eigenvalue", "Eigenvector"}, 1], Frame -> All]  
[特征向量] [边框] [全部]
```

Eigenvalue	Eigenvector
2	$\left\{ \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\}$
0	$\left\{ -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\}$

又  $\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ , 二者的直积  $\sigma_z \otimes A$  为

```
In[ ]:= KroneckerProduct[PauliMatrix[3], A] // TraditionalForm  
[克罗内克积] [泡利自旋矩阵] [传统格式]
```

```
Out[ ]:= TraditionalForm=
```

$$\begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & -1 \end{pmatrix}$$

同理，可以求得  $\sigma_z \otimes A$  的本征值和本征态，

```
In[ ]:= ZA = KroneckerProduct[PauliMatrix[3], A];
          [克罗内克积] [泡利自旋矩阵]

Grid[Insert[Transpose[{Eigenvalues[ZA], Normalize /@
          [格子] [插入] [转置] [特征值] [正规化]
          Eigenvectors[ZA]}], {"Eigenvalue", "Eigenvector"}, 1], Frame -> All]
          [特征向量] [边框] [全部]
```

Out[ ]:=

Eigenvalue	Eigenvector
-2	$\{0, 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\}$
2	$\{\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0\}$
0	$\{0, 0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\}$
0	$\{-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0\}$

易知， $\sigma_z \otimes A$  的本征值和本征态有如下关系：

本征值	本征态
$\lambda$	$ 0\rangle \lambda\rangle$
$-\lambda$	$ 1\rangle \lambda\rangle$

□

## 题 2

“w-state”, for N-spin- $\frac{1}{2}$  particles, one can construct

$$|w_N\rangle \equiv \frac{1}{\sqrt{N}}(|10\dots 0\rangle + |010\dots 0\rangle + \dots + |0\dots 01\rangle)$$

with all the permutation of one of the particles at state  $|1\rangle$  and the other particles at  $|0\rangle$  state. What is the probability of measuring  $|1\rangle$  state for the first particle?

If we measure  $|0\rangle$  for the first particle, find the relation of the remaining state and  $|w_{N-1}\rangle$ .

## 题 3

Evolution of coupled spin- $\frac{1}{2}$  system.

$$H = \Omega(\sigma_z \otimes I + I \otimes \sigma_z), \quad |\psi(t=0)\rangle = \frac{1}{2}(|01\rangle + |10\rangle),$$

find  $|\psi(t)\rangle$ .

Hint: find the eigenstates and eigenvalues of  $H$  first