

a
Covariate tracking (hypothesis 2)
Complete VR
No national VR
Mean R² across causes
0.40
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
+0.131
(0.076 to 0.192)
+0.092
(0.057 to 0.154)
0.28
0.26
0.15
0.17
Deaths
(112 causes)
Prevalence
(110 causes)
b
Covariate R², cause by cause
95 of 112 causes
above the diagonal
R², countries without national VR
R², countries with complete VR
0.0
0.1
0.2
0.3
0.4
0.5
0.6
c
All 112 causes
Complete VR
No national VR
Cardiovascular
Neoplasms
Chronic respiratory
Digestive
Neuro & mental
Diabetes & kidney
Infectious & maternal
Other NCDs
0.1
0.2
0.3
0.4
0.5
0.6
R² of 5-year change in log death rate
on changes in nine covariates
d
Partial R²
SDI 0.015 0.052
Median age 0.013 0.029
Smoking 0.017 0.026
High SBP 0.009 0.021
Unsafe water 0.008 0.019
High BMI 0.009 0.019
Solid fuel 0.026 0.013
High FPG 0.016 0.012
PM2.5 0.009 0.006
Full
None
National VR
e
CI width
EAPC CI width, relative to cause median
10
3
1
0.3
0.1
Full
Part
None
National VR
f
Median, by category
Cardiovascular 1.31 1.09 0.65
Neoplasms 1.21 1.07 0.74
Chronic respiratory 1.54 0.97 0.61
Digestive 1.44 0.97 0.63
Neuro & mental 1.22 1.10 0.78
Diabetes & kidney 1.56 0.99 0.57
Infectious & maternal 1.34 0.98 0.76
Other NCDs 1.53 0.97 0.62
Full
Part
None
National VR