

a
GBD minus raw VR trend (z statistic), 29 causes × 78 countries with complete VR
% differ
100
50
0
per cause
Cardiovascular
Neoplasms
Chronic respiratory
Digestive
Neuro & mental
Diabetes & kidney
Infectious & maternal
Countries ordered by population (small → large)
GBD lower
0
GBD higher
0
50
100
% of countries differing
b
GBD v raw trends
n=2232; r=0.68
dashed: line of identity
EAPC, GBD (%/year)
EAPC, raw VR (%/year)
c
Disagreement by precision
Differ
Agree
% of series
66
61
57
52
53
55
52
52
47
33
Decile of raw-trend SE (1 = most precise)
d
By country size
% of series differing
Population, 2019 (millions)
e
50 exposure–outcome pairs
raw P<0.05 (19)
raw P≥0.05
Association, GBD
Association, raw VR
red: median ratio 0.63
f
|GBD| / |raw| association for each pair (bold: raw P<0.05; grey cell: pair not analysed)
SDI
Smoking
High BMI
High SBP
High FPG
Alcohol
PM2.5
Solid fuel
IHD
Stroke
COPD
Lung ca
Diabetes
CKD
Liver ca
Colorectal ca
Stomach ca
LRI
Dementia
CLD
Breast ca
Pancreatic ca
Oesophageal ca
Bladder ca
Kidney ca
HHD
AF
≥2
1.5
1
0.5
0