

1 Chapter 1

All:1.1,1.2,1.7,1.14 and 6 examples

1.1:

$$\begin{aligned}x &= \tan\theta d \quad \dot{x} = \frac{\dot{\theta}}{\cos^2\theta} d = \frac{\omega}{\cos^2\theta} d \\ \cos\theta &= \frac{d}{\sqrt{d^2 + x^2}} \Rightarrow \frac{\omega(d^2 + x^2)}{d} \\ \ddot{x} &= \frac{2x\dot{x}\omega}{d} = \frac{2x\omega}{d} \frac{\omega(d^2 + x^2)}{d} = \frac{2x\omega^2(d^2 + x^2)}{d^2}\end{aligned}$$

1.2:

$$\begin{aligned}\begin{cases} x = \sin\theta b \\ y = \cos\theta d \end{cases} \\ \frac{d}{dt} \cos\theta(b+d) &= -c \\ \Rightarrow \dot{\theta} &= \frac{c}{\sin\theta(b+d)} \\ \begin{cases} \dot{x} = \frac{cb\cos\theta}{\sin\theta(b+d)} \\ \dot{y} = -\frac{dc}{(b+d)} \end{cases} \\ \begin{cases} \ddot{x} = -\frac{c^2 b}{\sin^3\theta(b+d)^2} \\ \ddot{y} = 0 \end{cases}\end{aligned}$$

1.7:

$$\begin{aligned}\vec{v} &= v_x \vec{i} + v_y \vec{j} \quad v_x^2 + v_y^2 = c \\ \vec{a} &= \dot{v}_x \vec{i} + \dot{v}_y \vec{j} \\ \vec{a} \cdot \vec{v} &= v_x \dot{v}_x + v_y \dot{v}_y = \frac{1}{2}(\dot{v}_x^2 + \dot{v}_y^2) = \frac{1}{2}\dot{c} = 0\end{aligned}$$

1.14:

$$\begin{aligned}
& \begin{cases} m\ddot{x} = -mk\dot{x} \\ m\ddot{y} = -mk\dot{y} - mg \end{cases} \\
& \begin{cases} \dot{x} = C_1 e^{-kt} \\ \dot{y} = C_2 e^{-kt} - \frac{g}{k} \end{cases} \\
& t = 0 \quad C_1 = v_{x0}; C_2 = \frac{g}{k} + v_{y0} \\
& -\tan\alpha = \frac{(\frac{g}{k} + v_{y0})e^{-kt} - \frac{g}{k}}{v_{x0}e^{-kt}} \\
& -\tan\alpha = (\frac{g}{kv_{x0}} + \tan\alpha) - \frac{g}{kv_{x0}}e^{kt} \\
& \Rightarrow e^{kt} = \frac{kv_{x0}}{g}(\frac{g}{kv_{x0}} + 2\tan\alpha) \\
& t = \frac{1}{k}\ln[\frac{kv_{x0}}{g}(\frac{g}{kv_{x0}} + 2\tan\alpha)] = \frac{1}{k}\ln[1 + 2\frac{k}{g}\sin\alpha v_0]
\end{aligned}$$

exp1:

$$\begin{aligned}
v &= \begin{cases} v_x = v_0 \\ v_y = gt \end{cases} \\
a_n &= \frac{v^2}{\rho} \Rightarrow \rho = \frac{v^2}{a_n} \\
a_n &= \vec{a} \cdot \vec{n} = \sqrt{a^2 - (a \cdot v \frac{1}{|v|})^2} \\
\vec{a} &= g\vec{j}; \vec{a} \cdot \vec{v} = g^2 t \\
a_n &= \sqrt{g^2 - \frac{g^4 t^2}{v_0^2 + g^2 t^2}} = \sqrt{\frac{v_0^2 g^2}{v_0^2 + g^2 t^2}} = \frac{v_0 g}{\sqrt{v_0^2 + g^2 t^2}} \\
\rho &= \frac{(v_0^2 + g^2 t^2)^{\frac{3}{2}}}{v_0 g}
\end{aligned}$$

exp2:

$$\begin{aligned}
m\ddot{y} &= -mg - mk\dot{y} \\
\ddot{y} + k\dot{y} &= -g \Rightarrow y = A + Be^{-kt} - \frac{gt}{k} \\
\dot{y}|_{t=0} &= 0 \Rightarrow \dot{y} = \frac{g}{k}e^{-kt} - \frac{g}{k}
\end{aligned}$$

exp3:

$$l = r \times p = 2 \times 0.5 \times 4 = 1 = v_B \times 0.5 \times 2 \\ \Rightarrow v_B = 1$$

exp4:

$$mg = kx \Rightarrow x = \frac{mg}{k} \\ \frac{1}{2}mv^2 - mgx + \frac{1}{2}kx^2 = 0 \\ \Rightarrow \frac{1}{2}mv^2 = \frac{m^2g^2}{k} - \frac{1}{2}k\frac{m^2g^2}{k^2} = \frac{m^2g^2}{2k} \\ v = \sqrt{\frac{m}{k}}g$$

exp5:

$$V = - \int (ax + by^2)dx + (az + 2bxy)dy + (ay + bz^2)dz \\ = - \frac{1}{2}ax^2 - \frac{1}{3}bz^3 - \int by^2dx + azdy + bxdy^2 + aydz \\ = - \frac{1}{2}ax^2 - \frac{1}{3}bz^3 - by^2x - azy + C \\ C = 0$$

exp6:

$$\frac{d^2x}{dt^2} = 2 + 6x^2 \\ \int_0^x adx = 2x + 2x^3 = \int_0^x \frac{d^2x}{dt^2}dx = \int_0^x vdv = \frac{1}{2}v^2 - \frac{1}{2}v_0^2 \\ v = \sqrt{100 + 4x + 4x^3}$$

2 Chapter 2

2.8,2.13,2 examples

2.8:

$$\vec{r} = r\vec{e}_r; \vec{v} = \dot{r}\vec{e}_r + r\omega\vec{e}_\theta;$$

$$\vec{a} = \ddot{r}\vec{e}_r + \dot{r}\omega\vec{e}_\theta + (\dot{r}\omega + r\dot{\omega})\vec{e}_\theta - r\omega^2\vec{e}_r = (\ddot{r} - r\omega^2)\vec{e}_r + (2\dot{r}\omega + r\dot{\omega})\vec{e}_\theta$$

$$r \times p = h \Rightarrow r^2\omega = \frac{h}{m}$$

$$\dot{r} = \frac{dr}{dt} = \frac{dr}{d\theta}\omega = \frac{dr}{d\theta}\frac{h}{mr^2} = -\frac{h}{m}\frac{d\frac{1}{r}}{d\theta}$$

$$r = ae^{b\theta} \Rightarrow \dot{r} = -\frac{h}{m}\frac{1}{a}(-b)e^{-b\theta} = \frac{bh}{m}\frac{1}{r}$$

$$\ddot{r} = -\frac{hb}{m}\frac{1}{r^2}\left(\frac{hb}{m}\frac{1}{r}\right) = -\frac{h^2b^2}{m^2}\frac{1}{r^3}$$

$$\ddot{r} - r\omega^2 = \frac{F}{m} = -\frac{k}{mr^3} = -\frac{h^2b^2}{m^2}\frac{1}{r^3} - r\frac{h^2}{m^2r^4}$$

$$\Rightarrow k = \frac{h^2b^2}{m} + \frac{h^2}{m} \Rightarrow h = \sqrt{\frac{km}{b^2 + 1}}$$

$$E = T + V = \frac{1}{2}m(\dot{r}^2 + r^2\omega^2) - \frac{k}{2r^2} = \frac{1}{2}m\left(\frac{h^2b^2}{m^2r^2} + \frac{h^2}{m^2r^2}\right) - \frac{k}{2r^2}$$

$$\Rightarrow \frac{1}{2r^2}\left(\frac{h^2b^2}{m} + \frac{h^2}{m} - k\right) = 0$$