

Chapitre 1. Random Variables

Prof. REMITA Mohamed Riad

National Higher School of Artificial Intelligence.

2022-2023

Definitions and notations

Example

A coin is tossed twice.

Definitions and notations

Example

A coin is tossed twice. The possible results are $\{PP, PF, FP \text{ et } FF\}$. We define variable X representing the number of tails P obtained.

Definitions and notations

Example

A coin is tossed twice. The possible results are $\{PP, PF, FP \text{ et } FF\}$. We define variable X representing the number of tails P obtained. Then the values of X are $\{0, 1 \text{ and } 2\}$.

Definitions and notations

Example

A coin is tossed twice. The possible results are $\{PP, PF, FP \text{ et } FF\}$. We define variable X representing the number of tails P obtained. Then the values of X are $\{0, 1 \text{ and } 2\}$.

Example

A die is rolled until a 6 is rolled.

Definitions and notations

Example

A coin is tossed twice. The possible results are $\{PP, PF, FP \text{ et } FF\}$. We define variable X representing the number of tails P obtained. Then the values of X are $\{0, 1 \text{ and } 2\}$.

Example

A die is rolled until a 6 is rolled. The possible outcomes are $\{6, (1, 6), (2, 6), \dots, (5, 6), (1, 1, 6), \dots, (5, 5, 6), \dots\}$. We define a variable X representing the number of throws needed until a 6 is obtained.

Definitions and notations

Example

A coin is tossed twice. The possible results are $\{PP, PF, FP \text{ et } FF\}$. We define variable X representing the number of tails P obtained. Then the values of X are $\{0, 1 \text{ and } 2\}$.

Example

A die is rolled until a 6 is rolled. The possible outcomes are $\{6, (1, 6), (2, 6), \dots, (5, 6), (1, 1, 6), \dots, (5, 5, 6), \dots\}$. We define a variable X representing the number of throws needed until a 6 is obtained. Then the values of X are $\{1, 2, 3, \dots\} = \mathbb{N}^*$.

Definitions and notations

Example

A coin is tossed twice. The possible results are $\{PP, PF, FP \text{ et } FF\}$. We define variable X representing the number of tails P obtained. Then the values of X are $\{0, 1 \text{ and } 2\}$.

Example

A die is rolled until a 6 is rolled. The possible outcomes are $\{6, (1, 6), (2, 6), \dots, (5, 6), (1, 1, 6), \dots, (5, 5, 6), \dots\}$. We define a variable X representing the number of throws needed until a 6 is obtained. Then the values of X are $\{1, 2, 3, \dots\} = \mathbb{N}^*$.

Example

The service life of a spare part can be represented by a r.r.v.

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

$$\forall B \in \mathcal{B}_{\mathbb{R}}, X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{A}.$$

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

$$\forall B \in \mathcal{B}_{\mathbb{R}}, X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{A}.$$

B can be presented in several forms

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

$$\forall B \in \mathcal{B}_{\mathbb{R}}, X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{A}.$$

B can be presented in several forms

- If $B =]a, b] : X^{-1}(B) = \{a < X \leq b\}$

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

$$\forall B \in \mathcal{B}_{\mathbb{R}}, X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{A}.$$

B can be presented in several forms

- If $B =]a, b] : X^{-1}(B) = \{a < X \leq b\}$
- If $B = \{a\} : X^{-1}(B) = \{X = a\}$

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

$$\forall B \in \mathcal{B}_{\mathbb{R}}, X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{A}.$$

B can be presented in several forms

- If $B =]a, b] : X^{-1}(B) = \{a < X \leq b\}$
- If $B = \{a\} : X^{-1}(B) = \{X = a\}$
- If $B = [a, +\infty[: X^{-1}(B) = \{X \geq a\}$

Definitions and notations

We have a probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ and a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Definition

We call a real random variable, noted r.r.v. any application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ such that:

$$\forall B \in \mathcal{B}_{\mathbb{R}}, X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{A}.$$

B can be presented in several forms

- If $B =]a, b] : X^{-1}(B) = \{a < X \leq b\}$
- If $B = \{a\} : X^{-1}(B) = \{X = a\}$
- If $B = [a, +\infty[: X^{-1}(B) = \{X \geq a\}$
- If $B =]-\infty, a] : X^{-1}(B) = \{X \leq a\}$

Definitions and notations

Since the Borel σ -algebra is generated by all the intervals $] -\infty, x]$, then a random variable can be defined by the following definition:

Definitions and notations

Since the Borel σ -algebra is generated by all the intervals $] -\infty, x]$, then a random variable can be defined by the following definition:

Definition

The application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ is a real random variable if for all $x \in \mathbb{R}$ and $\forall B \in \mathcal{B}_{\mathbb{R}}$ the subset $A_x = X^{-1} (]-\infty, x]) = \{\omega \in \Omega : X(\omega) \leq x\} \in \mathcal{A}$.

Definitions and notations

Since the Borel σ -algebra is generated by all the intervals $] -\infty, x]$, then a random variable can be defined by the following definition:

Definition

The application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ is a real random variable if for all $x \in \mathbb{R}$ and $\forall B \in \mathcal{B}_{\mathbb{R}}$ the subset $A_x = X^{-1} (]-\infty, x]) = \{\omega \in \Omega : X(\omega) \leq x\} \in \mathcal{A}$.

Example

We throw two coins and let X be the number of tails obtained. We know that $\Omega = \{(F, F); (F, P); (P, F); (P, P)\}$ and the values of X are $\{0, 1, 2\}$.

Definitions and notations

Since the Borel σ -algebra is generated by all the intervals $] -\infty, x]$, then a random variable can be defined by the following definition:

Definition

The application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ is a real random variable if for all $x \in \mathbb{R}$ and $\forall B \in \mathcal{B}_{\mathbb{R}}$ the subset $A_x = X^{-1}(] -\infty, x]) = \{\omega \in \Omega : X(\omega) \leq x\} \in \mathcal{A}$.

Example

We throw two coins and let X be the number of tails obtained. We know that $\Omega = \{(F, F); (F, P); (P, F); (P, P)\}$ and the values of X are $\{0, 1, 2\}$.

1. Show that X is a random variable on Ω endowed with the algebra $\mathcal{P}(\Omega)$.

Definitions and notations

Since the Borel σ -algebra is generated by all the intervals $] -\infty, x]$, then a random variable can be defined by the following definition:

Definition

The application X of (Ω, \mathcal{A}) in $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ is a real random variable if for all $x \in \mathbb{R}$ and $\forall B \in \mathcal{B}_{\mathbb{R}}$ the subset $A_x = X^{-1} (]-\infty, x]) = \{\omega \in \Omega : X(\omega) \leq x\} \in \mathcal{A}$.

Example

We throw two coins and let X be the number of tails obtained. We know that $\Omega = \{(F, F); (F, P); (P, F); (P, P)\}$ and the values of X are $\{0, 1, 2\}$.

1. Show that X is a random variable on Ω endowed with the algebra $\mathcal{P}(\Omega)$.
2. Show that X is not a random variable on Ω endowed with the algebra $\mathcal{A}_1 = \{\Omega, \emptyset, \{(F, F)\}, \{(F, F); (F, P); (P, F)\}\}$.

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A .

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A . We can show easily that $I_A(\cdot)$ is a random variable for the algebra $\mathcal{A}_{I_A(\cdot)} = \{\Omega, \emptyset, A, \bar{A}\}$.

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A . We can show easily that $I_A(\cdot)$ is a random variable for the algebra $\mathcal{A}_{I_A(\cdot)} = \{\Omega, \emptyset, A, \bar{A}\}$.

Properties. The indicator function $I_A(\cdot)$ satisfies the following properties:

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A . We can show easily that $I_A(\cdot)$ is a random variable for the algebra $\mathcal{A}_{I_A(\cdot)} = \{\Omega, \emptyset, A, \bar{A}\}$.

Properties. The indicator function $I_A(\cdot)$ satisfies the following properties:

1. $I_A(\omega) = 1 - I_{\bar{A}}(\omega), \forall A \in \mathcal{A},$

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A . We can show easily that $I_A(\cdot)$ is a random variable for the algebra $\mathcal{A}_{I_A(\cdot)} = \{\Omega, \emptyset, A, \bar{A}\}$.

Properties. The indicator function $I_A(\cdot)$ satisfies the following properties:

1. $I_A(\omega) = 1 - I_{\bar{A}}(\omega), \forall A \in \mathcal{A},$
2. $I_{\cap A_i}(\omega) = \prod_i I_{A_i}(\omega), \forall A_i \in \mathcal{A},$

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A . We can show easily that $I_A(\cdot)$ is a random variable for the algebra $\mathcal{A}_{I_A(\cdot)} = \{\Omega, \emptyset, A, \bar{A}\}$.

Properties. The indicator function $I_A(\cdot)$ satisfies the following properties:

1. $I_A(\omega) = 1 - I_{\bar{A}}(\omega), \forall A \in \mathcal{A},$
2. $I_{\cap A_i}(\omega) = \prod_i I_{A_i}(\omega), \forall A_i \in \mathcal{A},$
3. $\mathbb{P}(I_A(\omega) = 1) = \mathbb{P}(A), \forall A \in \mathcal{A},$

Definitions and notations

Let Ω be the space of trials associated to a Bernoulli random experiment and let $I_A(\cdot)$ be the function from Ω to $\{0, 1\}$ defined by

$$I_A(\cdot) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}.$$

$I_A(\cdot)$ is called indicator function (or Dirac measure) of the event A . We can show easily that $I_A(\cdot)$ is a random variable for the algebra $\mathcal{A}_{I_A(\cdot)} = \{\Omega, \emptyset, A, \bar{A}\}$.

Properties. The indicator function $I_A(\cdot)$ satisfies the following properties:

1. $I_A(\omega) = 1 - I_{\bar{A}}(\omega), \forall A \in \mathcal{A},$
2. $I_{\cap A_i}(\omega) = \prod_i I_{A_i}(\omega), \forall A_i \in \mathcal{A},$
3. $\mathbb{P}(I_A(\omega) = 1) = \mathbb{P}(A), \forall A \in \mathcal{A},$
4. $\mathbb{P}(I_A(\omega) = 0) = 1 - \mathbb{P}(A) = \mathbb{P}(\bar{A}), \forall A \in \mathcal{A}.$

Induced probability

Theorem

Let X be a r.r.v. defined on probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ with values in a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$. The application \mathbb{P}_X of $\mathcal{B}_{\mathbb{R}}$ in \mathbb{R} defined by $\mathbb{P}_X(B) = \mathbb{P}(X^{-1}(B))$, is a probability on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Induced probability

Theorem

Let X be a r.r.v. defined on probabilized space $(\Omega, \mathcal{A}, \mathbb{P})$ with values in a probabilizable space $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$. The application \mathbb{P}_X of $\mathcal{B}_{\mathbb{R}}$ in \mathbb{R} defined by $\mathbb{P}_X(B) = \mathbb{P}(X^{-1}(B))$, is a probability on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$.

Remark

The definition is due to the existence of \mathbb{P} on (Ω, \mathcal{T}) , hence the notion of induced probability.

Induced probability

Proof.

It is obvious that \mathbb{P} is an application with values in $[0, 1]$.

Induced probability

Proof.

It is obvious that \mathbb{P} is an application with values in $[0, 1]$.

Moreover \mathbb{P} verifies these conditions:

$$\mathbb{P}_X(\mathbb{R}) = \mathbb{P}(X^{-1}(\mathbb{R})) = \mathbb{P}(\Omega) = 1$$

Let $(B_i)_{i \geq 1}$ be two by two incompatible borelean sequences.

Induced probability

Proof.

It is obvious that \mathbb{P} is an application with values in $[0, 1]$.

Moreover \mathbb{P} verifies these conditions:

$$\mathbb{P}_X(\mathbb{R}) = \mathbb{P}(X^{-1}(\mathbb{R})) = \mathbb{P}(\Omega) = 1$$

Let $(B_i)_{i \geq 1}$ be two by two incompatible borelean sequences. Then

$$\begin{aligned} \mathbb{P}_X\left(\bigcup_{i \geq 1} B_i\right) &= \mathbb{P}\left(X^{-1}\left(\bigcup_{i \geq 1} B_i\right)\right) = \mathbb{P}\left(\bigcup_{i \geq 1} X^{-1}(B_i)\right) \\ &= \sum_{i \geq 1} \mathbb{P}(X^{-1}(B_i)) = \sum_{i \geq 1} \mathbb{P}_X(B_i), \end{aligned}$$

Induced probability

Proof.

It is obvious that \mathbb{P} is an application with values in $[0, 1]$.

Moreover \mathbb{P} verifies these conditions:

$$\mathbb{P}_X(\mathbb{R}) = \mathbb{P}(X^{-1}(\mathbb{R})) = \mathbb{P}(\Omega) = 1$$

Let $(B_i)_{i \geq 1}$ be two by two incompatible borelean sequences. Then

$$\begin{aligned} \mathbb{P}_X\left(\bigcup_{i \geq 1} B_i\right) &= \mathbb{P}\left(X^{-1}\left(\bigcup_{i \geq 1} B_i\right)\right) = \mathbb{P}\left(\bigcup_{i \geq 1} X^{-1}(B_i)\right) \\ &= \sum_{i \geq 1} \mathbb{P}(X^{-1}(B_i)) = \sum_{i \geq 1} \mathbb{P}_X(B_i), \end{aligned}$$

noting that $X^{-1}(B_i)$ and $X^{-1}(B_j)$ are incompatible $\forall i \neq j$. □

Cumulative distribution function of a random variable

Definition

The cumulative distribution function of a r.r.v. is the function F or F_X defined by:

$$F(x) = F_X(x) = \mathbb{P}(X \leq x).$$

Cumulative distribution function of a random variable

Definition

The cumulative distribution function of a r.r.v. is the function F or F_X defined by:

$$F(x) = F_X(x) = \mathbb{P}(X \leq x).$$

Properties of a cumulative distribution function

Cumulative distribution function of a random variable

Definition

The cumulative distribution function of a r.r.v. is the function F or F_X defined by:

$$F(x) = F_X(x) = \mathbb{P}(X \leq x).$$

Properties of a cumulative distribution function

Definition

A sequence of events $(A_n)_{n \geq 1}$ is increasing (resp. decreasing) if $A_n \subset A_{n+1}$ (resp. $A_{n+1} \subset A_n$) for all $n \geq 1$. $(A_n)_{n \geq 1}$ is said to be monotonic if it is increasing or decreasing.

Cumulative distribution function of a random variable

Definition

The cumulative distribution function of a r.r.v. is the function F or F_X defined by:

$$F(x) = F_X(x) = \mathbb{P}(X \leq x).$$

Properties of a cumulative distribution function

Definition

A sequence of events $(A_n)_{n \geq 1}$ is increasing (resp. decreasing) if $A_n \subset A_{n+1}$ (resp. $A_{n+1} \subset A_n$) for all $n \geq 1$. $(A_n)_{n \geq 1}$ is said to be monotonic if it is increasing or decreasing. In this case we put $\lim_{n \rightarrow \infty} A_n = \bigcup_{n \geq 1} A_n$ if it is increasing (resp. $\lim_{n \rightarrow \infty} A_n = \bigcap_{n \geq 1} A_n$ if it is decreasing).

Cumulative distribution function of a random variable

Remark

$\lim_{n \rightarrow \infty} A_n$ exists if and only if the sequence $(A_n)_{n \geq 1}$ is monotonic.

Cumulative distribution function of a random variable

Remark

$\lim_{n \rightarrow \infty} A_n$ exists if and only if the sequence $(A_n)_{n \geq 1}$ is monotonic.

Lemma

(Property of the continuity of \mathbb{P})

If $(A_n)_{n \geq 1}$ is a monotonic sequence of events, then we have:

$$\lim_{n \rightarrow \infty} \mathbb{P}(A_n) = \mathbb{P}\left(\lim_{n \rightarrow \infty} A_n\right).$$

Cumulative distribution function of a random variable

Theorem

If F is the cumulative distribution function of X then

Cumulative distribution function of a random variable

Theorem

If F is the cumulative distribution function of X then

1. $\forall x \in \mathbb{R} \ 0 \leq F(x) \leq 1;$

Cumulative distribution function of a random variable

Theorem

If F is the cumulative distribution function of X then

1. $\forall x \in \mathbb{R} \ 0 \leq F(x) \leq 1;$
2. F is an increasing function;

Cumulative distribution function of a random variable

Theorem

If F is the cumulative distribution function of X then

1. $\forall x \in \mathbb{R} \ 0 \leq F(x) \leq 1;$
2. F is an increasing function;
3. F is right continuous;

Cumulative distribution function of a random variable

Theorem

If F is the cumulative distribution function of X then

1. $\forall x \in \mathbb{R} \ 0 \leq F(x) \leq 1;$
2. F is an increasing function;
3. F is right continuous;
4. $\lim_{x \rightarrow \infty} F(x) = 1$ and $\lim_{x \rightarrow -\infty} F(x) = 0.$

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$.

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$. It follows that $\mathbb{P}(X^{-1}(] -\infty, x_1])) \leq \mathbb{P}(X^{-1}(] -\infty, x_2]))$ hence $F(x_1) \leq F(x_2)$.

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$. It follows that $\mathbb{P}(X^{-1}(] -\infty, x_1])) \leq \mathbb{P}(X^{-1}(] -\infty, x_2]))$ hence $F(x_1) \leq F(x_2)$.
3. Let us show that for any real sequence (ε_n) decreasing and converging to 0, $\lim_{n \rightarrow \infty} F(x + \varepsilon_n) = F(x)$.

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$. It follows that $\mathbb{P}(X^{-1}(] -\infty, x_1])) \leq \mathbb{P}(X^{-1}(] -\infty, x_2]))$ hence $F(x_1) \leq F(x_2)$.
3. Let us show that for any real sequence (ε_n) decreasing and converging to 0, $\lim_{n \rightarrow \infty} F(x + \varepsilon_n) = F(x)$. We set $A_n =]x, x + \varepsilon_n]$.

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$. It follows that $\mathbb{P}(X^{-1}(] -\infty, x_1])) \leq \mathbb{P}(X^{-1}(] -\infty, x_2]))$ hence $F(x_1) \leq F(x_2)$.
3. Let us show that for any real sequence (ε_n) decreasing and converging to 0, $\lim_{n \rightarrow \infty} F(x + \varepsilon_n) = F(x)$. We set $A_n =]x, x + \varepsilon_n]$. The (A_n) are decreasing and $\lim_{n \rightarrow \infty} A_n = \emptyset$, hence from the lemma $\lim_{n \rightarrow \infty} \mathbb{P}_X(A_n) = \mathbb{P}_X(\lim_{n \rightarrow \infty} A_n) = \mathbb{P}_X(\emptyset) = 0$.

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$. It follows that $\mathbb{P}(X^{-1}(] -\infty, x_1])) \leq \mathbb{P}(X^{-1}(] -\infty, x_2]))$ hence $F(x_1) \leq F(x_2)$.
3. Let us show that for any real sequence (ε_n) decreasing and converging to 0, $\lim_{n \rightarrow \infty} F(x + \varepsilon_n) = F(x)$. We set $A_n =]x, x + \varepsilon_n]$. The (A_n) are decreasing and $\lim_{n \rightarrow \infty} A_n = \emptyset$, hence from the lemma $\lim_{n \rightarrow \infty} \mathbb{P}_X(A_n) = \mathbb{P}_X(\lim_{n \rightarrow \infty} A_n) = \mathbb{P}_X(\emptyset) = 0$. Since

$$\begin{aligned}\mathbb{P}_X(A_n) &= \mathbb{P}(x < X \leq x + \varepsilon_n) = \mathbb{P}(X \leq x + \varepsilon_n) - \mathbb{P}(X \leq x) \\ &= F(x + \varepsilon_n) - F(x),\end{aligned}$$

Cumulative distribution function of a random variable

Proof.

1. Obvious because $F(X) = \mathbb{P}(X \leq x)$ so $0 \leq F(X) \leq 1$.
2. Suppose that $x_1 \leq x_2$, hence $] -\infty, x_1] \subset] -\infty, x_2]$ and $X^{-1}(] -\infty, x_1]) \subset X^{-1}(] -\infty, x_2])$. It follows that $\mathbb{P}(X^{-1}(] -\infty, x_1])) \leq \mathbb{P}(X^{-1}(] -\infty, x_2]))$ hence $F(x_1) \leq F(x_2)$.
3. Let us show that for any real sequence (ε_n) decreasing and converging to 0, $\lim_{n \rightarrow \infty} F(x + \varepsilon_n) = F(x)$. We set $A_n =]x, x + \varepsilon_n]$. The (A_n) are decreasing and $\lim_{n \rightarrow \infty} A_n = \emptyset$, hence from the lemma

$\lim_{n \rightarrow \infty} \mathbb{P}_X(A_n) = \mathbb{P}_X(\lim_{n \rightarrow \infty} A_n) = \mathbb{P}_X(\emptyset) = 0$. Since

$$\begin{aligned}\mathbb{P}_X(A_n) &= \mathbb{P}(x < X \leq x + \varepsilon_n) = \mathbb{P}(X \leq x + \varepsilon_n) - \mathbb{P}(X \leq x) \\ &= F(x + \varepsilon_n) - F(x),\end{aligned}$$

then $\lim_{n \rightarrow \infty} F(x + \varepsilon_n) = F(x)$.

Cumulative distribution function of a random variable

Proof.

4. Let $B_n =]-\infty, x_n]$ where the x_n is decreasing and converging to $-\infty$.

Cumulative distribution function of a random variable

Proof.

4. Let $B_n =]-\infty, x_n]$ where the x_n is decreasing and converging to $-\infty$. We deduce that (B_n) is a decreasing sequence and $\lim B_n = \emptyset$, and according to the lemma

$$\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \mathbb{P}_X\left(\lim_{n \rightarrow \infty} B_n\right) = \mathbb{P}_X(\emptyset) = 0$$

Cumulative distribution function of a random variable

Proof.

4. Let $B_n =]-\infty, x_n]$ where the x_n is decreasing and converging to $-\infty$. We deduce that (B_n) is a decreasing sequence and $\lim B_n = \emptyset$, and according to the lemma

$$\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \mathbb{P}_X\left(\lim_{n \rightarrow \infty} B_n\right) = \mathbb{P}_X(\emptyset) = 0$$

or $\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \lim_{n \rightarrow \infty} \mathbb{P}_X(X \leq x_n) =$
 $\lim_{n \rightarrow \infty} F(x_n) = 0.$

Cumulative distribution function of a random variable

Proof.

4. Let $B_n =]-\infty, x_n]$ where the x_n is decreasing and converging to $-\infty$. We deduce that (B_n) is a decreasing sequence and $\lim B_n = \emptyset$, and according to the lemma

$$\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \mathbb{P}_X\left(\lim_{n \rightarrow \infty} B_n\right) = \mathbb{P}_X(\emptyset) = 0$$

or $\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \lim_{n \rightarrow \infty} \mathbb{P}_X(X \leq x_n) =$
 $\lim_{n \rightarrow \infty} F(x_n) = 0$.

Let us consider the sequence defined by $C_n =]-\infty, y_n]$ where (y_n) is an increasing real sequence such that $\lim_{n \rightarrow \infty} y_n = +\infty$.

Cumulative distribution function of a random variable

Proof.

4. Let $B_n =]-\infty, x_n]$ where the x_n is decreasing and converging to $-\infty$. We deduce that (B_n) is a decreasing sequence and $\lim B_n = \emptyset$, and according to the lemma

$$\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \mathbb{P}_X\left(\lim_{n \rightarrow \infty} B_n\right) = \mathbb{P}_X(\emptyset) = 0$$

or $\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \lim_{n \rightarrow \infty} \mathbb{P}_X(X \leq x_n) =$
 $\lim_{n \rightarrow \infty} F(x_n) = 0$.

Let us consider the sequence defined by $C_n =]-\infty, y_n]$ where (y_n) is an increasing real sequence such that $\lim_{n \rightarrow \infty} y_n = +\infty$. We deduce that the (C_n) are increasing and $\lim_{n \rightarrow \infty} C_n = \mathbb{R}$.

Cumulative distribution function of a random variable

Proof.

4. Let $B_n =]-\infty, x_n]$ where the x_n is decreasing and converging to $-\infty$. We deduce that (B_n) is a decreasing sequence and $\lim B_n = \emptyset$, and according to the lemma

$$\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \mathbb{P}_X\left(\lim_{n \rightarrow \infty} B_n\right) = \mathbb{P}_X(\emptyset) = 0$$

or $\lim_{n \rightarrow \infty} \mathbb{P}_X(B_n) = \lim_{n \rightarrow \infty} \mathbb{P}_X(X \leq x_n) =$
 $\lim_{n \rightarrow \infty} F(x_n) = 0$.

Let us consider the sequence defined by $C_n =]-\infty, y_n]$ where (y_n) is an increasing real sequence such that $\lim_{n \rightarrow \infty} y_n = +\infty$. We deduce that the (C_n) are increasing and $\lim_{n \rightarrow \infty} C_n = \mathbb{R}$.

We have $\lim_{n \rightarrow +\infty} F(y_n) = \lim_{n \rightarrow +\infty} \mathbb{P}(X \leq y_n) =$
 $\lim_{n \rightarrow +\infty} \mathbb{P}_X(C_n) = \mathbb{P}_X(\lim_{n \rightarrow +\infty} C_n) = \mathbb{P}_X(\mathbb{R}) = 1$.