

BlackHoleImages: A Wolfram Mathematica packet for analytical black hole imaging

David Podrápský¹ and Vojtěch Witzany¹

¹ Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 180 00 Praha 8, Czech Republic

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Black holes are astrophysical objects with gravitational fields so extreme that they bend even the trajectories of light. Specifically, in the geometrical optics limit, the paths of particles and fields moving at the speed of light follow null geodesics in the curved spacetime geometry. Consequently, the presence of a black hole can lead to strong lensing effects on the images emerging from their surroundings. Additionally, the passage of light through the gravitational field leads to changes in the frequency and intensity of the light, which is particularly important to understand the observational imprints of accretion disks forming around black holes.

BlackHoleImages is a Wolfram Mathematica packet designed to compute the aforementioned effects in the field of isolated rotating black holes in Einstein's General Relativity, which are known as the Kerr family of spacetimes. The packet can be used for computing:

- photon orbits (null geodesics) in the fields of black holes,
- generating images of stellar backgrounds distorted by black holes, and
- generating images of accretion disks near black holes.

The packet contains three packages, `KerrNullGeodesics`, `AlphaDiskModel`, and `KerrImages`. `KerrNullGeodesics` implements the analytical solutions for null geodesics in Kerr space-times as described in detail by Gralla & Lupsasca (2020). The `AlphaDiskModel` then implements the standard α -disk model for the accretion disk near a black hole as proposed by Shakura & Sunyaev (1973), specifically its relativistic incarnation by Novikov & Thorne (1973) and Page & Thorne (1974). Finally, the `KerrImages` package uses the previous two packages to provide functions that compute the disk images and the lensed images of faraway backgrounds.

We expect this packet to be used primarily for investigating the observational characteristics of accretion disks of black holes, including dynamical processes in the disk, since the time delay of each geodesic is also computable in this packet. The packet can also be used for educational purposes, such as plotting null geodesics in the Kerr metric or illustrating the lensing effects, or for any other purpose where the computation of photon trajectories is necessary.

Statement of need

Black holes are one of the most intriguing objects in modern physics. They were one of the earliest predictions of Einstein's general relativity and starting from the 1960s they were deduced to possibly exist in the Universe through observations in various electromagnetic bands (Narayan & McClintock, 2015). Since most observations were and still are unable to resolve the spatial scales of the so-called event horizon of the black hole, one needs to understand the radiation that emerges from the interactions of the black hole with its surroundings, in particular its overall emergent spectrum. This was provided by the seminal α accretion disk model of Shakura & Sunyaev (1973) and Novikov & Thorne (1973). It was thanks to these

seminal results that the community of astronomers and astrophysicists identified the first candidates for black holes in the Universe.

Currently, the evidence for dark compact objects consistent with black holes is overwhelming. Throughout the 1990s, two teams of astronomers followed stars orbiting very close to our Galactic center and deduced on the presence of a dark compact object with the mass of¹ $4 \cdot 10^6 M_{\odot}$ (Genzel et al., 2010; Ghez & others, 2008); the leaders of the teams later shared the 2020 Nobel prize in physics (Nobel Prize Outreach, 2020). In 2015, the LIGO collaboration detected the first merger of black holes by gravitational waves (LIGO and Virgo collaborations, 2016) and more than 180 events were detected to date (LVK Collaboration, 2025). In 2017, the Event Horizon Telescope collaboration reconstructed a radio-band image of the accretion flow in the immediate vicinity of the massive black hole in the center of the M87 galaxy by the methods of very long baseline interferometry (Event Horizon Telescope collaboration, 2019a). They followed up this result in 2020 by presenting a similar image of the accretion flow from the center of our very own Galaxy the Milky Way (Event Horizon Telescope collaboration, 2022).

The images of the massive black holes from M87 and the Milky Way present a dark feature in their centers known as the “shadow” of the black hole. It corresponds to a region of the sky where the paths of the photons are bent so much that they end behind the black hole horizon, the point from beyond which not even light can escape. While this qualitative picture is correct, the precise quantitative science inferred from the observations of the Event Horizon Telescope requires the detailed modeling of the accretion flow, its radiation, and the emerging photon orbits (Event Horizon Telescope collaboration, 2019b). This is one of the important use cases our packet is aiming for.

Another important application of the packet is the interpretation of timing and spectral features of unresolved accreting black holes as observed in the burgeoning field of time-domain astronomy (Burns et al., 2025). Transients such as quasi-periodic eruptions, oscillations, or outflows emerging from systems containing accreting black holes are detected at an increasing rate. They often contain subtle timing features that require precise modelling of the accretion flows and other physical components, but possibly also the lensing effects by the black holes. Our packet can be easily extended to compute the lightcurves and/or images emerging from these systems.

Last but not least, the topic of black holes captures the imagination of the public. In the 2014 movie *Interstellar* directed by Christopher Nolan, the general movie audience was able to observe a faithful simulation of a black hole image for the first time (James et al., 2015). Our packet provides an intermediate step for relativity students and other interested parties to be able to dissect how such an image is made through a user-friendly Wolfram Mathematica packet.

On the more technical side, this is also one of the first open source releases of an implementation of the *analytical* solution for null geodesics in Kerr space-time (the only other implementation we know of is (Omwoyo, 2025)). Other implementations typically use numerical differential equation solvers (e.g. the EinsteinPy implementation (Shivotam & Bapat, 2020)). The arbitrary precision capabilities of *Wolfram Mathematica* thus allow to use this packet as an exact validation and reference solution for such machine precision codes. Using the symbolic calculus capabilities of Mathematica provided a good platform to implement the relatively complicated analytical solution as a starting point. The packet could now also be used as a reference when porting the solutions to more optimized lower-level implementations. For example, the Mathematica code *KerrGeodesics* (Warburton et al., 2023) for time-like geodesics (trajectories of massive test particles) in Kerr space-time was recently rewritten in Python as *KerrGeoPy* (Park & Nasipak, 2024), our packet could follow a similar route.

Burns, E., Fryer, C. L., Agullo, I., Andrews, J., Aydi, E., Baring, M. G., Baron, E., Boorman,

¹ $M_{\odot} \approx 2 \times 10^{30}$ kg is the solar mass.

- 91 P. G., Boroumand, M. A., Borowski, E., & others. (2025, February). *Multidisciplinary*
92 *Science in the Multimessenger Era*. <https://arxiv.org/abs/2502.03577>
- 93 Event Horizon Telescope collaboration. (2019a). First M87 Event Horizon Telescope Results.
94 I. The Shadow of the Supermassive Black Hole. *Astrophys. J. Lett.*, 875, L1. <https://doi.org/10.3847/2041-8213/ab0ec7>
95
- 96 Event Horizon Telescope collaboration. (2019b). First M87 Event Horizon Telescope Results.
97 VI. The Shadow and Mass of the Central Black Hole. *Astrophys. J. Lett.*, 875(1), L6.
98 <https://doi.org/10.3847/2041-8213/ab1141>
- 99 Event Horizon Telescope collaboration. (2022). First Sagittarius A* Event Horizon Telescope
100 Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way.
101 *Astrophys. J. Lett.*, 930(2), L12. <https://doi.org/10.3847/2041-8213/ac6674>
- 102 Genzel, R., Eisenhauer, F., & Gillessen, S. (2010). The Galactic Center Massive Black Hole
103 and Nuclear Star Cluster. *Rev. Mod. Phys.*, 82, 3121–3195. [https://doi.org/10.1103/](https://doi.org/10.1103/RevModPhys.82.3121)
104 [RevModPhys.82.3121](https://doi.org/10.1103/RevModPhys.82.3121)
- 105 Ghez, A. M., & others. (2008). Measuring Distance and Properties of the Milky Way's
106 Central Supermassive Black Hole with Stellar Orbits. *Astrophys. J.*, 689, 1044–1062.
107 <https://doi.org/10.1086/592738>
- 108 Gralla, S. E., & Lupsasca, A. (2020). Null geodesics of the Kerr exterior. *Phys. Rev. D*,
109 101(4), 044032. <https://doi.org/10.1103/PhysRevD.101.044032>
- 110 James, O., Tunzelmann, E. von, Franklin, P., & Thorne, K. S. (2015). Gravitational Lensing by
111 Spinning Black Holes in Astrophysics, and in the Movie *Interstellar*. *Class. Quant. Grav.*,
112 32(6), 065001. <https://doi.org/10.1088/0264-9381/32/6/065001>
- 113 LIGO and Virgo collaborations. (2016). Observation of Gravitational Waves from a Binary Black
114 Hole Merger. *Phys. Rev. Lett.*, 116(6), 061102. [https://doi.org/10.1103/PhysRevLett.](https://doi.org/10.1103/PhysRevLett.116.061102)
115 [116.061102](https://doi.org/10.1103/PhysRevLett.116.061102)
- 116 LVK Collaboration. (2025). *Gravitational wave open science centre*. <https://gwosc.org/>.
- 117 Narayan, R., & McClintock, J. E. (2015). Observational evidence for black holes. In A.
118 Ashtekar, B. K. Berger, J. Isenberg, & M. MacCallum (Eds.), *General relativity and*
119 *gravitation: A centennial perspective* (pp. 97–161). Cambridge University Press.
- 120 Nobel Prize Outreach. (2020). *Nobel prize in physics 2020 webpage*. [https://www.nobelprize.](https://www.nobelprize.org/prizes/physics/2020/summary/)
121 [org/prizes/physics/2020/summary/](https://www.nobelprize.org/prizes/physics/2020/summary/).
- 122 Novikov, I. D., & Thorne, K. S. (1973). Astrophysics of black holes. *Black Holes (Les Astres*
123 *Occlus)*, 1, 343–450.
- 124 Omwoyo, E. (2025). *KerrGeodesics:NullGeodesics branch on github*. [https://github.com/](https://github.com/BlackHolePerturbationToolkit/KerrGeodesics/tree/NullGeodesics)
125 [BlackHolePerturbationToolkit/KerrGeodesics/tree/NullGeodesics](https://github.com/BlackHolePerturbationToolkit/KerrGeodesics/tree/NullGeodesics).
- 126 Page, D. N., & Thorne, K. S. (1974). Disk-Accretion onto a Black Hole. Time-Averaged
127 Structure of Accretion Disk. *Astrophys. J.*, 191, 499–506. <https://doi.org/10.1086/152990>
- 128 Park, S., & Nasipak, Z. (2024). KerrGeoPy: A python package for computing timelike
129 geodesics in kerr spacetime. *Journal of Open Source Software*, 9(98), 6587. <https://doi.org/10.21105/joss.06587>
130
- 131 Shakura, N. I., & Sunyaev, R. A. (1973). Black holes in binary systems. Observational
132 appearance. *Astron. Astrophys.*, 24, 337–355.
- 133 Shivotam, J., & Bapat, S. (2020). *EinsteinPy-Geodesics on github*. [https://github.com/](https://github.com/einsteinpy/einsteinpy-geodesics)
134 [einsteinpy/einsteinpy-geodesics](https://github.com/einsteinpy/einsteinpy-geodesics).
- 135 Warburton, N., Wardell, B., Long, O., Upton, S., Lynch, P., Nasipak, Z., & Stein, L. C. (2023).
136 *KerrGeodesics* (Version 0.9.0). Zenodo. <https://doi.org/10.5281/zenodo.8108265>