ASSIGNMENT 6

Y.Nagarani

Download all python codes from

https://github.com/Y.Nagarani/ASSIGNMENT6/tree/main/CODES

and latex-tikz codes from

https://github.com/Y.Nagarani/ASSIGNMENT6/tree/main

1 Question No 2.5

In
$$\triangle ABC$$
, $A = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$, $B = \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}$, $C = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$. Find

∠B.

2 SOLUTION Let, $\mathbf{A} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$, $\mathbf{B} = \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}$, $\mathbf{C} = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$.

Now,

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} 2 \\ 2 \\ 3 \end{pmatrix} \tag{2.0.1}$$

$$\mathbf{C} - \mathbf{B} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \tag{2.0.2}$$

We know that,

$$\mathbf{B} = \arccos\left(\frac{(\mathbf{A} - \mathbf{B})^T (\mathbf{C} - \mathbf{B})}{\|\mathbf{A} - \mathbf{B}\|\|\mathbf{C} - \mathbf{B}\|}\right) \tag{2.0.3}$$

Then

$$\|\mathbf{A} - \mathbf{B}\| = \sqrt{17} \tag{2.0.4}$$

$$\|\mathbf{C} - \mathbf{B}\| = \sqrt{6} \tag{2.0.5}$$

$$(\mathbf{A} - \mathbf{B})^{\mathrm{T}}(\mathbf{C} - \mathbf{B}) = 10 \tag{2.0.6}$$

Substitute above values in (2.0.3) then,

$$\mathbf{B} = \arccos\left(\frac{10}{\sqrt{17}\sqrt{6}}\right) \tag{2.0.7}$$

$$\mathbf{B} = 66.15 \tag{2.0.8}$$




Fig. 0: △ABC