

# The selection and use of essential medicines, 2025

---

WHO  
Model List of Essential Medicines

24th list

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: “This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition”.

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

**Suggested citation.** The selection and use of essential medicines, 2025: WHO Model List of Essential Medicines, 24th list. Geneva: World Health Organization; 2025. <https://doi.org/10.2471/B09474>. Licence: CC BY-NC-SA 3.0 IGO.

**Cataloguing-in-Publication (CIP) data.** CIP data are available at <https://iris.who.int/>.

**Sales, rights and licensing.** To purchase WHO publications, see <https://www.who.int/publications/book-orders>. To submit requests for commercial use and queries on rights and licensing, see <https://www.who.int/copyright>.

**Third-party materials.** If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

**General disclaimers.** The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

This publication contains the collective views of an international group of experts and does not necessarily represent the decisions or the policies of WHO.

The recommendations contained in this publication are based on the advice of independent experts, who have considered the best available evidence, a risk-benefit analysis and other factors, as appropriate. This publication may include recommendations on the use of medicinal products for an indication, in a dosage form, dose regimen, population or other use parameters that are not included in the approved labelling. Relevant stakeholders should familiarize themselves with applicable national legal and ethical requirements. WHO does not accept liability for the procurement, distribution, and/or administration of any product for any use.

## CORRIGENDA (20 October 2025)

The selection and use of essential medicines, 2025: WHO Model List of Essential Medicines, 24th list.

WHO Digital Object Identifier (DOI): <https://doi.org/10.2471/B09474>

Page 41, lines 7–14

Delete:

<input type="checkbox"/> perindopril + <input type="checkbox"/> amlodipine + <input type="checkbox"/> indapamide  Therapeutic alternatives:  - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for perindopril)  - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)  - chlorthalidone, chlorothiazide, hydrochlorothiazide (for indapamide)	<b>Solid oral dosage form:</b> 5 mg + 5 mg + 1.25 mg; 5 mg + 10 mg + 2.5 mg; 10 mg + 5 mg + 1.25 mg; 10 mg + 10 mg + 2.5 mg.
--	--

Insert:

<input type="checkbox"/> perindopril + <input type="checkbox"/> amlodipine + <input type="checkbox"/> indapamide  Therapeutic alternatives:  - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for perindopril)  - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)  - chlorthalidone, chlorothiazide, hydrochlorothiazide (for indapamide)	<b>Solid oral dosage form:</b> 5 mg (arginine) + 5 mg + 1.25 mg; 5 mg (arginine) + 10 mg + 1.25 mg; 10 mg (arginine) + 5 mg + 2.5 mg; 10 mg (arginine) + 10 mg + 2.5 mg.
--	--

Page 41, lines 20–28

Delete:

<input type="checkbox"/> valsartan + <input type="checkbox"/> amlodipine + <input type="checkbox"/> hydrochlorothiazide  Therapeutic alternatives:  - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for valsartan)  - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)  - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	<b>Solid oral dosage form:</b> 5 mg + 160 mg + 12.5 mg; 5 mg + 160 mg + 25 mg; 10 mg + 160 mg + 12.5 mg; 10 mg + 160 mg + 25 mg; 10 mg + 320 mg + 25 mg.
--	--

Insert:

<input type="checkbox"/> valsartan + <input type="checkbox"/> amlodipine + <input type="checkbox"/> hydrochlorothiazide  Therapeutic alternatives:  - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for valsartan)  - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)  - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	<b>Solid oral dosage form:</b> 160 mg + 5 mg + 12.5 mg; 160 mg + 5 mg + 25 mg; 160 mg + 10 mg + 12.5 mg; 160 mg + 10 mg + 25 mg; 320 mg + 10 mg + 25 mg.
--	--

**Page 49, lines 30–35**

*Delete:*

<input type="checkbox"/> semaglutide* Therapeutic alternatives: - dulaglutide - liraglutide - tirzepatide <i>*including quality-assured biosimilars</i>	<b>Injection solution:</b> 0.68 mg/mL; 1.34 mg/mL.
--	--

*Insert:*

<input type="checkbox"/> semaglutide* Therapeutic alternatives: - dulaglutide - liraglutide - tirzepatide <i>*including quality-assured biosimilars</i>	<b>Injection solution:</b> 0.68 mg/mL; 1.34 mg/mL; 2.68 mg/mL.
--	--

---

# WHO Model List of Essential Medicines – 24th List (2025)

## Explanatory notes

---

The **core list** presents a list of minimum medicine needs for a basic health-care system, listing the most efficacious, safe and cost-effective medicines for priority conditions. Priority conditions are selected on the basis of current and estimated future public health relevance, and potential for safe and cost-effective treatment.

Where the **[c]** symbol is placed next to an individual medicine or strength of medicine on the core list it signifies that there is a specific indication for restricting its use to children.

The **complementary list** presents essential medicines for priority diseases, for which specialized diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training are needed. In case of doubt medicines may also be listed as complementary on the basis of consistent higher costs or less attractive cost-effectiveness in a variety of settings.

Where the **[c]** symbol is placed next to an individual medicine or strength of medicine on the complementary list it signifies that the medicine(s) require(s) specialist diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training for their use in children.

The **square box symbol (□)** is intended to indicate therapeutic alternatives to the listed medicine that may be considered for selection in national essential medicines lists. Alternatives may be individual medicines, or multiple medicines within a pharmacological class or chemical subgroup, defined at the 4th level of the Anatomical Therapeutic Chemical (ATC) classification, which have similar clinical effectiveness and safety. The listed medicine should be the example of the class or subgroup for which there is the best evidence for effectiveness and safety. In some cases, this may be the first medicine that is licensed for marketing; in other instances, subsequently licensed compounds may be safer or more effective. Where there is no difference in terms of efficacy and safety data, the listed medicine should be the one that is generally available at the lowest price, based on international drug price information sources. Not all square box listings are applicable to medicine selection for children. A square box is not used to indicate alternative generic brands of the same small molecule medicines, nor alternative biosimilars of biological medicines. However, the selection and use of quality-assured generics and biosimilars of essential medicines at country level is recommended.

National lists should not use a similar symbol and should be specific in their final selection, which would depend on local availability and price.

The **[a]** symbol indicates that there is an age or weight restriction on use of the medicine.

The presence of an entry on the Essential Medicines List carries no assurance as to pharmaceutical quality. It is the responsibility of the relevant national or regional drug regulatory authority to ensure that each product is of appropriate pharmaceutical quality (including stability) and that, when relevant, different products are interchangeable.

For recommendations and advice concerning all aspects of the quality assurance of medicines see the WHO website <https://www.who.int/teams/health-product-and-policy-standards/standards-and-specifications/norms-and-standards-for-pharmaceuticals/guidelines/quality-assurance>

Medicines and dosage forms are listed in alphabetical order within each section and the order of listing does not imply preference for one form over another. Standard treatment guidelines should be consulted for information on appropriate dosage forms.




The main terms used for dosage forms in the Essential Medicines List can be found in Table 1.

Definitions of many of these terms and pharmaceutical quality requirements applicable to the different categories are published in the current edition of *The International Pharmacopoeia*. <https://www.who.int/teams/health-product-policy-and-standards/standards-and-specifications/norms-and-standards-for-pharmaceuticals/international-pharmacopoeia>.

# WHO Model List of Essential Medicines – 24th List (2025)

<b>1. ANAESTHETICS, PREOPERATIVE MEDICINES AND MEDICAL GASES</b>	
<b>1.1 General anaesthetics and oxygen</b>	
<b>1.1.1 Inhalational medicines</b>	
isoflurane	Inhalation.
nitrous oxide*	Inhalation. *Piped nitrous oxide is a major source of atmospheric pollution from healthcare facilities. Point-of-care cylinders are the preferred delivery system over centrally-supplied (piped) delivery systems.
oxygen	Inhalation (medical gas).
sevoflurane	Inhalation.
<b>1.1.2 Injectable medicines</b>	
ketamine	Injection: 10 mg/mL [c]; 50 mg/mL (as hydrochloride) in vial.
<input type="checkbox"/> propofol Therapeutic alternatives: - thiopental	Injection: 10 mg/mL; 20 mg/mL.
<b>1.2 Local anaesthetics</b>	
<input type="checkbox"/> bupivacaine Therapeutic alternatives to be reviewed	Injection: 0.25%; 0.5% (hydrochloride). Injection for spinal anaesthesia: 0.5% (hydrochloride) in 4 mL ampoule in 8% glucose solution.
<input type="checkbox"/> lidocaine Therapeutic alternatives to be reviewed	Injection: 0.5% [c]; 1%; 2% (hydrochloride). Injection for spinal anaesthesia: 5% (hydrochloride) in 2 mL ampoule in 7.5% glucose solution. Topical forms: 2% to 4% (hydrochloride).
lidocaine + epinephrine (adrenaline)	Dental cartridge: 2% (hydrochloride) + epinephrine 1:80 000. Injection: 1%; 2% (hydrochloride or sulfate) + epinephrine 1:200 000.
<b>Complementary List</b>	
<i>ephedrine*</i>	Injection: 30 mg/mL (hydrochloride) in 1 mL ampoule. *For use in spinal anaesthesia during delivery, to prevent hypotension.
<b>1.3 Preoperative medication and sedation for short-term procedures</b>	
atropine	Injection: 400 micrograms/mL [c]; 1 mg/mL (sulfate) in 1 mL ampoule or vial.
<input type="checkbox"/> midazolam Therapeutic alternatives to be reviewed	Injection: 1 mg/mL in 5 mL vial; 5 mg/mL in 1 mL or 3 mL vial. Oral liquid: 2 mg/mL [c]. Solid oral dosage form: 7.5 mg; 15 mg.
morphine	Injection: 1 mg/mL [c]; 2 mg/mL [c]; 10 mg/mL (sulfate or hydrochloride) in 1 mL ampoule.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>1.4 Medical gases</b>	
oxygen*	<b>Inhalation</b> For use in the management of hypoxaemia. *No more than 30% oxygen should be used to initiate resuscitation of neonates less than or equal to 32 weeks of gestation.
<b>2. MEDICINES FOR PAIN AND PALLIATIVE CARE</b>	
<b>2.1 Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)</b>	
acetylsalicylic acid	<b>Suppository:</b> 50 mg to 150 mg. <b>Tablet:</b> 100 mg to 500 mg.
ibuprofen 	<b>Oral liquid:</b> 100 mg/5 mL [c], 200 mg/5 mL. <b>Tablet:</b> 200 mg; 400 mg; 600 mg.  Not in children less than 3 months.
paracetamol (acetaminophen)*	<b>Oral liquid:</b> 120 mg/5 mL or 125 mg/5 mL**, 250 mg/5 mL [c]. **The presence of both 120 mg/5 mL and 125 mg/5 mL strengths on the same market would cause confusion in prescribing and dispensing and should be avoided. <b>Suppository:</b> 100 mg, 250 mg [c]. <b>Tablet:</b> 250 mg, 325 mg, 500 mg. <b>Tablet (dispersible):</b> 100 mg, 250 mg [c]. *Not recommended for anti-inflammatory use due to lack of proven benefit to that effect.
<b>2.2 Opioid analgesics</b>	
codeine	<b>Tablet:</b> 30 mg (phosphate).
fentanyl*	<b>Transdermal patch:</b> 12 micrograms/hr; 25 micrograms/hr; 50 micrograms/hr; 75 micrograms/hr; 100 micrograms/hr. *For the management of cancer pain
 morphine Therapeutic alternatives: - hydromorphone - oxycodone	<b>Granules (slow release; to mix with water):</b> 20 mg; 30 mg; 60 mg; 100 mg; 200 mg (morphine sulfate). <b>Injection:</b> 1 mg/mL [c]; 2 mg/mL [c]; 10 mg/mL (morphine hydrochloride or morphine sulfate) in 1 mL ampoule. <b>Oral liquid:</b> 5 mg/5mL [c]; 10 mg/5 mL (morphine hydrochloride or morphine sulfate). <b>Solid oral dosage form (slow release):</b> 5 mg [c]; 10 mg; 30 mg; 60 mg; 100 mg; 200 mg (morphine hydrochloride or morphine sulfate). <b>Tablet (immediate release):</b> 10 mg (morphine sulfate).
<b>Complementary list</b>	
methadone*	<b>Tablet:</b> 5 mg; 10 mg (hydrochloride). <b>Oral liquid:</b> 5 mg/5 mL; 10 mg/5 mL (hydrochloride). <b>Concentrate for oral liquid:</b> 25 mg/5mL; 50 mg/5mL (hydrochloride). *For the management of cancer pain.

## WHO Model List of Essential Medicines – 24th List (2025)

2.3 Medicines for other common symptoms in palliative care	
amitriptyline	<b>Oral liquid:</b> 25 mg/5 mL [c]. <b>Tablet:</b> 10 mg; 25 mg; 75 mg.
cyclizine [c]	<b>Injection:</b> 50 mg/mL (lactate). <b>Tablet:</b> 50 mg (hydrochloride).
dexamethasone	<b>Injection:</b> 4 mg/mL dexamethasone phosphate (as sodium phosphate) (equivalent to 3.3 mg/mL dexamethasone base) in 1 mL ampoule. <b>Oral liquid:</b> 0.5 mg/5 mL [c]; 2 mg/5 mL (as sodium phosphate). <b>Tablet:</b> 0.5 mg; 0.75 mg; 1.5 mg; 2 mg; 4 mg (as dexamethasone base).
diazepam	<b>Injection:</b> 5 mg/mL. <b>Oral liquid:</b> 2 mg/5 mL. <b>Rectal gel:</b> 5 mg/mL in 0.5 mL, 2 mL, 4 mL rectal delivery system. <b>Rectal solution:</b> 2 mg/mL in 1.25 mL, 2.5 mL rectal tube; 4 mg/mL in 2.5 mL rectal tube. <b>Tablet (scored):</b> 2 mg [c]; 5 mg; 10 mg.
docusate sodium	<b>Capsule:</b> 100 mg. <b>Oral liquid:</b> 12.5 mg/5 mL [c]; 50 mg/5 mL.
fluoxetine	<b>Solid oral dosage form:</b> 20 mg (as hydrochloride).
haloperidol	<b>Injection:</b> 5 mg in 1 mL ampoule. <b>Oral liquid:</b> 2 mg/mL. <b>Solid oral dosage form:</b> 0.5 mg; 2 mg; 5 mg.
hyoscine butylbromide	<b>Injection:</b> 20 mg/mL.
hyoscine hydrobromide [c]	<b>Injection:</b> 400 micrograms/mL; 600 micrograms/mL. <b>Transdermal patches:</b> 1 mg/72 hours.
lactulose [c]	<b>Oral liquid:</b> 3.3 to 3.4 g/5 mL.
loperamide	<b>Solid oral dosage form:</b> 2 mg.
metoclopramide	<b>Injection:</b> 5 mg/mL (hydrochloride) in 2 mL ampoule. <b>Oral liquid:</b> 5 mg/5 mL. <b>Solid oral form:</b> 10 mg (hydrochloride).
midazolam	<b>Injection*:</b> 1 mg/mL in 5 mL vial; 5 mg/mL in 1 mL, 3 mL vial. *May be used for buccal administration when solution for oromucosal administration is not available. <b>Oral liquid:</b> 2 mg/mL [c]. <b>Solid oral dosage form:</b> 7.5 mg; 15 mg. <b>Solution for oromucosal administration:</b> 5 mg/mL in 0.5 mL, 1 mL, 1.5 mL, 2 mL pre-filled syringe; 10 mg/mL in 0.25 mL, 0.5 mL, 0.75 mL, 1 mL prefilled syringe.



## WHO Model List of Essential Medicines – 24th List (2025)

<input type="checkbox"/> ondansetron Therapeutic alternatives: - dolasetron - granisetron - palonosetron - tropisetron	<b>Injection:</b> 2 mg/mL in 2 mL, 4 mL ampoule (as hydrochloride dihydrate). <b>Oral liquid:</b> 4 mg/5 mL (as hydrochloride dihydrate). <b>Solid oral dosage form:</b> 4 mg; 8 mg (as hydrochloride dihydrate).
senna	<b>Oral liquid:</b> 7.5 mg/5 mL.
<b>3. ANTIALLERGICS AND MEDICINES USED IN ANAPHYLAXIS</b>	
dexamethasone	<b>Injection:</b> 4 mg/mL dexamethasone phosphate (as sodium phosphate) (equivalent to 3.3 mg/mL dexamethasone base) in 1 mL ampoule. <b>Oral liquid:</b> 0.5 mg/5 mL [c]; 2 mg/5 mL (as sodium phosphate). <b>Tablet:</b> 0.5 mg; 0.75 mg; 1.5 mg; 2 mg; 4 mg (as dexamethasone base).
epinephrine (adrenaline)	<b>Injection:</b> 1 mg/mL (as hydrochloride or hydrogen tartrate) in 1 mL ampoule.
hydrocortisone	<b>Powder for injection:</b> 100 mg (as sodium succinate) in vial.
<input type="checkbox"/> loratadine* Therapeutic alternatives: - cetirizine - fexofenadine	<b>Oral liquid:</b> 1 mg/mL. <b>Tablet:</b> 10 mg. <b>Tablet (chewable):</b> 5 mg [c]; 10 mg [c]. <i>*There may be a role for sedating antihistamines for limited indications (EMLC).</i>
<input type="checkbox"/> prednisolone Therapeutic alternatives: - prednisone	<b>Oral liquid:</b> 5 mg/mL [c]. <b>Tablet:</b> 5 mg; 25 mg.
<b>4. ANTIDOTES AND OTHER SUBSTANCES USED IN POISONINGS</b>	
<b>4.1 Non-specific</b>	
charcoal, activated	<b>Granules for oral suspension*:</b> 50 mg. <i>*Alternative formulations of activated charcoal may be used if granules are not available.</i>
<b>4.2 Specific</b>	
acetylcysteine	<b>Injection:</b> 200 mg/mL in 10 mL ampoule.
atropine	<b>Injection:</b> 1 mg/mL (sulfate) in 1 mL ampoule or vial.
calcium gluconate	<b>Injection:</b> 100 mg/mL (10%) in 10 mL ampoule or vial.
methylnthioninium chloride (methylene blue)	<b>Injection:</b> 10 mg/mL in 10 mL ampoule.
naloxone	<b>Injection:</b> 400 micrograms (hydrochloride) in 1 mL ampoule.
penicillamine	<b>Solid oral dosage form:</b> 250 mg.
potassium ferric hexacyano-ferrate(II) ·2H <sub>2</sub> O (Prussian blue)	<b>Powder for oral administration.</b>
sodium nitrite	<b>Injection:</b> 30 mg/mL in 10 mL ampoule.
sodium thiosulfate	<b>Injection:</b> 250 mg/mL in 50 mL ampoule.

## WHO Model List of Essential Medicines – 24th List (2025)

<i>Complementary List</i>	
<i>deferoxamine</i>	<b>Powder for injection:</b> 500 mg (mesilate) in vial.
<i>dimercaprol</i>	<b>Injection in oil:</b> 50 mg/mL in 2 mL ampoule; 100 mg/mL in 3 mL ampoule.
<i>fomepizole</i>	<b>Injection:</b> 5 mg/mL (sulfate) in 20 mL ampoule or 1 g/mL (base) in 1.5 mL ampoule or vial.
<i>sodium calcium edetate</i>	<b>Injection:</b> 200 mg/mL in 5 mL ampoule.
<i>succimer</i>	<b>Solid oral dosage form:</b> 100 mg.
<b>5. MEDICINES FOR NEUROLOGICAL DISORDERS</b>	
<b>5.1 Medicines for central nervous system disorders</b>	
<b>5.1.1 Antiseizure medicines</b>	
carbamazepine	<b>Oral liquid:</b> 100 mg/5 mL. <b>Tablet (chewable):</b> 100 mg; 200 mg. <b>Tablet (scored):</b> 100 mg; 200 mg; 400 mg.
diazepam	<b>Rectal gel:</b> 5 mg/mL in 0.5 mL, 2 mL, 4 mL rectal delivery system. <b>Rectal solution:</b> 2 mg/mL in 1.25 mL, 2.5 mL rectal tube; 4 mg/mL in 2.5 mL rectal tube.
lamotrigine*	<b>Tablet:</b> 25 mg; 50 mg; 100 mg; 200 mg. <b>Tablet (chewable, dispersible):</b> 2 mg; 5 mg; 25 mg; 50 mg; 100 mg; 200 mg. *For use as adjunctive therapy for treatment-resistant partial or generalized seizures.
levetiracetam	<b>Oral liquid:</b> 100 mg/mL <b>Tablet:</b> 250 mg; 500 mg; 750 mg; 1000 mg.
<input type="checkbox"/> lorazepam Therapeutic alternatives: - diazepam (injection) - midazolam (injection)	<b>Injection:</b> 2 mg/mL in 1 mL ampoule; 4 mg/mL in 1 mL ampoule.
magnesium sulfate*	<b>Injection:</b> 0.5 g/mL in 2 mL ampoule (equivalent to 1 g in 2 mL; 50% weight/volume); 0.5 g/mL in 10 mL ampoule (equivalent to 5 g in 10 mL; 50% weight/volume). *For use in eclampsia and severe pre-eclampsia and not for other convulsant disorders.
midazolam	<b>Solution for oromucosal administration:</b> 5 mg/mL in 0.5 mL, 1 mL, 1.5 mL, 2 mL pre-filled syringe; 10 mg/mL in 0.25 mL, 0.5 mL, 0.75 mL, 1 mL pre-filled syringe. <b>Injection*:</b> 1 mg/mL in 5 mL vial; 5 mg/mL in 1 mL or 3 mL vial. *For buccal administration when solution for oromucosal administration is not available.
phenobarbital	<b>Injection:</b> 30 mg/mL or 60 mg/mL [c], 200 mg/mL (sodium). <b>Oral liquid:</b> 15 mg/5 mL. <b>Tablet:</b> 15 mg; 30 mg; 60 mg; 100 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

phenytoin	<b>Injection:</b> 50 mg/mL (phenytoin sodium). <b>Oral liquid:</b> 30 mg/5 mL (phenytoin). <b>Solid oral dosage form:</b> 25 mg; 50 mg; 100 mg (phenytoin sodium). <b>Tablet (chewable):</b> 50 mg (phenytoin).
prednisolone [c]	<b>Oral liquid:</b> 1 mg/mL. <b>Tablet:</b> 1 mg; 5 mg; 10 mg.
valproic acid (sodium valproate)* <i>*Valproic acid (sodium valproate) is not recommended in women and girls of childbearing potential owing to the high risk of birth defects and neurodevelopmental disorders in children exposed to valproic acid (sodium valproate) in the womb.</i>	<b>Oral liquid:</b> 200 mg/5 mL. <b>Tablet (crushable):</b> 100 mg. <b>Tablet (enteric-coated):</b> 200 mg; 500 mg.
<b>Complementary List</b>	
ethosuximide	<b>Capsule:</b> 250 mg. <b>Oral liquid:</b> 250 mg/5 mL.
levetiracetam	<b>Concentrate solution for infusion:</b> 100 mg/mL in 5 mL ampoule or vial. <b>Solution for infusion:</b> 5 mg/mL; 10 mg/mL; 15 mg/mL in 100 mL bag.
valproic acid (sodium valproate)* <i>*Valproic acid (sodium valproate) is not recommended in women and girls of childbearing potential owing to the high risk of birth defects and neurodevelopmental disorders in children exposed to valproic acid (sodium valproate) in the womb.</i>	<b>Injection:</b> 100 mg/mL in 3 mL, 4 mL, 10 mL ampoule.
<b>5.1.2 Medicines for multiple sclerosis</b>	
<b>Complementary List</b>	
cladribine	<b>Tablet:</b> 10 mg.
glatiramer acetate	<b>Injection (subcutaneous):</b> 20 mg/mL; 40 mg/mL in pre-filled syringe.
rituximab* <i>*including quality-assured biosimilars</i>	<b>Injection (intravenous):</b> 500 mg/50 mL in 50 mL vial.
<b>5.1.3 Medicines for parkinsonism</b>	
<input type="checkbox"/> biperiden Therapeutic alternatives: – trihexyphenidyl	<b>Injection:</b> 5 mg (lactate) in 1 mL ampoule. <b>Tablet:</b> 2 mg (hydrochloride).
levodopa + <input type="checkbox"/> carbidopa Therapeutic alternatives: – benserazide (for carbidopa)	<b>Tablet:</b> 100 mg + 10 mg; 100 mg + 25 mg; 250 mg + 25 mg.
<b>5.1.4 Medicines for cerebral palsy</b>	
<b>Complementary List</b>	
baclofen	<b>Intrathecal injection:</b> 500 micrograms/mL in ampoule. <b>Oral liquid:</b> 10 mg/5 mL. <b>Tablet:</b> 10 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

5.1.5 Medicines for headache disorders		
5.1.5.1 Medicines for acute migraine attacks		
acetylsalicylic acid	Tablet: 300 mg to 500 mg.	
<input type="checkbox"/> ibuprofen  Therapeutic alternatives: - naproxen	Oral liquid: 100 mg/5 mL [c]. Tablet: 200 mg; 400 mg.	
paracetamol (acetaminophen)	Oral liquid: 120 mg/5 mL or 125 mg/5 mL*; 250 mg/5 mL [c].  *The presence of both 120 mg/5 mL and 125 mg/5 mL strengths on the same market would cause confusion in prescribing and dispensing and should be avoided.  Suppository: 250 mg [c].  Tablet: 250 mg; 325 mg; 500 mg.  Tablet (dispersible): 100 mg, 250 mg [c].	
<input type="checkbox"/> sumatriptan  Therapeutic alternatives: - eletriptan	Tablet: 50 mg.	
5.1.5.2 Medicines for migraine prophylaxis		
propranolol	Tablet: 10 mg [c]; 40 mg (hydrochloride).	
5.1.5.3 Medicines for cluster headache		
prednisolone	Tablet: 5 mg; 25 mg.	
sumatriptan	Injection (subcutaneous): 6 mg/ 0.5 mL in pre-filled syringe or pen.	
verapamil	Tablet (immediate-release): 40 mg; 80 mg; 120 mg.	
	Tablet (extended-release): 120 mg; 180 mg; 240 mg.	
5.1.6 Medicines for central nervous system infections		
5.1.6.1 Medicines for bacterial central nervous system infections		
amoxicillin	Powder for injection: 250 mg; 500 mg; 1 g (as sodium) in vial. Powder for oral liquid: 125 mg/5 mL; 250 mg/5 mL (as trihydrate) [c]. Solid oral dosage form: 250 mg; 500 mg (as trihydrate). Tablet (dispersible, scored): 250 mg; 500 mg (as trihydrate) [c].	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis
ampicillin	Powder for injection: 500 mg; 1 g (as sodium) in vial.	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis
benzylpenicillin	Powder for injection: 600 mg (= 1 million IU); 3 g (= 5 million IU) (sodium or potassium salt) in vial.	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis

## WHO Model List of Essential Medicines – 24th List (2025)

cefotaxime*	Powder for injection: 250 mg; 500 mg; 1 g; 2 g (as sodium) in vial. *3rd generation cephalosporin of choice for use in hospitalized neonates.	
	FIRST CHOICE – Acute bacterial meningitis	SECOND CHOICE
ceftriaxone*[a]	Powder for injection: 250 mg; 500 mg; 1 g (as sodium) in vial. *Do not administer with calcium and avoid in infants with hyperbilirubinaemia. [a] > 41 weeks corrected gestational age.	
	FIRST CHOICE – Acute bacterial meningitis	SECOND CHOICE
chloramphenicol	Oily suspension for injection*: 0.5 g/mL (as sodium succinate) in 2 mL ampoule. *Only for the presumptive treatment of epidemic meningitis in children older than 2 years. Powder for injection: 1 g (sodium succinate) in vial.	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis
gentamicin	Injection: 10 mg/mL (as sulfate); 40 mg/mL (as sulfate) in 2 mL vial.	
	FIRST CHOICE – Acute bacterial meningitis in neonates [c]	SECOND CHOICE
Complementary List		
meropenem*[a]	Powder for injection: 500 mg (as trihydrate); 1 g (as trihydrate) in vial [a] > 3 months.	
	FIRST CHOICE	SECOND CHOICE – Acute bacterial meningitis in neonates [c]
5.1.6.2 Medicines for viral central nervous system infections		
<input type="checkbox"/> aciclovir Therapeutic alternatives: - valaciclovir (oral)	Oral liquid: 200 mg/5 mL [c]. Powder for solution for infusion: 250 mg (as sodium dihydrate) in vial. Solution for infusion: 25 mg/mL (as sodium) in vial. Tablet: 200 mg.	
5.2 Medicines for peripheral nervous system disorders		
5.2.1 Medicines for Guillain-Barré syndrome		
Complementary List		
normal immunoglobulin	Intravenous administration: 5%; 10% protein solution.	
5.2.2 Medicines for myasthenia gravis		
neostigmine	Injection: 500 micrograms/mL (methylsulfate) in 1 mL ampoule; 2.5 mg/mL (methylsulfate) in 1 mL ampoule. Tablet: 15 mg (bromide).	
Complementary List		
pyridostigmine	Injection: 5 mg/mL (bromide) in ampoule or vial. Tablet (scored): 60 mg (bromide).	

## WHO Model List of Essential Medicines – 24th List (2025)

<b>6. ANTI-INFECTIVE MEDICINES</b>	
<b>6.1 Anthelmintics</b>	
<b>6.1.1 Intestinal anthelmintics</b>	
albendazole	Tablet (chewable, scored): 400 mg.
ivermectin	Tablet: 3 mg.
levamisole	Tablet: 50 mg; 150 mg (as hydrochloride).
mebendazole	Tablet (chewable): 100 mg; 500 mg.
niclosamide	Tablet (chewable): 500 mg.
praziquantel	Tablet: 150 mg, 500 mg. Tablet (scored): 600 mg.
pyrantel	Tablet (chewable): 250 mg (as embonate or pamoate).
<b>6.1.2 Antifilarials</b>	
albendazole	Tablet (chewable, scored): 400 mg.
diethylcarbamazine	Tablet: 50 mg; 100 mg (dihydrogen citrate).
<input type="checkbox"/> ivermectin Therapeutic alternatives: - moxidectin	Tablet: 3 mg.
<b>6.1.3 Antischistosomes and other antitrepatode medicines</b>	
<input type="checkbox"/> praziquantel* Therapeutic alternatives: - arpraziquantel (Tablet (dispersible): 150 mg) [c] *The square box applies only to the listing of praziquantel on the EMLc for schistosomiasis	Tablet: 150 mg; 500 mg. Tablet (scored): 600 mg.
triclabendazole	Tablet (scored): 250 mg.
<b>Complementary List</b>	
oxamniquine*	Capsule: 250 mg. Oral liquid: 250 mg/5 mL. *For use when praziquantel treatment fails.
<b>6.1.4 Cysticidal medicines</b>	
<b>Complementary List</b>	
albendazole	Tablet (chewable): 200 mg [c]. Tablet (chewable, scored): 400 mg.
mebendazole	Tablet (chewable): 100 mg [c], 500 mg.
praziquantel	Tablet: 150 mg, 500 mg. Tablet (scored): 600 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

### 6.2 Antibacterials

To assist in the development of tools for antibiotic stewardship at local, national and global levels and to reduce antimicrobial resistance, the Access, Watch, Reserve (AWaRe) classification of antibiotics has been developed by WHO – where antibiotics are classified into different groups to emphasize the importance of their appropriate use.

#### ACCESS GROUP ANTIBIOTICS

This group includes antibiotics that have activity against a wide range of commonly encountered susceptible pathogens while also showing lower resistance potential than antibiotics in the other groups. Selected Access group antibiotics are recommended as essential first or second choice empiric treatment options for infectious syndromes reviewed by the EML Expert Committee and are listed as individual medicines on the Model Lists to improve access and promote appropriate use. They are essential antibiotics that should be widely available, affordable and quality assured.

#### WATCH GROUP ANTIBIOTICS

This group includes antibiotic classes that have higher resistance potential and includes most of the highest priority agents among the Critically Important Antimicrobials for Human Medicine and/or antibiotics that are at relatively high risk of selection of bacterial resistance. These medicines should be prioritized as key targets of stewardship programs and monitoring. Selected Watch group antibiotics are recommended as essential first or second choice empiric treatment options for a limited number of specific infectious syndromes and are listed as individual medicines on the Model Lists.

#### RESERVE GROUP ANTIBIOTICS

This group includes antibiotics and antibiotic classes that should be reserved for treatment of confirmed or suspected infections due to multi-drug-resistant organisms. Reserve group antibiotics should be treated as “last resort” options. Selected Reserve group antibiotics are listed as individual medicines on the Model Lists when they have a favourable risk-benefit profile and proven activity against “Critical Priority” or “High Priority” pathogens identified by the WHO Priority Pathogens List, notably carbapenem resistant *Enterobacteriaceae*. These antibiotics should be accessible, but their use should be tailored to highly specific patients and settings, when all alternatives have failed or are not suitable. These medicines could be protected and prioritized as key targets of national and international stewardship programmes involving monitoring and utilization reporting, to preserve their effectiveness.

## WHO Model List of Essential Medicines – 24th List (2025)

6.2.1 Access group antibiotics		
amikacin	<b>Injection:</b> 50 mg/mL (as sulfate) [c]; 250 mg/mL (as sulfate) in 2 mL vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– High-risk febrile neutropenia</li> <li>– Pyelonephritis or prostatitis (severe)</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Sepsis in neonates and children [c]</li> </ul>
amoxicillin	<b>Powder for injection:</b> 250 mg; 500 mg; 1 g (as sodium) in vial. <b>Powder for oral liquid:</b> 125 mg/5 mL; 250 mg/5 mL (as trihydrate) [c]. <b>Solid oral dosage form:</b> 250 mg; 500 mg; 1 g (as trihydrate). <b>Tablet (dispersible, scored):</b> 250 mg; 500 mg (as trihydrate) [c].	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Community acquired pneumonia (mild to moderate)</li> <li>– Community acquired pneumonia (severe) [c]</li> <li>– Complicated severe acute malnutrition [c]</li> <li>– Exacerbations of COPD</li> <li>– Otitis media</li> <li>– Pharyngitis</li> <li>– Progressive apical dental abscess</li> <li>– Sepsis in neonates and children [c]</li> <li>– Sinusitis</li> <li>– Uncomplicated severe acute malnutrition [c]</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Acute bacterial meningitis</li> </ul>
amoxicillin + clavulanic acid	<b>Powder for injection:</b> 500 mg (as sodium) + 100 mg (as potassium salt); 1000 mg (as sodium) + 200 mg (as potassium salt) in vial. <b>Powder for oral liquid:</b> 125 mg (as trihydrate) + 31.25 mg (as potassium salt)/5 mL; 250 mg (as trihydrate) + 62.5 mg (as potassium salt)/5 mL [c]. <b>Tablet:</b> 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt). <b>Tablet (dispersible):</b> 200 mg (as trihydrate) + 28.5 mg (as potassium salt) [c]; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) [c].	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Community acquired pneumonia (severe) [c]</li> <li>– Complicated intraabdominal infections (mild to moderate)</li> <li>– Exacerbations of COPD</li> <li>– Hospital acquired pneumonia</li> <li>– Low-risk febrile neutropenia</li> <li>– Lower urinary tract infections</li> <li>– Sinusitis</li> <li>– Skin and soft tissue infections</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Bone and joint infections</li> <li>– Community-acquired pneumonia (mild to moderate)</li> <li>– Community acquired pneumonia (severe)</li> <li>– Otitis media</li> <li>– Surgical prophylaxis</li> </ul>



## WHO Model List of Essential Medicines – 24th List (2025)

ampicillin	<b>Powder for injection:</b> 500 mg; 1 g (as sodium) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated intraabdominal infections</i> [c]</li> <li>– <i>Complicated severe acute malnutrition</i> [c]</li> <li>– <i>Sepsis in neonates and children</i> [c]</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> </ul>
benzathine benzylpenicillin	<b>Powder for injection:</b> 1.2 million IU ( $\approx$ 900 mg) in vial [c]; 2.4 million IU ( $\approx$ 1.8 g) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Syphilis</i></li> </ul>	<b>SECOND CHOICE</b>
benzylpenicillin	<b>Powder for injection:</b> 600 mg (= 1 million IU); 3 g (= 5 million IU) (sodium or potassium salt) in vial.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (severe)</i> [c]</li> <li>– <i>Complicated severe acute malnutrition</i> [c]</li> <li>– <i>Sepsis in neonates and children</i> [c]</li> <li>– <i>Syphilis</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> </ul>
cefalexin	<b>Powder for oral liquid:</b> 125 mg/5 mL; 250 mg/5 mL (anhydrous). <b>Solid oral dosage form:</b> 250 mg; 500 mg (as monohydrate). <b>Tablet (dispersible):</b> 125 mg [c]; 250 mg [c].	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Skin and soft tissue infections</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Exacerbations of COPD</i></li> <li>– <i>Pharyngitis</i></li> </ul>
cefazolin <sup>a</sup>	<b>Powder for injection:</b> 1 g (as sodium salt) in vial. <sup>a</sup> > 1 month.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Surgical prophylaxis</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Bone and joint infections</i></li> </ul>
chloramphenicol	<b>Oily suspension for injection*:</b> 0.5 g/mL (as sodium succinate) in 2 mL ampoule. *Only for the presumptive treatment of epidemic meningitis in children older than 2 years and in adults. <b>Powder for injection:</b> 1 g (as sodium succinate) in vial.	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> </ul>

## WHO Model List of Essential Medicines – 24th List (2025)

clindamycin	<b>Capsule:</b> 150 mg (as hydrochloride). <b>Injection:</b> 150 mg/mL (as phosphate); 600 mg/4 mL (as phosphate); 900 mg/6 mL (as phosphate). <b>Powder for oral liquid:</b> 75 mg/5 mL (as palmitate hydrochloride) [c].	
	<b>FIRST CHOICE</b> – <i>Necrotizing fasciitis</i>	<b>SECOND CHOICE</b> – <i>Bone and joint infections</i>
<input type="checkbox"/> cloxacillin*  Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (J01CF Beta-lactamase resistant penicillins)	<b>Capsule:</b> 250 mg [c], 500 mg; 1 g (as sodium). <b>Powder for injection:</b> 250 mg [c], 500 mg (as sodium) in vial. <b>Powder for oral liquid:</b> 125 mg/5 mL, 250 mg/5 mL (as sodium) [c]. *cloxacillin, dicloxacillin and flucloxacillin are preferred for oral administration due to better bioavailability.	
	<b>FIRST CHOICE</b> – <i>Bone and joint infections</i> – <i>Skin and soft tissue infections</i>	<b>SECOND CHOICE</b> – <i>Sepsis in neonates and children</i> [c]
doxycycline <input type="checkbox"/> a	<b>Oral liquid:</b> 50 mg/5 mL (calcium) [c]. <b>Powder for oral liquid:</b> 25 mg/5 mL (monohydrate) [c]. <b>Powder for injection:</b> 100 mg in vial. <b>Solid oral dosage form:</b> 50 mg [c]; 100 mg (as hyclate). <b>Tablet (dispersible):</b> 100 mg (as monohydrate) [c]. <input type="checkbox"/> a Use in children <8 years only for life-threatening infections when no alternative exists.	
	<b>FIRST CHOICE</b> – <i>Cholera</i> – <i>Sexually transmitted infection due to Chlamydia trachomatis</i>	<b>SECOND CHOICE</b> – <i>Cholera</i> [c] – <i>Community acquired pneumonia (mild to moderate)</i> – <i>Exacerbations of COPD</i>
gentamicin	<b>Injection:</b> 10 mg/mL (as sulfate); 40 mg/mL (as sulfate) in 2 mL vial.	
	<b>FIRST CHOICE</b> – <i>Acute bacterial meningitis in neonates</i> [c] – <i>Community acquired pneumonia (severe)</i> [c] – <i>Complicated intraabdominal infections</i> [c] – <i>Complicated severe acute malnutrition</i> [c] – <i>Sepsis in neonates and children</i> [c]	<b>SECOND CHOICE</b> – <i>Gonorrhoea</i> – <i>Surgical prophylaxis</i>

## WHO Model List of Essential Medicines – 24th List (2025)

metronidazole	<b>Injection:</b> 500 mg in 100 mL vial. <b>Oral liquid:</b> 200 mg/5 mL (as benzoate). <b>Suppository:</b> 500 mg; 1 g. <b>Tablet:</b> 200 mg; 250 mg; 400 mg; 500 mg.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>C. difficile</i> infection</li> <li>– Complicated intraabdominal infections (mild to moderate)</li> <li>– Complicated intrabdominal infections (severe)</li> <li>– Necrotizing fasciitis</li> <li>– Surgical prophylaxis</li> <li>– Trichomoniasis</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Complicated intraabdominal infections (mild to moderate)</li> </ul>
nitrofurantoin	<b>Oral liquid:</b> 25 mg/5 mL [c]. <b>Solid oral dosage form:</b> 50 mg [c]; 100 mg.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Lower urinary tract infections</li> </ul>	<b>SECOND CHOICE</b>
phenoxymethylpenicillin	<b>Powder for oral liquid:</b> 250 mg/5 mL (as potassium). <b>Solid oral dosage form:</b> 250 mg; 500 mg (as potassium).	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Community acquired pneumonia (mild to moderate)</li> <li>– Pharyngitis</li> <li>– Progressive apical dental abscess</li> </ul>	<b>SECOND CHOICE</b>
procaine benzylpenicillin*	<b>Powder for injection:</b> 1 g (=1 million IU); 3 g (=3 million IU) in vial. *Procaine benzylpenicillin is not recommended as first-line treatment for neonatal sepsis except in settings with high neonatal mortality, when given by trained health workers in cases where hospital care is not achievable.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Syphilis (congenital) [c]</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Syphilis</li> </ul>
spectinomycin	<b>Powder for injection:</b> 2 g (as hydrochloride) in vial.	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Gonorrhoea</li> </ul>
sulfamethoxazole + trimethoprim	<b>Injection:</b> 80 mg + 16 mg/mL in 5 mL ampoule; 80 mg + 16 mg/mL in 10 mL ampoule. <b>Oral liquid:</b> 200 mg + 40 mg/5 mL. <b>Tablet:</b> 100 mg + 20 mg; 400 mg + 80 mg; 800 mg + 160 mg. <b>Tablet (dispersible):</b> 100 mg + 20 mg [c].	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– Lower urinary tract infections</li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– Acute invasive bacterial diarrhoea / dysentery</li> </ul>

## WHO Model List of Essential Medicines – 24th List (2025)

trimethoprim	<b>Tablet:</b> 100 mg; 200 mg. <b>Oral liquid:</b> 50 mg/5 mL [c].	
	<b>FIRST CHOICE</b> – Lower urinary tract infections	<b>SECOND CHOICE</b>
<b>6.2.2 Watch group antibiotics</b>		
azithromycin	<b>Solid oral dosage form:</b> 250 mg; 500 mg (anhydrous). <b>Powder for oral liquid:</b> 200 mg/5 mL (anhydrous) [c].	
	<b>FIRST CHOICE</b> – Cholera – Enteric fever – Gonorrhoea – Sexually transmitted infection due to <i>Chlamydia trachomatis</i> – Trachoma – Yaws	<b>SECOND CHOICE</b> – Acute invasive bacterial diarrhoea / dysentery – Gonorrhoea
cefixime	<b>Powder for oral liquid:</b> 100 mg/5 mL [c]. <b>Solid oral dosage form:</b> 200 mg; 400 mg (as trihydrate).	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> – Acute invasive bacterial diarrhoea / dysentery – Gonorrhoea
cefotaxime*	<b>Powder for injection:</b> 250 mg; 500 mg; 1 g; 2 g (as sodium) in vial. *3rd generation cephalosporin of choice for use in hospitalized neonates.	
	<b>FIRST CHOICE</b> – Acute bacterial meningitis – Community acquired pneumonia (severe) – Complicated intraabdominal infections (mild to moderate) – Complicated intraabdominal infections (severe) – Hospital acquired pneumonia – Pyelonephritis or prostatitis (severe)	<b>SECOND CHOICE</b> – Bone and joint infections – Pyelonephritis or prostatitis (mild to moderate) – Sepsis in neonates and children [c]

## WHO Model List of Essential Medicines – 24th List (2025)

ceftriaxone* <b>a</b>	<b>Powder for injection:</b> 250 mg; 500 mg; 1 g; 2 g (as sodium) in vial. *Do not administer with calcium and avoid in infants with hyperbilirubinaemia. <b>a</b> > 41 weeks corrected gestational age.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute bacterial meningitis</i></li> <li>– <i>Community acquired pneumonia (severe)</i></li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> <li>– <i>Complicated intraabdominal infections (severe)</i></li> <li>– <i>Endophthalmitis</i></li> <li>– <i>Enteric fever</i></li> <li>– <i>Gonorrhoea</i></li> <li>– <i>Hospital acquired pneumonia</i></li> <li>– <i>Necrotizing fasciitis</i></li> <li>– <i>Pyelonephritis or prostatitis (severe)</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute invasive bacterial diarrhoea / dysentery</i></li> <li>– <i>Bone and joint infections</i></li> <li>– <i>Pyelonephritis or prostatitis (mild to moderate)</i></li> <li>– <i>Sepsis in neonates and children [c]</i></li> </ul>
cefuroxime	<b>Powder for injection:</b> 250 mg; 750 mg; 1.5 g (as sodium) in vial.	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Surgical prophylaxis</i></li> </ul>
ciprofloxacin	<b>Oral liquid:</b> 250 mg/5 mL (anhydrous) <b>[c]</b> . <b>Solution for IV infusion:</b> 2 mg/mL (as hyclate) <b>[c]</b> . <b>Solid oral dosage form:</b> 100 mg <b>[c]</b> ; 250 mg; 500 mg (as hydrochloride).	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Acute invasive bacterial diarrhoea / dysentery</i></li> <li>– <i>Enteric fever</i></li> <li>– <i>Low-risk febrile neutropenia</i></li> <li>– <i>Pyelonephritis or prostatitis (mild to moderate)</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Cholera</i></li> <li>– <i>Complicated intraabdominal infections (mild to moderate)</i></li> </ul>
<input type="checkbox"/> clarithromycin† Therapeutic alternatives: - erythromycin* *as second choice treatment for pharyngitis in children (EMLc only)	<b>Powder for oral liquid:</b> 125 mg/5 mL; 250 mg/5 mL. <b>Powder for injection:</b> 500 mg in vial. <b>Solid oral dosage form:</b> 250 mg <b>[c]</b> ; 500 mg. †clarithromycin is also listed for use in combination regimens for eradication of <i>H. pylori</i> in adults.	
	<b>FIRST CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Community acquired pneumonia (severe)</i></li> </ul>	<b>SECOND CHOICE</b> <ul style="list-style-type: none"> <li>– <i>Pharyngitis</i></li> </ul>

## WHO Model List of Essential Medicines – 24th List (2025)

piperacillin + tazobactam	<b>Powder for injection:</b> 2 g (as sodium) + 250 mg (as sodium); 4 g (as sodium) + 500 mg (as sodium) in vial.	
	<b>FIRST CHOICE</b> – <i>Complicated intraabdominal infections (severe)</i> – <i>High-risk febrile neutropenia</i> – <i>Hospital acquired pneumonia</i> – <i>Necrotizing fasciitis</i>	<b>SECOND CHOICE</b>
vancomycin*	<b>Capsule:</b> 125 mg; 250 mg (as hydrochloride). *vancomycin powder for injection may also be used for oral administration	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> – <i>C. difficile infection</i>
<b>Complementary List</b>		
ceftazidime	<b>Powder for injection:</b> 250 mg; 1 g (as pentahydrate) in vial.	
	<b>FIRST CHOICE</b> – <i>Endophthalmitis</i>	<b>SECOND CHOICE</b>
<input type="checkbox"/> meropenem* <sup>[a]</sup>  <i>Therapeutic alternatives*:</i> – imipenem + cilastatin  <i>*complicated intraabdominal infections and high-risk febrile neutropenia only. Meropenem is the preferred choice for acute bacterial meningitis in neonates.</i>	<b>Powder for injection:</b> 500 mg (as trihydrate); 1 g (as trihydrate) in vial. <sup>[a]</sup> > 3 months.	
	<b>FIRST CHOICE</b>	<b>SECOND CHOICE</b> – <i>Acute bacterial meningitis in neonates [c]</i> – <i>Complicated intraabdominal infections (severe)</i> – <i>High-risk febrile neutropenia</i>
vancomycin	<b>Powder for injection:</b> 250 mg; 500 mg; 1 g (as hydrochloride) in vial.	
	<b>FIRST CHOICE</b> – <i>Endophthalmitis</i> – <i>Necrotizing fasciitis</i>	<b>SECOND CHOICE</b> – <i>High-risk febrile neutropenia</i>
<b>6.2.3 Reserve group antibiotics</b>		
<b>Complementary List</b>		
cefiderocol	<b>Powder for injection:</b> 1 g (as sulfate toxylate) in vial.	
ceftazidime + avibactam	<b>Powder for injection:</b> 2 g + 0.5 g in vial.	
ceftolozane + tazobactam	<b>Powder for injection:</b> 1 g + 0.5 g in vial.	
colistin	<b>Powder for injection:</b> 1 million IU (as colistemetate sodium) (equivalent to 34 mg colistin base activity) in vial.	
fosfomycin	<b>Powder for injection:</b> 2 g; 4 g (as sodium) in vial.	
<input type="checkbox"/> linezolid  <i>Therapeutic alternatives:</i> – tedizolid phosphate	<b>Injection for intravenous administration:</b> 2 mg/mL in 300 mL bag.  <b>Powder for oral liquid:</b> 100 mg/5 mL.  <b>Tablet:</b> 600 mg.  <b>Tablet (dispersible):</b> 150 mg [c].	

## WHO Model List of Essential Medicines – 24th List (2025)

<i>meropenem + vaborbactam</i>	<b>Powder for injection:</b> 1 g (as trihydrate) + 1 g in vial.
<i>plazomicin</i>	<b>Injection:</b> 500 mg/10 mL.
<i>polymyxin B</i>	<b>Powder for injection:</b> 500 000 IU (equivalent to 50 mg polymyxin B base) in vial.
<b>6.2.4 Antileprosy medicines</b>	
Medicines used in the treatment of leprosy should never be used except in combination. Combination therapy is essential to prevent the emergence of drug resistance. Colour-coded blister packs (MDT blister packs) containing standard two-medicine (paucibacillary leprosy) or three-medicine (multibacillary leprosy) combinations for adult and childhood leprosy should be used. MDT blister packs can be supplied free of charge through WHO.	
clofazimine	<b>Solid oral dosage form:</b> 50 mg; 100 mg.
dapsone	<b>Tablet:</b> 25 mg; 50 mg; 100 mg.
rifampicin	<b>Oral liquid:</b> 20 mg/mL [c]. <b>Solid oral dosage form:</b> 150 mg; 300 mg.
<b>6.2.5 Antituberculosis medicines</b>	
WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.	
amikacin	<b>Injection:</b> 250 mg/mL (as sulfate) in 2 mL vial.
amoxicillin + clavulanic acid*	<b>Powder for oral liquid:</b> 250 mg (as trihydrate) + 62.5 mg (as potassium salt)/5 mL [c]. <b>Tablet:</b> 500 mg (as trihydrate) + 125 mg (as potassium salt). <b>Tablet (dispersible):</b> 250 mg (as trihydrate) + 62.5 mg (as potassium salt) [c]. *For use only in combination with meropenem or imipenem+cilastatin.
bedaquiline	<b>Tablet:</b> 20 mg [c]; 100 mg.
clofazimine	<b>Solid oral dosage form:</b> 50 mg; 100 mg.
<input type="checkbox"/> cycloserine Therapeutic alternatives: - terizidone	<b>Solid oral dosage form:</b> 125 mg [c]; 250 mg.
delamanid	<b>Tablet (dispersible):</b> 25 mg [c]. <b>Tablet:</b> 50 mg.
ethambutol	<b>Tablet:</b> 100 mg; 400 mg (hydrochloride). <b>Tablet (dispersible):</b> 100 mg [c]
ethambutol + isoniazid + pyrazinamide + rifampicin	<b>Tablet:</b> 275 mg + 75 mg + 400 mg + 150 mg.
ethambutol + isoniazid + rifampicin	<b>Tablet:</b> 275 mg + 75 mg + 150 mg.
<input type="checkbox"/> ethionamide Therapeutic alternatives*: - prothionamide *for multidrug-resistant tuberculosis	<b>Tablet:</b> 250 mg. <b>Tablet (dispersible):</b> 125 mg [c].
isoniazid	<b>Tablet:</b> 100 mg; 300 mg. <b>Tablet (dispersible, scored):</b> 100 mg [c].

## WHO Model List of Essential Medicines – 24th List (2025)

isoniazid + pyrazinamide + rifampicin	<b>Tablet (dispersible):</b> 50 mg + 150 mg + 75 mg [c].
isoniazid + rifampicin	<b>Tablet:</b> 75 mg + 150 mg; 150 mg + 300 mg. <b>Tablet (dispersible):</b> 50 mg + 75 mg [c].
isoniazid + rifapentine	<b>Tablet (scored):</b> 300 mg + 300 mg.
levofloxacin	<b>Tablet:</b> 250 mg; 500 mg; 750 mg. <b>Tablet (dispersible):</b> 100 mg [c].
linezolid	<b>Tablet:</b> 600 mg. <b>Tablet (dispersible, scored):</b> 150 mg [c].
<input type="checkbox"/> meropenem Therapeutic alternatives: - imipenem + cilastatin	<b>Powder for injection:</b> 500 mg (as trihydrate); 1 g (as trihydrate) in vial.
moxifloxacin	<b>Tablet:</b> 400 mg. <b>Tablet (dispersible):</b> 100 mg [c].
p-aminosalicylate sodium	<b>Powder for oral solution:</b> 5.52 g in sachet (equivalent to 4 g p-aminosalicylic acid).
pretomanid	<b>Tablet:</b> 200 mg.
pyrazinamide	<b>Tablet:</b> 400 mg; 500 mg <b>Tablet (dispersible):</b> 150 mg.
rifabutin	<b>Solid oral dosage form:</b> 150 mg.*
rifampicin	<b>Oral liquid:</b> 20 mg/mL [c]. <b>Solid oral dosage form:</b> 150 mg; 300 mg.
rifapentine	<b>Tablet:</b> 150 mg; 300 mg. <b>Tablet (dispersible, scored):</b> 150 mg [c].
streptomycin [c]	<b>Powder for injection:</b> 1 g (as sulfate) in vial.
<b>6.3 Antifungal medicines</b>	
amphotericin B*	<b>Powder for injection:</b> 50 mg (liposomal complex) in vial. <b>Powder for injection:</b> 50 mg (as sodium deoxycholate) in vial *Liposomal amphotericin B has a better safety profile than the sodium deoxycholate formulation and should be prioritized for selection and use depending on local availability and cost.
clotrimazole	<b>Vaginal cream:</b> 1%; 10%. <b>Vaginal tablet:</b> 100 mg; 500 mg.
fluconazole	<b>Capsule:</b> 50 mg. <b>Injection:</b> 2 mg/mL in vial. <b>Oral liquid:</b> 50 mg/5 mL. <b>Powder for oral liquid:</b> 50 mg/5 mL [c].
flucytosine	<b>Capsule:</b> 250 mg. <b>Infusion:</b> 2.5 g in 250 mL.



## WHO Model List of Essential Medicines – 24th List (2025)

griseofulvin	<b>Oral liquid:</b> 125 mg/5 mL [c]. <b>Solid oral dosage form:</b> 125 mg; 250 mg.
itraconazole*	<b>Capsule:</b> 100 mg. <b>Oral liquid:</b> 10 mg/mL. *For treatment of chronic pulmonary aspergillosis, histoplasmosis, sporotrichosis, paracoccidioidomycosis, mycoses caused by <i>T. marneffe</i> and chromoblastomycosis; and prophylaxis of histoplasmosis and infections caused by <i>T. marneffe</i> in AIDS patients.
nystatin	<b>Lozenge:</b> 100 000 IU. <b>Oral liquid:</b> 100 000 IU/mL [c]. <b>Pessary:</b> 100 000 IU. <b>Solid oral dosage form:</b> 500 000 IU.
voriconazole*	<b>Tablet:</b> 50 mg; 200 mg <b>Powder for injection:</b> 200 mg in vial <b>Powder for oral liquid:</b> 40 mg/mL *For treatment of chronic pulmonary aspergillosis and acute invasive aspergillosis.
<b>Complementary List</b>	
<input type="checkbox"/> micafungin Therapeutic alternatives: - anidulafungin - caspofungin	<b>Powder for injection:</b> 50 mg (as sodium); 100 mg (as sodium) in vial.
potassium iodide	<b>Saturated solution.</b>
<b>6.4 Antiviral medicines</b>	
<b>6.4.1 Antiherpes medicines</b>	
<input type="checkbox"/> aciclovir Therapeutic alternatives: - valaciclovir (oral)	<b>Oral liquid:</b> 200 mg/5 mL [c]. <b>Powder for solution for infusion:</b> 250 mg (as sodium dihydrate) in vial. <b>Solution for infusion:</b> 25 mg/mL (as sodium) in vial. <b>Tablet:</b> 200 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>6.4.2 Antiretrovirals</b>	
Based on current evidence and experience of use, medicines in the following classes of antiretrovirals are included as essential medicines for treatment and prevention of HIV (prevention of mother-to-child transmission, pre-exposure prophylaxis (where indicated) and post-exposure prophylaxis). WHO emphasizes the importance of using these products in accordance with global and national guidelines. WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.	
Scored tablets can be used in children and therefore can be considered for inclusion in the listing of tablets, provided that adequate quality products are available.	
<b>6.4.2.1 Nucleoside/Nucleotide reverse transcriptase inhibitors</b>	
abacavir	<b>Tablet:</b> 300 mg (as sulfate).
lamivudine	<b>Oral liquid:</b> 50 mg/5 mL [c]. <b>Tablet:</b> 150 mg.
tenofovir disoproxil fumarate†	<b>Tablet:</b> 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil). †also indicated for pre-exposure prophylaxis.
zidovudine	<b>Capsule:</b> 250 mg. <b>Oral liquid:</b> 50 mg/5 mL. <b>Solution for IV infusion:</b> 10 mg/mL in 20 mL vial. <b>Tablet:</b> 300 mg.
<b>6.4.2.2 Non-nucleoside reverse transcriptase inhibitors</b>	
efavirenz	<b>Tablet:</b> 600 mg.
nevirapine <sup>a</sup>	<b>Oral liquid:</b> 50 mg/5 mL. <b>Tablet (dispersible):</b> 50 mg; 200 mg. <sup>a</sup> > 6 weeks
<b>6.4.2.3 Protease inhibitors</b>	
Selection of protease inhibitor(s) from the Model List will need to be determined by each country after consideration of international and national treatment guidelines and experience. Ritonavir is recommended for use in combination as a pharmacological booster, and not as an antiretroviral in its own right. All other protease inhibitors should be used in boosted forms (e.g. with ritonavir).	
atazanavir + ritonavir	<b>Tablet (heat stable):</b> 300 mg (as sulfate) + 100 mg.
darunavir <sup>a</sup>	<b>Tablet:</b> 75 mg; 400 mg; 600 mg; 800 mg <sup>a</sup> > 3 years
lopinavir + ritonavir	<b>Solid oral dosage form:</b> 40 mg + 10 mg [c]. <b>Tablet (heat stable):</b> 100 mg + 25 mg; 200 mg + 50 mg.
ritonavir	<b>Tablet (heat stable):</b> 25 mg; 100 mg.
<b>6.4.2.4 Integrase inhibitors</b>	
dolutegravir <sup>a</sup>	<b>Tablet (dispersible, scored):</b> 10 mg [c]. <sup>a</sup> ≥ 4 weeks and ≥ 3 kg <b>Tablet:</b> 50 mg <sup>a</sup> ≥ 25 kg

## WHO Model List of Essential Medicines – 24th List (2025)

raltegravir*	<p>Granules for oral suspension: 100 mg in sachet.</p> <p>Tablet (chewable): 25 mg.</p> <p>Tablet: 400 mg.</p> <p>*For use in pregnant women and in second-line regimens in accordance with WHO treatment guidelines.</p>
<b>6.4.2.5 Fixed-dose combinations of antiretroviral medicines</b>	
abacavir + dolutegravir + lamivudine [c]	Tablet (dispersible): 60 mg (as sulfate) + 5 mg + 30 mg.
abacavir + lamivudine	Tablet (dispersible, scored): 120 mg (as sulfate) + 60 mg.
dolutegravir + lamivudine + tenofovir	Tablet: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil)
efavirenz + □ emtricitabine + tenofovir Therapeutic alternatives: - lamivudine (for emtricitabine)	Tablet: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil).
efavirenz + lamivudine + tenofovir	Tablet: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil)
□ emtricitabine + tenofovir† Therapeutic alternatives: - lamivudine (for emtricitabine)	Tablet: 200 mg + 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil). † combination also indicated for pre-exposure prophylaxis
lamivudine + zidovudine	Tablet: 30 mg + 60 mg [c]; 150 mg + 300 mg.
<b>6.4.2.6 Medicines for prevention of HIV-related opportunistic infections</b>	
isoniazid + pyridoxine + sulfamethoxazole + trimethoprim	Tablet (scored): 300 mg + 25 mg + 800 mg + 160 mg
<b>6.4.3 Other antivirals</b>	
valganciclovir*	<p>Tablet: 450 mg (as hydrochloride).</p> <p>*For the treatment of cytomegalovirus retinitis (CMVr).</p>
<b>Complementary list</b>	
oseltamivir*	<p>Capsule: 30 mg; 45 mg; 75 mg (as phosphate).</p> <p>Powder for oral liquid: 6 mg/mL (as phosphate) [c].</p> <p>*Severe illness due to confirmed or suspected influenza virus infection in critically ill hospitalized patients</p>
valganciclovir*[c]	<p>Powder for oral solution: 50 mg/mL (as hydrochloride).</p> <p>Tablet: 450 mg (as hydrochloride).</p> <p>*For the treatment of cytomegalovirus retinitis (CMVr).</p>

## WHO Model List of Essential Medicines – 24th List (2025)

<b>6.4.4 Antihepatitis medicines</b>	
<b>6.4.4.1 Medicines for hepatitis B</b>	
<b>6.4.4.1.1 Nucleoside/Nucleotide reverse transcriptase inhibitors</b>	
entecavir	<b>Oral liquid:</b> 0.05 mg/mL <b>Tablet:</b> 0.5 mg; 1 mg
tenofovir disoproxil fumarate	<b>Tablet:</b> 300 mg (tenofovir disoproxil fumarate – equivalent to 245 mg tenofovir disoproxil).
<b>6.4.4.2 Medicines for hepatitis C</b>	
Pangenotypic direct-acting antivirals should be considered as therapeutic alternatives for the purposes of selection and procurement at national level.	
<b>6.4.4.2.1 <input type="checkbox"/> Pangenotypic direct-acting antiviral combinations</b>	
daclatasvir*	<b>Tablet:</b> 30 mg; 60 mg (as dihydrochloride). *Pangenotypic when used in combination with sofosbuvir
daclatasvir + sofosbuvir	<b>Tablet:</b> 60 mg (as dihydrochloride) + 400 mg.
glecaprevir + pibrentasvir	<b>Granules:</b> 50 mg + 20 mg in sachet [c]. <b>Tablet:</b> 100 mg + 40 mg.
ravidasvir*	<b>Tablet:</b> 200 mg. *Pangenotypic when used in combination with sofosbuvir
sofosbuvir*	<b>Granules:</b> 200 mg in sachet [c]. <b>Tablet:</b> 200 mg; 400 mg. *Pangenotypic when used in combination with daclatasvir or ravidasvir
sofosbuvir + velpatasvir	<b>Granules:</b> 150 mg + 37.5 mg; 200 mg + 50 mg in sachet [c]. <b>Tablet:</b> 200 mg + 50 mg [c]; 400 mg + 100 mg.
<b>6.4.4.2.2 Non-pangenotypic direct-acting antiviral combinations</b>	
ledipasvir + sofosbuvir	<b>Tablet:</b> 90 mg + 400 mg.
<b>6.4.4.2.3 Other antivirals for hepatitis C</b>	
ribavirin*	<b>Injection for intravenous administration:</b> 800 mg; 1 g in 10 mL phosphate buffer solution. <b>Solid oral dosage form:</b> 200 mg; 400 mg; 600 mg. *For the treatment of hepatitis C, in combination with direct acting anti-viral medicines
<b>6.5 Antiprotozoal medicines</b>	
<b>6.5.1 Antiamoebic and anti giardiasis medicines</b>	
diloxanide* <input type="checkbox"/> *proposed for deletion in 2027	<b>Tablet:</b> 500 mg (furoate). <input type="checkbox"/> > 25 kg.
<input type="checkbox"/> metronidazole Therapeutic alternatives: - tinidazole	<b>Injection:</b> 500 mg in 100 mL vial. <b>Oral liquid:</b> 200 mg/5 mL (as benzoate). <b>Tablet:</b> 200 mg; 250 mg; 400 mg; 500 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>6.5.2 Antileishmaniasis medicines</b>	
amphotericin B*	<p><b>Powder for injection:</b> 50 mg (liposomal complex) in vial.</p> <p><b>Powder for injection:</b> 50 mg (as sodium deoxycholate) in vial.</p> <p>*Liposomal amphotericin B has a better safety profile than the sodium deoxycholate formulation and should be prioritized for selection and use depending on local availability and cost.</p>
meglumine antimoniate	<b>Injection:</b> 1.5 g/5 mL in 5 mL ampoule.
miltefosine	<b>Solid oral dosage form:</b> 10 mg; 50 mg.
paromomycin	<b>Solution for intramuscular injection:</b> 750 mg of paromomycin base (as sulfate).
sodium stibogluconate	<b>Injection:</b> 100 mg/mL in 30 mL vial.
<b>6.5.3 Antimalarial medicines</b>	
<b>6.5.3.1 Medicines for curative treatment</b>	
Medicines for the treatment of <i>P. falciparum</i> malaria cases should be used in combination. The list currently recommends combinations according to WHO treatment guidelines for malaria.	
artemether	<p><b>Oily injection:</b> 20 mg/mL; 40 mg/mL in 1 mL ampoule.</p> <p>For use in the management of severe malaria.</p>
artemether + lumefantrine	<p><b>Tablet:</b> 20 mg + 120 mg.</p> <p><b>Tablet (dispersible):</b> 20 mg + 120 mg [c].</p>
artesunate	<p><b>Powder for injection:</b> 30 mg; 60 mg; 120 mg in vial.</p> <p>For use in the management of severe malaria.</p> <p><b>Rectal dosage form:</b> 100 mg [c].</p> <p>For pre-referral treatment of severe malaria only.</p>
artesunate + amodiaquine	<b>Tablet:</b> 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg.
artesunate + mefloquine	<b>Tablet:</b> 25 mg + 50 mg (as hydrochloride); 100 mg + 200 mg (as hydrochloride).
artesunate + pyronaridine	<p><b>Granules:</b> 20 mg + 60 mg (tetraphosphate) [c].</p> <p><b>Tablet:</b> 60 mg + 180 mg (tetraphosphate).</p>
artesunate – sulfadoxine + pyrimethamine	<b>Co-packaged scored tablets:</b> artesunate 50 mg [3] and sulfadoxine + pyrimethamine 500 mg + 25 mg [1]
chloroquine	<p><b>Oral liquid:</b> 50 mg/5 mL (base).</p> <p><b>Tablet:</b> 150 mg (base).</p> <p>For use only in the treatment of <i>Plasmodium vivax</i> infection.</p>
dihydroartemisinin + piperaquine	<p><b>Tablet:</b> 20 mg + 160 mg (phosphate); 40 mg + 320 mg (phosphate); 60 mg + 480 mg (phosphate); 80 mg + 640 mg (phosphate).</p> <p><b>Tablet (dispersible):</b> 20 mg + 160 mg (phosphate); 40 mg + 320 mg (phosphate) [c].</p>
primaquine*	<p><b>Tablet:</b> 7.5 mg; 15 mg (as phosphate).</p> <p>For use to reduce the transmission of <i>Plasmodium falciparum</i> and for radical cure of <i>Plasmodium vivax</i> and <i>Plasmodium ovale</i> infections.</p>

## WHO Model List of Essential Medicines – 24th List (2025)

quinine	<b>Solution for infusion:</b> 60 mg/mL [c]; 300 mg/mL (hydrochloride) in 2 mL ampoule.  For use in the management of severe malaria.
<b>6.5.3.2 Medicines for chemoprevention</b>	
amodiaquine – sulfadoxine + pyrimethamine [c]	<b>Co-packaged dispersible tablets:</b>  amodiaquine 75 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 250 mg + 12.5 mg [1].  amodiaquine 76.5 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 250 mg + 12.5 mg [1].  amodiaquine 150 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 500 mg + 25 mg [1].  amodiaquine 153 mg (as hydrochloride) [3] and sulfadoxine + pyrimethamine 500 mg + 25 mg [1].
sulfadoxine + pyrimethamine	<b>Tablet (dispersible):</b> 250 mg + 12.5 mg [c]; 500 mg + 25 mg.
<b>6.5.3.3 Medicines for chemoprophylaxis in travellers</b>	
chloroquine	<b>Oral liquid:</b> 50 mg/5 mL (base).  <b>Tablet:</b> 150 mg (base).  For use only for prophylaxis of <i>Plasmodium vivax</i> infection.
doxycycline [a]	<b>Oral liquid:</b> 50 mg/5 mL (calcium).  <b>Powder for oral liquid:</b> 25 mg/5 mL (monohydrate).  <b>Solid oral dosage form:</b> 50 mg; 100 mg (as hyclate).  <b>Tablet (dispersible):</b> 100 mg (as monohydrate).  [a] > 8 years
mefloquine	<b>Tablet (scored):</b> 250 mg (as hydrochloride).
<b>6.5.4 Antipneumocystosis and antitoxoplasmosis medicines</b>	
pyrimethamine	<b>Tablet:</b> 25 mg.
sulfadiazine	<b>Tablet:</b> 500 mg.
sulfamethoxazole + trimethoprim	<b>Injection:</b> 80 mg + 16 mg/mL in 5 mL ampoule; 80 mg + 16 mg/mL in 10 mL ampoule.  <b>Oral liquid:</b> 200 mg + 40 mg/5 mL [c].  <b>Tablet:</b> 100 mg + 20 mg [c]; 400 mg + 80 mg [c]; 800 mg + 160 mg.  <b>Tablet (dispersible):</b> 100 mg + 20 mg [c].
<b>Complementary List</b>	
pentamidine	<b>Tablet:</b> 200 mg; 300 mg (as isethionate).

## WHO Model List of Essential Medicines – 24th List (2025)

<b>6.5.5 Antitrypanosomal medicines</b>	
<b>6.5.5.1 African trypanosomiasis</b>	
fexinidazole*	<b>Tablet:</b> 600 mg *For the treatment of 1 <sup>st</sup> and 2 <sup>nd</sup> stage of human African trypanosomiasis due to <i>Trypanosoma brucei gambiense</i> and <i>Trypanosoma brucei rhodesiense</i> infection.
<b>Medicines for the treatment of 1<sup>st</sup> stage African trypanosomiasis</b>	
pentamidine*	<b>Powder for injection:</b> 300 mg (as isetionate) in vial. *To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
suramin sodium*	<b>Powder for injection:</b> 1 g in vial. *To be used for the treatment of the initial phase of <i>Trypanosoma brucei rhodesiense</i> infection.
<b>Medicines for the treatment of 2<sup>nd</sup> stage African trypanosomiasis</b>	
eflornithine*	<b>Injection:</b> 200 mg/mL (hydrochloride) in 50 mL bottle. *To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
melarsoprol	<b>Injection:</b> 180 mg/5 mL in 5 mL ampoule (3.6% solution).
nifurtimox*	<b>Tablet (scored):</b> 30 mg; 120 mg. *Only to be used in combination with eflornithine, for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
<b>Complementary List</b>	
<i>melarsoprol [c]</i>	<b>Injection:</b> 180 mg/5 mL in 5 mL ampoule (3.6% solution).
<b>6.5.5.2 American trypanosomiasis</b>	
benznidazole	<b>Tablet:</b> 12.5 mg [c] <b>Tablet (scored):</b> 50 mg; 100 mg.
nifurtimox	<b>Tablet (scored):</b> 30 mg; 120 mg.
<b>6.6 Medicines for ectoparasitic infections</b>	
ivermectin	<b>Tablet:</b> 3 mg.
<b>6.7 Medicines for Ebola virus disease</b>	
ansuvimab	<b>Powder for injection:</b> 400 mg.
atoltivimab + maftivimab + odesivimab	<b>Injection:</b> 241.7 mg + 241.7 mg + 241.7 mg in 14.5 mL vial.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>6.8 Medicines for COVID-19</b>	
<p>WHO recommends that effective and safe therapeutics for prevention and treatment of COVID-19 should be considered as essential medicines in the context of the public health emergency. WHO recommendations are revised and updated regularly in WHO living guidelines for therapeutics for the treatment and prevention of COVID-19.</p> <p>Selection of essential therapeutics for COVID-19 at the national level should be informed by recommendations in these guidelines, and consideration of the latest evidence, epidemiology and national priorities.</p> <p>The latest WHO Therapeutics and COVID-19: living guideline is available online at: <a href="https://app.magicapp.org/#/guideline/nBkO1E">https://app.magicapp.org/#/guideline/nBkO1E</a></p> <p>The latest WHO Drugs to prevent COVID-19: living guideline is available online at: <a href="https://app.magicapp.org/#/guideline/L6RxYL">https://app.magicapp.org/#/guideline/L6RxYL</a></p>	
<b>7. MEDICINES FOR CYSTIC FIBROSIS</b>	
elexacaftor + tezacaftor + ivacaftor	<p><b>Granules:</b> 80 mg + 40 mg + 60 mg; 100 mg + 50 mg + 75 mg in sachet.</p> <p><b>Tablet:</b> 50 mg + 25 mg + 37.5 mg; 100 mg + 50 mg + 75 mg.</p>
ivacaftor	<p><b>Granules:</b> 59.5 mg; 75 mg in sachet.</p> <p><b>Tablet:</b> 75 mg; 150 mg.</p>
<i>Complementary List</i>	
pancreatic enzymes [c]	<p><b>Capsule (modified release)*:</b> 10 000 lipase units + 8000 amylase units + 600 protease units; 25 000 lipase units + 18 000 amylase units + 1000 protease units.</p> <p><i>*Units expressed in Ph.Eur</i></p>
<b>8. IMMUNOMODULATORS AND ANTINEOPLASTICS</b>	
<b>8.1 Immunomodulators for non-malignant disease</b>	
<i>Complementary List</i>	
<p>□ <i>adalimumab*</i></p> <p><i>Therapeutic alternatives*:</i></p> <ul style="list-style-type: none"> <li>- certolizumab pegol</li> <li>- etanercept</li> <li>- golimumab</li> <li>- infliximab</li> </ul> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Injection:</b> 10 mg/0.2 mL [c]; 20 mg/0.2 mL [c]; 20 mg/0.4 mL [c]; 40 mg/0.4 mL; 40 mg/0.8 mL; 80 mg/0.8 mL in pre-filled syringe or pre-filled pen.</p>
azathioprine	<p><b>Oral liquid:</b> 10 mg/mL [c].</p> <p><b>Powder for injection:</b> 50 mg [c]; 100 mg (as sodium salt) in vial.</p> <p><b>Tablet:</b> 25 mg [c].</p> <p><b>Tablet (scored):</b> 50 mg.</p>
ciclosporin	<p><b>Capsule:</b> 25 mg.</p> <p><b>Concentrate for injection:</b> 50 mg/mL in 1 mL ampoule.</p> <p><b>Oral liquid:</b> 100 mg/mL [c].</p>
tacrolimus	<p><b>Capsule (immediate-release):</b> 0.5 mg; 0.75 mg; 1 mg; 2 mg; 5 mg.</p> <p><b>Granules for oral suspension:</b> 0.2 mg; 1 mg.</p> <p><b>Injection:</b> 5 mg/mL in 1 mL vial.</p>



## WHO Model List of Essential Medicines – 24th List (2025)

8.2 Antineoplastics and supportive medicines	
Medicines listed below should be used according to protocols for treatment of the diseases.	
8.2.1 Cytotoxic medicines	
Complementary List	
<i>arsenic trioxide</i>	<b>Concentrate for solution for infusion:</b> 1 mg/mL; 2 mg/mL. – Acute promyelocytic leukaemia
<i>asparaginase*</i> <i>*including quality-assured biosimilars</i>	<b>Powder for injection:</b> 10 000 IU in vial. – Acute lymphoblastic leukaemia.
<i>bendamustine</i>	<b>Injection:</b> 45 mg/0.5 mL; 180 mg/2 mL. – Chronic lymphocytic leukaemia – Follicular lymphoma
<i>bleomycin</i>	<b>Powder for injection:</b> 15 000 IU (as sulfate) in vial. – Hodgkin lymphoma – Kaposi sarcoma – Ovarian germ cell tumour – Testicular germ cell tumour
<i>calcium folinate (leucovorin calcium)</i>	<b>Injection:</b> 3 mg/mL in 10 mL ampoule; 7.5 mg/mL in 2 mL ampoule; 10 mg/mL in 5 mL ampoule. <b>Tablet:</b> 5 mg; 15 mg; 25 mg. – Burkitt lymphoma – Early stage colon cancer – Early stage rectal cancer – Gestational trophoblastic neoplasia – Metastatic colorectal cancer – Osteosarcoma
<i>capecitabine</i>	<b>Tablet:</b> 150 mg; 500 mg. – Early stage colon cancer – Early stage rectal cancer – Metastatic breast cancer – Metastatic colorectal cancer
<i>carboplatin</i>	<b>Injection:</b> 50 mg/5 mL; 150 mg/15 mL; 450 mg/45 mL; 600 mg/60 mL. – Cervical cancer – Early stage breast cancer – Epithelial ovarian cancer – Head and neck cancer (as a radio-sensitizer) – Low-grade glioma – Nasopharyngeal cancer – Nephroblastoma (Wilms tumour) – Non-small cell lung cancer – Osteosarcoma – Ovarian germ cell tumour – Retinoblastoma – Testicular germ cell tumour
<i>chlorambucil</i>	<b>Tablet:</b> 2 mg. – Chronic lymphocytic leukaemia

## WHO Model List of Essential Medicines – 24th List (2025)

<i>cisplatin</i>	<p><b>Injection:</b> 10 mg/10 mL; 20 mg/20 mL; 50 mg/50 mL; 100 mg/100 mL.</p> <ul style="list-style-type: none"> <li>– Cervical cancer</li> <li>– Head and neck cancer (as a radio-sensitizer)</li> <li>– Low-grade glioma</li> <li>– Nasopharyngeal cancer (as a radio-sensitizer)</li> <li>– Non-small cell lung cancer</li> <li>– Osteosarcoma</li> <li>– Ovarian germ cell tumour</li> <li>– Testicular germ cell tumour</li> </ul>
<i>cyclophosphamide</i>	<p><b>Powder for injection:</b> 500 mg; 1 g; 2 g in vial.</p> <p><b>Solid oral dosage form:</b> 25 mg; 50 mg.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Anaplastic large cell lymphoma</li> <li>– Burkitt lymphoma</li> <li>– Chronic lymphocytic leukaemia</li> <li>– Diffuse large B-cell lymphoma</li> <li>– Early stage breast cancer</li> <li>– Ewing sarcoma</li> <li>– Follicular lymphoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Hodgkin lymphoma</li> <li>– Low-grade glioma</li> <li>– Metastatic breast cancer</li> <li>– Multiple myeloma</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Rhabdomyosarcoma</li> </ul>
<i>cytarabine</i>	<p><b>Injection:</b> 100 mg/mL in vial.</p> <p><b>Powder for injection:</b> 100 mg in vial.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Acute myeloid leukaemia</li> <li>– Acute promyelocytic leukaemia</li> <li>– Anaplastic large cell lymphoma</li> <li>– Burkitt lymphoma</li> <li>– Langerhans cell histiocytosis</li> </ul>
<i>dacarbazine</i>	<p><b>Powder for injection:</b> 100 mg; 200 mg in vial.</p> <ul style="list-style-type: none"> <li>– Hodgkin lymphoma</li> </ul>
<i>dactinomycin</i>	<p><b>Powder for injection:</b> 500 micrograms in vial.</p> <ul style="list-style-type: none"> <li>– Ewing sarcoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Nephroblastoma (Wilms tumour)</li> <li>– Rhabdomyosarcoma</li> </ul>
<i>daunorubicin</i>	<p><b>Injection:</b> 2 mg/mL; 5 mg/mL (as hydrochloride) in vial.</p> <p><b>Powder for injection:</b> 20 mg; 50 mg (as hydrochloride) in vial.</p> <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Acute myeloid leukaemia</li> <li>– Acute promyelocytic leukaemia</li> </ul>

## WHO Model List of Essential Medicines – 24th List (2025)

<i>docetaxel</i>	<b>Injection:</b> 20 mg/mL; 40 mg/mL. – Early stage breast cancer – Metastatic breast cancer – Metastatic prostate cancer
<i>doxorubicin</i>	<b>Injection:</b> 2 mg/mL (hydrochloride) in vial. <b>Powder for injection:</b> 10 mg; 50 mg (hydrochloride) in vial. – Acute lymphoblastic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Diffuse large B-cell lymphoma – Early stage breast cancer – Ewing sarcoma – Follicular lymphoma – Hodgkin lymphoma – Kaposi sarcoma – Metastatic breast cancer – Multiple myeloma – Nephroblastoma (Wilms tumour) – Osteosarcoma
<i>doxorubicin (as pegylated liposomal)</i>	<b>Injection:</b> 2 mg/mL (hydrochloride) in 10 mL, 25 mL vial – Kaposi sarcoma
<i>etoposide</i>	<b>Capsule:</b> 50 mg, 100 mg. <b>Injection:</b> 20 mg/mL in 5 mL ampoule. <b>Powder for injection:</b> 100 mg (as phosphate) in vial. – Acute lymphoblastic leukaemia – Acute myeloid leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Ewing sarcoma – Gestational trophoblastic neoplasia – Hodgkin lymphoma – Nephroblastoma (Wilms tumour) – Non-small cell lung cancer – Osteosarcoma – Ovarian germ cell tumour – Retinoblastoma – Testicular germ cell tumour
<i>fludarabine</i>	<b>Powder for injection:</b> 50 mg (phosphate) in vial. <b>Tablet:</b> 10 mg – Chronic lymphocytic leukaemia.
<i>fluorouracil</i>	<b>Injection:</b> 50 mg/mL in vial. – Early stage breast cancer – Early stage colon cancer – Early stage rectal cancer – Metastatic colorectal cancer – Nasopharyngeal cancer
<i>gemcitabine</i>	<b>Powder for injection:</b> 200 mg; 1 g in vial. – Epithelial ovarian cancer – Non-small cell lung cancer

## WHO Model List of Essential Medicines – 24th List (2025)

<i>hydroxyurea (hydroxycarbamide)</i>	<p><b>Solid oral dosage form:</b> 100 mg [c]; 200 mg; 300 mg; 400 mg; 500 mg; 1 g.</p> <p>– Chronic myeloid leukaemia</p>
<i>ifosfamide</i>	<p><b>Powder for injection:</b> 500 mg; 1 g; 2 g in vial.</p> <p>– Anaplastic large cell lymphoma – Burkitt lymphoma – Ewing sarcoma – Nephroblastoma (Wilms tumour) – Osteosarcoma – Ovarian germ cell tumour – Rhabdomyosarcoma – Testicular germ cell tumour</p>
<i>irinotecan</i>	<p><b>Injection:</b> 40 mg/2 mL in 2 mL vial; 100 mg/5 mL in 5 mL vial; 500 mg/25 mL in 25 mL vial.</p> <p>– Metastatic colorectal cancer – Nephroblastoma (Wilms tumour) – Rhabdomyosarcoma</p>
<i>melphalan</i>	<p><b>Tablet:</b> 2 mg.</p> <p><b>Powder for injection:</b> 50 mg in vial.</p> <p>– Multiple myeloma</p>
<i>mercaptopurine</i>	<p><b>Tablet:</b> 50 mg.</p> <p><b>Oral liquid:</b> 20 mg/mL [c].</p> <p>– Acute lymphoblastic leukaemia – Acute promyelocytic leukaemia. – Langerhans cell histiocytosis</p>
<i>methotrexate</i>	<p><b>Concentrated injection:</b> 1000 mg/10 mL.</p> <p><b>Injection:</b> 50 mg/2 mL.</p> <p><b>Powder for injection:</b> 50 mg (as sodium) in vial.</p> <p><b>Tablet:</b> 2.5 mg (as sodium).</p> <p>– Acute lymphoblastic leukaemia – Acute promyelocytic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Early stage breast cancer – Gestational trophoblastic neoplasia – Langerhans cell histiocytosis – Osteosarcoma</p>
<i>oxaliplatin</i>	<p><b>Injection:</b> 50 mg/10 mL in 10 mL vial; 100 mg/20 mL in 20 mL vial; 200 mg/40 mL in 40 mL vial.</p> <p><b>Powder for injection:</b> 50 mg; 100 mg in vial.</p> <p>– Early stage colon cancer – Metastatic colorectal cancer</p>

## WHO Model List of Essential Medicines – 24th List (2025)

<i>paclitaxel</i>	<b>Injection:</b> 6 mg/mL in vial. <ul style="list-style-type: none"> <li>– Cervical cancer</li> <li>– Epithelial ovarian cancer</li> <li>– Early stage breast cancer</li> <li>– Metastatic breast cancer</li> <li>– Kaposi sarcoma</li> <li>– Nasopharyngeal cancer</li> <li>– Non-small cell lung cancer</li> <li>– Ovarian germ cell tumour</li> </ul>
<i>pegaspargase*</i> <i>*including quality-assured biosimilars</i>	<b>Injection:</b> 3750 units/5 mL in vial. <b>Powder for injection:</b> 3750 units in vial. <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> </ul>
<i>procarbazine [c]</i>	<b>Capsule:</b> 50 mg (as hydrochloride). <ul style="list-style-type: none"> <li>– Hodgkin lymphoma</li> </ul>
<i>realgar-Indigo naturalis formulation</i>	<b>Tablet:</b> 270 mg (containing tetra-arsenic tetra-sulfide 30 mg). <ul style="list-style-type: none"> <li>– Acute promyelocytic leukaemia</li> </ul>
<i>tioguanine [c]</i>	<b>Solid oral dosage form:</b> 40 mg. <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> </ul>
<i>vinblastine</i>	<b>Injection:</b> 10 mg/10 mL (sulfate) in vial. <b>Powder for injection:</b> 10 mg (sulfate) in vial. <ul style="list-style-type: none"> <li>– Anaplastic large cell lymphoma</li> <li>– Hodgkin lymphoma</li> <li>– Kaposi sarcoma</li> <li>– Langerhans cell histiocytosis</li> <li>– Low-grade glioma</li> <li>– Ovarian germ cell tumour</li> <li>– Testicular germ cell tumour</li> </ul>
<i>vincristine</i>	<b>Injection:</b> 1 mg/mL (sulfate); 2 mg/2 mL (sulfate) in vial. <b>Powder for injection:</b> 1 mg; 5 mg (sulfate) in vial. <ul style="list-style-type: none"> <li>– Acute lymphoblastic leukaemia</li> <li>– Burkitt lymphoma</li> <li>– Diffuse large B-cell lymphoma</li> <li>– Ewing sarcoma</li> <li>– Follicular lymphoma</li> <li>– Gestational trophoblastic neoplasia</li> <li>– Hodgkin lymphoma</li> <li>– Kaposi sarcoma</li> <li>– Langerhans cell histiocytosis</li> <li>– Low-grade glioma</li> <li>– Neuroblastoma (Wilms tumour)</li> <li>– Retinoblastoma</li> <li>– Rhabdomyosarcoma</li> </ul>
<i>vinorelbine</i>	<b>Capsule:</b> 20 mg; 30 mg; 80 mg. <b>Injection:</b> 10 mg/mL in 1 mL, 5 mL vial. <ul style="list-style-type: none"> <li>– Non-small cell lung cancer</li> <li>– Metastatic breast cancer</li> <li>– Rhabdomyosarcoma</li> </ul>

## WHO Model List of Essential Medicines – 24th List (2025)

8.2.2 Targeted therapies	
Complementary List	
<i>all-trans retinoid acid (ATRA)</i>	<b>Capsule:</b> 10 mg. – Acute promyelocytic leukaemia.
<i>bortezomib</i>	<b>Powder for injection:</b> 3.5 mg in vial. – Multiple myeloma
<i>dasatinib</i>	<b>Tablet:</b> 20 mg; 50 mg; 70 mg; 80 mg; 100 mg; 140 mg. – Imatinib-resistant chronic myeloid leukaemia
<input type="checkbox"/> <i>erlotinib</i> Therapeutic alternatives: - <i>afatinib</i> - <i>gefitinib</i>	<b>Tablet:</b> 100 mg, 150 mg. – EGFR mutation-positive advanced non-small cell lung cancer
<i>everolimus</i>	<b>Tablet:</b> 2.5 mg; 5 mg; 7.5 mg; 10 mg. <b>Tablet (dispersible):</b> 2 mg; 3 mg; 5 mg. – Subependymal giant cell astrocytoma
<input type="checkbox"/> <i>ibrutinib</i> Therapeutic alternatives: - <i>zanubrutinib</i>	<b>Capsule:</b> 140 mg. – Relapsed/refractory chronic lymphocytic leukaemia/small lymphocytic lymphoma
<i>imatinib</i>	<b>Solid oral dosage form:</b> 100 mg; 400 mg. – Chronic myeloid leukaemia – Gastrointestinal stromal tumour – Philadelphia chromosome positive acute lymphoblastic leukaemia
<i>nilotinib</i>	<b>Capsule:</b> 150 mg; 200 mg. – Imatinib-resistant chronic myeloid leukaemia
<i>rituximab*</i> <i>*including quality-assured biosimilars</i>	<b>Injection (intravenous):</b> 100 mg/10 mL in 10 mL vial; 500 mg/50 mL in 50 mL vial. – Burkitt lymphoma – Diffuse large B-cell lymphoma – Chronic lymphocytic leukaemia – Follicular lymphoma
<i>trastuzumab*</i> <i>*including quality-assured biosimilars</i>	<b>Powder for injection:</b> 60 mg; 150 mg; 440 mg in vial. – Early stage HER2-positive breast cancer – Metastatic HER2-positive breast cancer

## WHO Model List of Essential Medicines – 24th List (2025)

8.2.3 Immunomodulators	
Complementary List	
<p><i>blinatumomab*</i></p> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Powder for concentrate for solution for infusion:</b> 35 micrograms; 38.5 micrograms in vial.</p> <p>– B-cell acute lymphoblastic leukemia</p>
<p><i>filgrastim*</i></p> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Injection:</b> 120 micrograms/0.2 mL; 300 micrograms/0.5 mL; 480 micrograms/0.8 mL in pre-filled syringe.</p> <p><b>Injection:</b> 300 micrograms/mL in 1 mL vial; 480 micrograms/1.6 mL in 1.6 mL vial.</p> <p>– Primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy.</p> <p>– Secondary prophylaxis for patients who have experienced neutropenia following prior myelotoxic chemotherapy</p> <p>– To facilitate administration of dose dense chemotherapy regimens</p>
<p><i>lenalidomide</i></p>	<p><b>Capsule:</b> 25 mg.</p> <p>– Multiple myeloma</p>
<p><i>pegfilgrastim*</i></p> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Injection:</b> 6 mg/0.6 mL in pre-filled syringe.</p> <p>– Primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy</p> <p>– Secondary prophylaxis for patients who have experienced neutropenia following prior myelotoxic chemotherapy</p> <p>– To facilitate administration of dose dense chemotherapy regimens</p>
<p><i>pembrolizumab*</i></p> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Concentrate for solution for infusion:</b> 25 mg/mL in 4 mL vial.</p> <p>– Metastatic cervical cancer <math>\geq 1\%</math> PD-L1 expression<sup>^</sup></p> <p><sup>^</sup>in combination with platinum-based chemotherapy</p> <p>– Metastatic colorectal cancer<sup>#</sup></p> <p><sup>#</sup>as monotherapy for deficient mismatch repair (dMMR) / microsatellite instability-high (MSI-H) tumours</p>
<p><input type="checkbox"/> <i>pembrolizumab*</i></p> <p>Therapeutic alternatives*:</p> <p>– atezolizumab*</p> <p>– cemiplimab*</p> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Concentrate for solution for infusion:</b> 25 mg/mL in 4 mL vial.</p> <p>– Metastatic non-small cell lung cancer, oncogene-driver wild-type and <math>\geq 50\%</math> PD-L1 expression<sup>^</sup></p> <p><sup>^</sup>as monotherapy</p>
<p><input type="checkbox"/> <i>pembrolizumab*</i></p> <p>Therapeutic alternatives*:</p> <p>– nivolumab*</p> <p><i>*including quality-assured biosimilars</i></p>	<p><b>Concentrate for solution for infusion:</b> 25 mg/mL in 4 mL vial.</p> <p>– Metastatic melanoma<sup>^</sup></p> <p><sup>^</sup> as monotherapy</p>
<p><i>thalidomide</i></p>	<p><b>Capsule:</b> 50 mg.</p> <p>– Multiple myeloma</p>

## WHO Model List of Essential Medicines – 24th List (2025)

8.2.4 Hormones and antihormones	
Complementary List	
<input type="checkbox"/> <i>abiraterone</i> Therapeutic alternatives: - <i>enzalutamide</i>	<b>Tablet:</b> 250 mg; 500 mg. – Metastatic castration-resistant prostate cancer
<input type="checkbox"/> <i>anastrozole</i> Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (L02BG Aromatase inhibitors)	<b>Tablet:</b> 1 mg. – Early stage breast cancer – Metastatic breast cancer
<input type="checkbox"/> <i>bicalutamide</i> Therapeutic alternatives: - <i>flutamide</i> - <i>nilutamide</i>	<b>Tablet:</b> 50 mg. – Metastatic prostate cancer
<i>dexamethasone</i>	<b>Injection:</b> 4 mg/mL dexamethasone phosphate (as sodium phosphate) (equivalent to 3.3 mg/mL dexamethasone base) in 1 mL ampoule. <b>Oral liquid:</b> 2 mg/5 mL (as sodium phosphate) [c]. <b>Tablet:</b> 2 mg [c]; 4 mg (as dexamethasone base). – Acute lymphoblastic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Multiple myeloma
<i>hydrocortisone</i>	<b>Powder for injection:</b> 100 mg (as sodium succinate) in vial. – Acute lymphoblastic leukaemia – Burkitt lymphoma
<input type="checkbox"/> <i>leuprorelin</i> Therapeutic alternatives: - <i>goserelin</i> - <i>triptorelin</i>	<b>Injection:</b> 7.5 mg; 22.5 mg in pre-filled syringe. – Early stage breast cancer – Metastatic prostate cancer.
<i>methylprednisolone [c]</i>	<b>Powder for injection:</b> 40 mg (as sodium succinate); 125 mg (as sodium succinate) in vial. – Acute lymphoblastic leukemia – Burkitt lymphoma
<input type="checkbox"/> <i>prednisolone</i> Therapeutic alternatives: - <i>prednisone</i>	<b>Oral liquid:</b> 5 mg/mL [c]. <b>Tablet:</b> 5 mg; 25 mg. – Acute lymphoblastic leukaemia – Anaplastic large cell lymphoma – Burkitt lymphoma – Chronic lymphocytic leukaemia – Diffuse large B-cell lymphoma – Follicular lymphoma – Hodgkin lymphoma – Langerhans cell histiocytosis – Metastatic castration-resistant prostate cancer – Multiple myeloma



## WHO Model List of Essential Medicines – 24th List (2025)

<i>tamoxifen</i>	<b>Tablet:</b> 10 mg; 20 mg (as citrate). – Early stage breast cancer – Metastatic breast cancer
<b>8.2.5 Supportive medicines</b>	
<b>Complementary List</b>	
<i>allopurinol [c]</i>	<b>Powder for injection:</b> 500 mg (as sodium) <b>Tablet:</b> 100 mg; 300 mg. – Tumour lysis syndrome
<i>mesna</i>	<b>Injection*:</b> 100 mg/mL in 2 mL, 4 mL, 10 mL ampoule. <b>Tablet:</b> 400 mg; 600 mg. *May also be used for oral administration. – Burkitt lymphoma – Ewing sarcoma – Nephroblastoma (Wilms tumour) – Osteosarcoma – Ovarian germ cell tumour – Rhabdomyosarcoma – Testicular germ cell tumour
<i>rasburicase</i>	<b>Powder and solvent for solution for infusion:</b> 1.5 mg (with 1 mL solvent); 7.5 mg (with 5 mL solvent) in vial. – Tumour lysis syndrome
<i>zoledronic acid</i>	<b>Concentrate solution for infusion:</b> 4 mg/5 mL in 5 mL vial. <b>Solution for infusion:</b> 4 mg/100 mL in 100 mL bottle. – Malignancy-related bone disease
<b>9. THERAPEUTIC FOODS</b>	
ready-to-use therapeutic food [c]	<b>Biscuit or paste*.</b> *of nutritional composition as determined by the UN joint statement on the community-based management of severe acute malnutrition and Codex alimentarius guidelines.
<b>10. MEDICINES AFFECTING THE BLOOD</b>	
<b>10.1 Antianaemia medicines</b>	
ferrous salt	<b>Oral liquid:</b> equivalent to 9 mg/mL elemental iron [c]; equivalent to 25 mg/mL elemental iron. <b>Tablet:</b> equivalent to 60 mg – 65 mg elemental iron.
ferrous salt + folic acid	<b>Tablet:</b> equivalent to 60 mg elemental iron + 400 micrograms folic acid.* *nutritional supplement for use during pregnancy. <b>Tablet:</b> equivalent to 60 mg elemental iron + 2.8 mg folic acid.** **for weekly iron and folic acid supplementation.
folic acid	<b>Oral liquid:</b> 1 mg/mL [c]. <b>Tablet:</b> 400 micrograms*; 1 mg; 5 mg. *periconceptual use for prevention of first occurrence of neural tube defects.

## WHO Model List of Essential Medicines – 24th List (2025)

hydroxocobalamin	<b>Injection:</b> 1 mg/mL (as acetate, as hydrochloride or as sulfate) in 1 mL ampoule.
<b>Complementary List</b>	
<input type="checkbox"/> erythropoiesis-stimulating agents* <i>Therapeutic alternatives:</i> - epoetin alfa, beta and theta - darbepoetin alfa - methoxy polyethylene glycol-epoetin beta *including quality-assured biosimilars	<b>Injection: pre-filled syringe</b> 1000 IU/0.5 mL; 2000 IU/0.5 mL; 3000 IU/0.3 mL; 4000 IU/0.4 mL; 5000 IU/0.5 mL; 6000 IU/0.6 mL; 8000 IU/0.8 mL; 10 000 IU/1 mL; 20 000 IU/0.5 mL; 40 000 IU/1 mL.
<b>10.2 Medicines affecting coagulation</b>	
<input type="checkbox"/> dabigatran <i>Therapeutic alternatives:</i> - apixaban - edoxaban - rivaroxaban	<b>Capsule:</b> 110 mg; 150 mg.
desmopressin [c]	<b>Injection:</b> 4 micrograms/mL (acetate) in 1 mL ampoule. <b>Nasal spray:</b> 150 micrograms (acetate) per actuation.
emicizumab	<b>Injection:</b> 12 mg/0.4 mL [c]; 30 mg/mL [c]; 60 mg/0.4 mL; 105 mg/0.7 mL; 150 mg/mL; 300 mg/2 mL in vial.
<input type="checkbox"/> enoxaparin* <i>Therapeutic alternatives*:</i> - dalteparin - nadroparin *including quality-assured biosimilars	<b>Injection: ampoule or pre-filled syringe</b> 20 mg/0.2 mL; 40 mg/0.4 mL; 60 mg/0.6 mL; 80 mg/0.8 mL; 100 mg/1 mL; 120 mg/0.8 mL; 150 mg/1 mL.
heparin sodium	<b>Injection:</b> 1000 IU/mL; 5000 IU/mL; 20 000 IU/mL in 1 mL ampoule or vial.
phytomenadione	<b>Injection:</b> 1 mg/0.5mL [c]; 1 mg/mL [c]; 10 mg/mL in ampoule. <b>Injection (mixed micelle solution):</b> 2 mg/0.2 mL; 10 mg/mL in ampoule. <b>Tablet:</b> 5 mg.
protamine sulfate	<b>Injection:</b> 10 mg/mL in 5 mL ampoule or vial.
tranexamic acid	<b>Injection:</b> 100 mg/mL in 10 mL ampoule.
<input type="checkbox"/> warfarin <i>Therapeutic alternatives:</i> - acenocoumarol	<b>Tablet (scored):</b> 1 mg; 2 mg; 3 mg; 5 mg (sodium).
<b>Complementary List</b>	
heparin sodium [c]	<b>Injection:</b> 1000 IU/mL; 5000 IU/mL in 1 mL ampoule or vial.
protamine sulfate [c]	<b>Injection:</b> 10 mg/mL in 5 mL ampoule or vial.
<input type="checkbox"/> warfarin [c] <i>Therapeutic alternatives:</i> - acenocoumarol	<b>Tablet (scored):</b> 0.5 mg; 1 mg; 2 mg; 3 mg; 5 mg (sodium).

## WHO Model List of Essential Medicines – 24th List (2025)

<b>10.3 Medicines for haemoglobinopathies</b>	
<b>10.3.1 Medicines for sickle-cell disease</b>	
<input type="checkbox"/> deferasirox Therapeutic alternatives: - deferiprone	<b>Tablet (dispersible):</b> 100 mg; 125 mg; 250 mg; 400 mg; 500 mg. <b>Tablet (film-coated):</b> 90 mg; 180 mg; 360 mg.
hydroxyurea (hydroxycarbamide)	<b>Solid oral dosage form:</b> 100 mg [c]; 200 mg; 500 mg; 1 g.
<b>Complementary List</b>	
<i>deferroxamine</i>	<b>Powder for injection:</b> 500 mg (mesilate) in vial.
<b>10.3.2 Medicines for thalassaemias</b>	
<input type="checkbox"/> deferasirox Therapeutic alternatives: - deferiprone	<b>Tablet (dispersible):</b> 100 mg; 125 mg; 250 mg; 400 mg; 500 mg. <b>Tablet (film-coated):</b> 90 mg; 180 mg; 360 mg.
<b>Complementary List</b>	
<i>deferroxamine</i>	<b>Powder for injection:</b> 500 mg (mesilate) in vial.
<b>11. BLOOD PRODUCTS, COAGULATION FACTORS, AND PLASMA SUBSTITUTES</b>	
<b>11.1 Blood and blood components</b>	
<p>In accordance with the World Health Assembly resolution WHA63.12, WHO recognizes that achieving self-sufficiency, unless special circumstances preclude it, in the supply of safe blood components based on voluntary, non-remunerated blood donation, and the security of that supply are important national goals to prevent blood shortages and meet the transfusion requirements of the patient population.</p> <p>All blood and plasma-derived products should comply with the WHO requirements.</p>	
<input type="checkbox"/> cryoprecipitate, pathogen-reduced Therapeutic alternatives: - cryoprecipitate, native* *native cryoprecipitate should only be used in situations of life-threatening haemorrhage when pathogen-reduced cryoprecipitate is not available.	<b>Injection:</b> frozen liquid in bag or lyophilized powder in vial containing: <ul style="list-style-type: none"> <li>- &gt; 50 IU Factor VIII</li> <li>- &gt; 100 IU vWF</li> <li>- &gt; 140 mg clottable fibrinogen per unit</li> </ul>
fresh-frozen plasma	
platelets	
red blood cells	
whole blood	
<b>11.2 Human immunoglobulins</b>	
anti-D immunoglobulin	<b>Injection:</b> 250 micrograms in single-dose vial.
anti-rabies immunoglobulin	<b>Injection:</b> 150 IU/mL in vial.
anti-tetanus immunoglobulin	<b>Injection:</b> 500 IU in vial.

## WHO Model List of Essential Medicines – 24th List (2025)

<i>Complementary List</i>	
<i>normal immunoglobulin</i>	<p><b>Intramuscular administration:</b> 16% protein solution.</p> <p><b>Subcutaneous administration:</b> 15%; 16% protein solution.</p> <ul style="list-style-type: none"> <li>– Primary immune deficiency.</li> </ul> <p><b>Intravenous administration:</b> 5%; 10% protein solution.</p> <ul style="list-style-type: none"> <li>– Primary immune deficiency</li> <li>– Kawasaki disease</li> <li>– Langerhans cell histiocytosis</li> </ul>
<b>11.3 Coagulation factors</b>	
coagulation factor VIII, plasma-derived	<b>Powder for injection:</b> 250 IU; 500 IU; 1000 IU in vial.
coagulation factor IX, plasma-derived	<b>Powder for injection:</b> 500 IU; 1000 IU in vial.
coagulation factor VIII, recombinant	<b>Lyophilized powder for solution for injection:</b> 250 IU, 500 IU, 1000 IU, 1500 IU, 2000 IU, 3000 IU, 4000 IU in vial.
coagulation factor IX, recombinant	<b>Lyophilized powder for solution for injection:</b> 250 IU, 500 IU, 1000 IU, 1500 IU, 2000 IU, 3000 IU, 4000 IU in vial.
<b>11.4 Plasma substitutes</b>	
<input type="checkbox"/> dextran 70 Therapeutic alternatives: - polygeline injectable solution 3.5%	<b>Injectable solution:</b> 6%.
<b>12. CARDIOVASCULAR MEDICINES</b>	
<b>12.1 Antianginal medicines</b>	
<input type="checkbox"/> bisoprolol Therapeutic alternatives: - carvedilol - metoprolol	<b>Tablet:</b> 1.25 mg; 5 mg.
glyceryl trinitrate	<b>Tablet (sublingual):</b> 500 micrograms.
isosorbide dinitrate	<b>Tablet (sublingual):</b> 5 mg.
verapamil	<b>Tablet:</b> 40 mg; 80 mg (hydrochloride).
<b>12.2 Antiarrhythmic medicines</b>	
<input type="checkbox"/> bisoprolol Therapeutic alternatives: - carvedilol - metoprolol	<b>Tablet:</b> 1.25 mg; 5 mg.
digoxin	<p><b>Injection:</b> 250 micrograms/mL in 2 mL ampoule.</p> <p><b>Oral liquid:</b> 50 micrograms/mL.</p> <p><b>Tablet:</b> 62.5 micrograms; 250 micrograms.</p>
epinephrine (adrenaline)	<b>Injection:</b> 100 micrograms/mL (as acid tartrate or hydrochloride) in 10 mL ampoule.
lidocaine	<b>Injection:</b> 20 mg/mL (hydrochloride) in 5 mL ampoule.

## WHO Model List of Essential Medicines – 24th List (2025)

verapamil	<b>Injection:</b> 2.5 mg/mL (hydrochloride) in 2 mL ampoule. <b>Tablet:</b> 40 mg; 80 mg (hydrochloride).
<i>Complementary List</i>	
amiodarone	<b>Injection:</b> 50 mg/mL (hydrochloride) in 3 mL ampoule. <b>Tablet:</b> 100 mg; 200 mg; 400 mg (hydrochloride).
<b>12.3 Antihypertensive medicines</b>	
<input type="checkbox"/> amlodipine Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives)	<b>Tablet:</b> 5 mg (as maleate, mesylate or besylate).
<input type="checkbox"/> bisoprolol Therapeutic alternatives: - atenolol* - carvedilol - metoprolol	<b>Tablet:</b> 1.25 mg; 5 mg.  *atenolol should not be used as a first-line agent in uncomplicated hypertension in patients > 60 years
<input type="checkbox"/> enalapril Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain)	<b>Oral liquid:</b> 1 mg/mL (as hydrogen maleate) [c]. <b>Tablet:</b> 2.5 mg; 5 mg; 10 mg (as hydrogen maleate).
hydralazine*	<b>Powder for injection:</b> 20 mg (hydrochloride) in ampoule. <b>Tablet:</b> 25 mg; 50 mg (hydrochloride).  *Hydralazine is listed for use only in the acute management of severe pregnancy-induced hypertension. Its use in the treatment of essential hypertension is not recommended in view of the evidence of greater efficacy and safety of other medicines.
<input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - chlorothiazide - chlorthalidone - indapamide	<b>Oral liquid:</b> 50 mg/5 mL. <b>Solid oral dosage form:</b> 12.5 mg; 25 mg.
<input type="checkbox"/> lisinopril + <input type="checkbox"/> amlodipine Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for lisinopril) - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	<b>Tablet:</b> 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg.
<input type="checkbox"/> lisinopril + <input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for lisinopril) - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	<b>Tablet:</b> 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

<input type="checkbox"/> losartan Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain)	<b>Tablet:</b> 25 mg; 50 mg; 100 mg.
methyldopa*	<b>Tablet:</b> 250 mg. *Methyldopa is listed for use only in the management of pregnancy-induced hypertension. Its use in the treatment of essential hypertension is not recommended in view of the evidence of greater efficacy and safety of other medicines.
<input type="checkbox"/> perindopril + <input type="checkbox"/> amlodipine + <input type="checkbox"/> indapamide Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for perindopril) - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine) - chlorthalidone, chlorothiazide, hydrochlorothiazide (for indapamide)	<b>Solid oral dosage form:</b> 5 mg (arginine) + 5 mg + 1.25 mg; 5 mg (arginine) + 10 mg + 1.25 mg; 10 mg (arginine) + 5 mg + 2.5 mg; 10 mg (arginine) + 10 mg + 2.5 mg.
<input type="checkbox"/> telmisartan + <input type="checkbox"/> amlodipine Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for telmisartan) - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	<b>Tablet:</b> 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg.
<input type="checkbox"/> telmisartan + <input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for telmisartan) - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	<b>Tablet:</b> 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg.
<input type="checkbox"/> valsartan + <input type="checkbox"/> amlodipine + <input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain) (for valsartan) - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine) - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	<b>Solid oral dosage form:</b> 160 mg + 5 mg + 12.5 mg; 160 mg + 5 mg + 25 mg; 160 mg + 10 mg + 12.5 mg; 160 mg + 10 mg + 25 mg; 320 mg + 10 mg + 25 mg.
<b>Complementary List</b>	
<i>sodium nitroprusside</i>	<b>Powder for infusion:</b> 50 mg in ampoule.
<b>12.4 Medicines used in heart failure</b>	
<input type="checkbox"/> bisoprolol Therapeutic alternatives: - carvedilol - metoprolol	<b>Tablet:</b> 1.25 mg; 5 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

digoxin	<b>Injection:</b> 250 micrograms/mL in 2 mL ampoule. <b>Oral liquid:</b> 50 micrograms/mL. <b>Tablet:</b> 62.5 micrograms; 250 micrograms.
<input type="checkbox"/> enalapril Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain)	<b>Tablet:</b> 2.5 mg; 5 mg; 10 mg (as hydrogen maleate).
<input type="checkbox"/> furosemide Therapeutic alternatives: - bumetanide - torasemide	<b>Injection:</b> 10 mg/mL in 2 mL, 5 mL ampoule. <b>Oral liquid:</b> 20 mg/5 mL; 50 mg/5 mL [c]. <b>Tablet:</b> 20 mg; 40 mg.
<input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - chlorothiazide - chlorthalidone - indapamide	<b>Oral liquid:</b> 50 mg/5 mL. <b>Solid oral dosage form:</b> 25 mg.
<input type="checkbox"/> losartan Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (C09CA Angiotensin II receptor blockers (ARBs), plain)	<b>Tablet:</b> 25 mg; 50 mg; 100 mg.
spironolactone	<b>Tablet:</b> 25 mg.
<b>Complementary List</b>	
digoxin [c]	<b>Injection:</b> 100 micrograms/mL in 1 mL ampoule; 250 micrograms/mL in 2 mL ampoule. <b>Oral liquid:</b> 50 micrograms/mL. <b>Tablet:</b> 62.5 micrograms; 125 micrograms; 250 micrograms.
dopamine	<b>Injection:</b> 40 mg/mL (hydrochloride) in 5 mL vial.
<b>12.5 Antithrombotic medicines</b>	
<b>12.5.1 Anti-platelet medicines</b>	
acetylsalicylic acid	<b>Tablet:</b> 100 mg.
clopidogrel	<b>Tablet:</b> 75 mg; 300 mg.
<b>12.5.2 Thrombolytic medicines</b>	
<b>Complementary List</b>	
alteplase	<b>Powder for injection:</b> 10 mg; 20 mg; 50 mg in vial.
streptokinase	<b>Powder for injection:</b> 1.5 million IU in vial.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>12.6 Lipid-lowering agents</b>	
<input type="checkbox"/> simvastatin* Therapeutic alternatives: - atorvastatin - fluvastatin - lovastatin - pravastatin	<b>Tablet:</b> 5 mg; 10 mg; 20 mg; 40 mg. *For use in high-risk patients.
<b>12.7 Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease</b>	
acetylsalicylic acid + <input type="checkbox"/> atorvastatin + <input type="checkbox"/> ramipril Therapeutic alternatives: - fluvastatin, lovastatin, pravastatin, simvastatin (for atorvastatin) - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for ramipril)	<b>Tablet:</b> 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg.
acetylsalicylic acid + <input type="checkbox"/> simvastatin + <input type="checkbox"/> ramipril + <input type="checkbox"/> atenolol + <input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - atorvastatin, fluvastatin, lovastatin, pravastatin (for simvastatin) - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for ramipril) - bisoprolol, carvedilol, metoprolol (for atenolol) - chlorthalidone, chlorothiazide, indapamide (for hydrochlorothiazide)	<b>Tablet:</b> 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg.
<input type="checkbox"/> atorvastatin + <input type="checkbox"/> perindopril + <input type="checkbox"/> amlodipine Therapeutic alternatives: - fluvastatin, lovastatin, pravastatin, simvastatin (for atorvastatin) - 4 <sup>th</sup> level ATC chemical subgroup (C09AA ACE inhibitors, plain) (for perindopril) - 4 <sup>th</sup> level ATC chemical subgroup (C08CA Dihydropyridine derivatives) (for amlodipine)	<b>Tablet:</b> 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg.
<b>13. DERMATOLOGICAL MEDICINES</b>	
<b>13.1 Antifungal medicines</b>	
<input type="checkbox"/> miconazole Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (D01AC Imidazole and triazole derivatives) excluding combinations	<b>Cream or ointment:</b> 2% (nitrate).
selenium sulfide	<b>Detergent-based suspension:</b> 2%.
sodium thiosulfate	<b>Solution:</b> 15%.
terbinafine	<b>Cream or ointment:</b> 1% (hydrochloride).
<b>13.2 Anti-infective medicines</b>	
mupirocin	<b>Cream:</b> 2% (as calcium). <b>Ointment:</b> 2%.
potassium permanganate	<b>Aqueous solution:</b> 1:10 000.
silver sulfadiazine <input type="checkbox"/>	<b>Cream:</b> 1%. <input type="checkbox"/> > 2 months.



## WHO Model List of Essential Medicines – 24th List (2025)

13.3 Anti-inflammatory and antipruritic medicines	
<input type="checkbox"/> betamethasone <b>a</b> Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (D07AC Corticosteroids, potent (group III))	<b>Cream or ointment:</b> 0.1% (as valerate). <b>a</b> Hydrocortisone preferred in neonates.
calamine	<b>Lotion.</b>
<input type="checkbox"/> hydrocortisone Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (D07AA Corticosteroids, weak (group I))	<b>Cream or ointment:</b> 1% (acetate).
13.4 Medicines affecting skin differentiation and proliferation	
benzoyl peroxide	<b>Cream or lotion:</b> 5%.
<input type="checkbox"/> calcipotriol Therapeutic alternatives: - calcitriol - tacalcitol	<b>Cream or ointment:</b> 50 micrograms/mL (0.005%). <b>Lotion:</b> 50 micrograms/mL (0.005%).
coal tar	<b>Solution:</b> 5%.
fluorouracil	<b>Ointment:</b> 5%.
<input type="checkbox"/> podophyllum resin Therapeutic alternatives: - podophyllotoxin	<b>Solution:</b> 10% to 25%.
salicylic acid	<b>Solution:</b> 5%.
urea	<b>Cream or ointment:</b> 5%; 10%.
Complementary List	
<input type="checkbox"/> <i>adalimumab</i> * <i>Therapeutic alternatives*:</i> - certolizumab pegol - etanercept - infliximab  <i>*including quality-assured biosimilars</i>	<b>Injection:</b> 10 mg/0.2 mL <b>[c]</b> ; 20 mg/0.2 mL <b>[c]</b> ; 20 mg/0.4 mL <b>[c]</b> ; 40 mg/0.4 mL; 40 mg/0.8 mL; 80 mg/0.8 mL in pre-filled syringe or pre-filled pen.
<i>methotrexate</i>	<b>Tablet:</b> 2.5 mg; 10 mg (as sodium).
<i>ustekinumab</i> * <i>*including quality-assured biosimilars</i>	<b>Injection:</b> 45 mg/0.5 mL in vial, pre-filled syringe, or pre-filled pen; 90 mg/mL in pre-filled syringe or pre-filled pen.
13.5 Scabicides and pediculicides	
<input type="checkbox"/> benzyl benzoate <b>a</b> Therapeutic alternatives: - precipitated sulfur topical ointment	<b>Lotion:</b> 25%. <b>a</b> > 2 years.
permethrin	<b>Cream:</b> 5%. <b>Lotion:</b> 1%.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>13.6 Moisturizers</b>	
urea	<b>Cream:</b> 5%.
glycerol	<b>Cream:</b> 15% to 20%.
<b>13.7 Sunscreens</b>	
sunscreen, broad-spectrum	<b>Topical:</b> Therapeutic broad-spectrum sunscreens should contain proven active ingredients in appropriate amounts to absorb or filter UVA and UVB radiation, and have a high sun protection factor (SPF).
<b>14. DIAGNOSTIC AGENTS</b>	
<b>14.1 Ophthalmic medicines</b>	
fluorescein	<b>Eye drops:</b> 1% (sodium salt).
<input type="checkbox"/> tropicamide Therapeutic alternatives: - atropine - cyclopentolate	<b>Eye drops:</b> 0.5%.
<b>14.2 Radiocontrast media</b>	
<input type="checkbox"/> amidotrizoate Therapeutic alternatives to be reviewed	<b>Injection:</b> 140 mg to 420 mg iodine/mL (as sodium or meglumine salt) in 20 mL ampoule.
barium sulfate	<b>Aqueous suspension.</b>
<input type="checkbox"/> iohexol Therapeutic alternatives to be reviewed	<b>Injection:</b> 140 mg to 350 mg iodine/mL in 5 mL, 10 mL, 20 mL ampoule.
<b><i>Complementary List</i></b>	
<i>barium sulfate [c]</i>	<b><i>Aqueous suspension.</i></b>
<input type="checkbox"/> meglumine iotroxate Therapeutic alternatives to be reviewed	<b><i>Solution:</i></b> 5 g to 8 g iodine in 100 mL to 250 mL.
<b>15. ANTISEPTICS AND DISINFECTANTS</b>	
<b>15.1 Antiseptics</b>	
<input type="checkbox"/> chlorhexidine Therapeutic alternatives to be reviewed	<b>Solution:</b> 5% (digluconate).
<input type="checkbox"/> ethanol Therapeutic alternatives: - propanol	<b>Solution:</b> 70% (denatured).
<input type="checkbox"/> povidone iodine Therapeutic alternatives: - iodine	<b>Solution:</b> 10% (equivalent to 1% available iodine).
<b>15.2 Disinfectants</b>	
alcohol based hand rub	<b>Solution:</b> containing ethanol 80% volume/volume. <b>Solution:</b> containing isopropyl alcohol 75% volume/volume.
chlorine base compound	<b>Liquid:</b> (0.1% available chlorine) for solution. <b>Powder:</b> (0.1% available chlorine) for solution. <b>Solid:</b> (0.1% available chlorine) for solution.

## WHO Model List of Essential Medicines – 24th List (2025)

<input type="checkbox"/> chloroxylenol Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (D08AE Phenol and derivatives)	<b>Solution:</b> 4.8%.
glutaral	<b>Solution:</b> 2%.
hypochlorous acid	<b>Solution (aqueous):</b> containing hypochlorous acid $\geq$ 150 parts per million.
<b>16. DIURETICS</b>	
amiloride	<b>Tablet:</b> 5 mg (hydrochloride).
<input type="checkbox"/> furosemide Therapeutic alternatives: - bumetanide - torasemide	<b>Injection:</b> 10 mg/mL in 2 mL, 5 mL ampoule. <b>Oral liquid:</b> 20 mg/5 mL; 50 mg/5 mL [c]. <b>Tablet:</b> 20 mg; 40 mg.
<input type="checkbox"/> hydrochlorothiazide Therapeutic alternatives: - chlorothiazide - chlortalidone - indapamide	<b>Solid oral dosage form:</b> 25 mg.
mannitol	<b>Injectable solution:</b> 10%; 20%.
spironolactone	<b>Tablet:</b> 25 mg.
<b>Complementary List</b>	
<input type="checkbox"/> hydrochlorothiazide [c] Therapeutic alternatives: - chlorothiazide - chlortalidone	<b>Solid oral dosage form:</b> 12.5 mg [c]; 25 mg.
mannitol [c]	<b>Solution for infusion:</b> 10%; 20%.
spironolactone [c]	<b>Oral liquid:</b> 25 mg/5 mL. <b>Tablet:</b> 12.5 mg; 25 mg.
<b>17. GASTROINTESTINAL MEDICINES</b>	
<b>Complementary List</b>	
pancreatic enzymes [c]	<b>Capsule (modified release)*:</b> 10 000 lipase units + 8000 amylase units + 600 protease units; 25 000 lipase units + 18 000 amylase units + 1000 protease units. *Units expressed in Ph.Eur
<b>17.1 Antiulcer medicines</b>	
<input type="checkbox"/> omeprazole Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (A02BC Proton pump inhibitors) excluding combinations	<b>Powder for injection:</b> 40 mg in vial. <b>Powder for oral liquid:</b> 1 mg/mL; 4 mg/mL. <b>Solid oral dosage form:</b> 10 mg; 20 mg; 40 mg.
<input type="checkbox"/> ranitidine Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (A02BA H <sub>2</sub> -receptor antagonists) excluding combinations	<b>Injection:</b> 25 mg/mL (as hydrochloride) in 2 mL ampoule. <b>Oral liquid:</b> 75 mg/5 mL (as hydrochloride). <b>Tablet:</b> 150 mg (as hydrochloride).

## WHO Model List of Essential Medicines – 24th List (2025)

17.2 Antiemetic medicines	
dexamethasone	<b>Injection:</b> 4 mg/mL dexamethasone phosphate (as sodium phosphate) (equivalent to 3.3 mg/mL dexamethasone base) in 1 mL ampoule. <b>Oral liquid:</b> 0.5 mg/5 mL; 2 mg/5 mL (as sodium phosphate). <b>Tablet:</b> 0.5 mg; 0.75 mg; 1.5 mg; 2 mg; 4 mg (as dexamethasone base).
metoclopramide	<b>Injection:</b> 5 mg/mL (hydrochloride) in 2 mL ampoule. <b>Oral liquid:</b> 5 mg/5 mL [c]. <b>Tablet (scored):</b> 10 mg (hydrochloride).
<input type="checkbox"/> ondansetron Therapeutic alternatives: - dolasetron - granisetron - palonosetron - tropisetron	<b>Injection:</b> 2 mg/mL in 2 mL, 4 mL ampoule (as hydrochloride dihydrate). <b>Oral liquid:</b> 4 mg/5 mL (as hydrochloride dihydrate). <b>Solid oral dosage form:</b> 4 mg; 8 mg; 24 mg (as hydrochloride dihydrate).
Complementary list	
aprepitant	<b>Capsule:</b> 80 mg; 125 mg; 165 mg <b>Powder for oral suspension:</b> 125 mg in sachet
17.3 Anti-inflammatory medicines	
<input type="checkbox"/> sulfasalazine Therapeutic alternatives: - mesalazine	<b>Retention enema:</b> 3 g/100 mL. <b>Suppository:</b> 500 mg. <b>Tablet:</b> 500 mg.
Complementary List	
hydrocortisone	<b>Retention enema:</b> 100 mg/60 mL. <b>Suppository:</b> 25 mg (acetate).
prednisolone	<b>Retention enema:</b> 20 mg/100 mL (as sodium phosphate).
17.4 Laxatives	
<input type="checkbox"/> senna Therapeutic alternatives: - bisacodyl	<b>Tablet:</b> 7.5 mg (sennosides) (or traditional dosage forms).

## WHO Model List of Essential Medicines – 24th List (2025)

<b>17.5 Medicines used in diarrhoea</b>																					
oral rehydration salts – zinc sulfate [c]	Co-package containing: <b>ORS powder for dilution</b> (see Section 17.5.1) – zinc sulfate <b>tablet (dispersible, scored)</b> 20 mg (see Section 17.5.2)																				
<b>17.5.1 Oral rehydration</b>																					
oral rehydration salts	<p><b>Powder for dilution</b> in 200 mL; 500 mL; 1 L.</p> <table> <tr><td>glucose:</td><td>75 mEq or mmol/L</td></tr> <tr><td>sodium:</td><td>75 mEq or mmol/L</td></tr> <tr><td>chloride:</td><td>65 mEq or mmol/L</td></tr> <tr><td>potassium:</td><td>20 mEq or mmol/L</td></tr> <tr><td>citrate:</td><td>10 mEq or mmol/L</td></tr> <tr><td>osmolarity:</td><td>245 mOsm/L</td></tr> <tr><td>glucose:</td><td>13.5 g/L</td></tr> <tr><td>sodium chloride:</td><td>2.6 g/L</td></tr> <tr><td>potassium chloride:</td><td>1.5 g/L</td></tr> <tr><td>trisodium citrate dihydrate*:</td><td>2.9 g/L</td></tr> </table> <p>*trisodium citrate dihydrate may be replaced by sodium hydrogen carbonate (sodium bicarbonate) 2.5 g/L. However, as the stability of this latter formulation is very poor under tropical conditions, it is recommended only when manufactured for immediate use.</p>	glucose:	75 mEq or mmol/L	sodium:	75 mEq or mmol/L	chloride:	65 mEq or mmol/L	potassium:	20 mEq or mmol/L	citrate:	10 mEq or mmol/L	osmolarity:	245 mOsm/L	glucose:	13.5 g/L	sodium chloride:	2.6 g/L	potassium chloride:	1.5 g/L	trisodium citrate dihydrate*:	2.9 g/L
glucose:	75 mEq or mmol/L																				
sodium:	75 mEq or mmol/L																				
chloride:	65 mEq or mmol/L																				
potassium:	20 mEq or mmol/L																				
citrate:	10 mEq or mmol/L																				
osmolarity:	245 mOsm/L																				
glucose:	13.5 g/L																				
sodium chloride:	2.6 g/L																				
potassium chloride:	1.5 g/L																				
trisodium citrate dihydrate*:	2.9 g/L																				
<b>17.5.2 Medicines for diarrhoea</b>																					
zinc sulfate*	<p><b>Tablet (dispersible, scored):</b> 20 mg.</p> <p>*In acute diarrhoea zinc sulfate should be used as an adjunct to oral rehydration salts.</p>																				
<b>18. MEDICINES FOR ENDOCRINE DISORDERS</b>																					
<b>18.1 Adrenal hormones and synthetic substitutes</b>																					
fludrocortisone	<p><b>Oral liquid:</b> 100 micrograms/mL (acetate) [c].</p> <p><b>Tablet:</b> 100 micrograms (acetate).</p>																				
hydrocortisone	<p><b>Granules:</b> 0.5 mg; 1 mg; 2 mg; 5 mg in capsule [c].</p> <p><b>Tablet:</b> 5 mg; 10 mg; 20 mg.</p>																				
<input type="checkbox"/> prednisolone Therapeutic alternatives: - prednisone	<b>Tablet:</b> 1 mg.																				
<b>18.2 Androgens</b>																					
<i>Complementary List</i>																					
testosterone	<b>Injection:</b> 200 mg (enanthate) in 1 mL ampoule.																				
<b>18.3 Estrogens</b>																					
<b>18.4 Progestogens</b>																					
<input type="checkbox"/> medroxyprogesterone acetate Therapeutic alternatives: - norethisterone	<b>Tablet:</b> 5 mg.																				

## WHO Model List of Essential Medicines – 24th List (2025)

<b>18.5 Medicines for diabetes</b>	
<b>18.5.1 Insulins</b>	
<input type="checkbox"/> insulin (analogue, long-acting)* Therapeutic alternatives: - insulin glargine - insulin degludec - insulin detemir  <i>*including quality-assured biosimilars</i>	<b>Injection solution:</b> 100 IU/mL in 10 mL vial; 100 IU/mL in 3 mL cartridge or pre-filled pen.
<input type="checkbox"/> insulin (analogue, rapid-acting)* Therapeutic alternatives: - insulin lispro - insulin aspart - insulin glulisine  <i>*including quality-assured biosimilars</i>	<b>Injection solution:</b> 100 IU/mL in 10 mL vial; 100 IU/mL in 3 mL cartridge or pre-filled pen.
insulin (human, intermediate-acting)* <i>*including quality-assured biosimilars</i>	<b>Injection suspension:</b> 40 IU/mL in 10 mL vial; 100 IU/mL in 10 mL vial; 100 IU/mL in 3 mL cartridge or pre-filled pen (as compound insulin zinc suspension or isophane insulin).
insulin (human, short-acting)* <i>*including quality-assured biosimilars</i>	<b>Injection solution:</b> 40 IU/mL in 10 mL vial; 100 IU/mL in 10 mL vial; 100 IU/mL in 3 mL cartridge or pre-filled pen.
<b>18.5.2 Hypoglycaemic agents</b>	
<input type="checkbox"/> empagliflozin Therapeutic alternatives: - canagliflozin - dapagliflozin	<b>Tablet:</b> 10 mg; 25 mg.
<input type="checkbox"/> gliclazide* Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (A10BB Sulfonylureas)	<b>Solid oral dosage form:</b> (controlled-release tablets) 30 mg; 60 mg; 80 mg.  <i>*glibenclamide not suitable above 60 years.</i>
metformin	<b>Tablet:</b> 500 mg (hydrochloride).
<input type="checkbox"/> semaglutide* Therapeutic alternatives: - dulaglutide - liraglutide - tirzepatide  <i>*including quality-assured biosimilars</i>	<b>Injection solution:</b> 0.68 mg/mL; 1.34 mg/mL; 2.68 mg/mL.
<b>Complementary List</b>	
metformin [c]	<b>Tablet:</b> 500 mg (hydrochloride).
<b>18.6 Medicines for hypoglycaemia</b>	
glucagon	<b>Injection:</b> 1 mg/mL as powder and diluent.
<b>Complementary List</b>	
diazoxide [c]	<b>Oral liquid:</b> 50 mg/mL. <b>Tablet:</b> 50 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>18.7 Thyroid hormones and antithyroid medicines</b>	
levothyroxine	<b>Tablet:</b> 25 micrograms [c]; 50 micrograms; 100 micrograms (sodium salt).
potassium iodide	<b>Tablet (scored):</b> 65 mg.
<input type="checkbox"/> methimazole Therapeutic alternatives: - carbimazole (depending on local availability)	<b>Tablet:</b> 5mg, 10mg.
propylthiouracil*	<b>Tablet:</b> 50 mg. *For use when alternative first-line treatment is not appropriate or available; and in patients during the first trimester of pregnancy.
<b>Complementary List</b>	
iodine + potassium iodide (Lugol's solution) [c]	<b>Oral liquid (aqueous):</b> 5% w/v + 10% w/v.
<input type="checkbox"/> methimazole [c] Therapeutic alternatives: - carbimazole (depending on local availability)	<b>Tablet:</b> 5 mg; 10 mg.
potassium iodide [c]	<b>Tablet (scored):</b> 65 mg.
propylthiouracil* [c]	<b>Tablet:</b> 50 mg. *For use when alternative first-line treatment is not appropriate or available
<b>18.8 Medicines for disorders of the pituitary hormone system</b>	
<input type="checkbox"/> cabergoline Therapeutic alternatives: - bromocriptine	<b>Tablet:</b> 0.5 mg; 1 mg.
<b>Complementary List</b>	
octreotide	<b>Injection (immediate-release):</b> 0.05 mg/mL; 0.1 mg/mL; 0.5 mg/mL (as acetate) in 1 mL vial. <b>Injection (modified-release):</b> 20 mg (as acetate) in vial plus diluent.
<b>19. IMMUNOLOGICALS</b>	
<b>19.1 Diagnostic agents</b>	
All tuberculin should comply with the WHO requirements for tuberculin.	
tuberculin, purified protein derivative (PPD)	<b>Injection.</b>
<b>19.2 Sera, immunoglobulins and monoclonal antibodies</b>	
All plasma fractions should comply with the WHO requirements.	
anti-rabies virus monoclonal antibodies* *including quality-assured biosimilars	<b>Injection:</b> 40 IU/mL in 1.25 mL, 2.5 mL vial; 100 IU/mL in 2.5 mL vial (human). <b>Injection:</b> 300 IU/mL in 10 mL vial; 600 IU/mL in 1 mL, 2.5 mL and 5 mL vial (murine).
antivenom immunoglobulin*	<b>Injection.</b> *Exact type to be defined locally.
diphtheria antitoxin	<b>Injection:</b> 10 000 IU; 20 000 IU in vial.
equine rabies immunoglobulin	<b>Injection:</b> 150 IU/mL; 200 IU/mL; 300 IU/mL; 400 IU/mL in vial.

## WHO Model List of Essential Medicines – 24th List (2025)

### 19.3 Vaccines

WHO immunization policy recommendations are published in vaccine position papers based on recommendations made by the Strategic Advisory Group of Experts (SAGE) on Immunization.

WHO vaccine position papers are periodically revised to assess the need for an update. The list below details the vaccines for which there is a recommendation from WHO and a corresponding WHO vaccine position paper as at May 2025. The most recent versions of the WHO position papers, reflecting the current evidence related to a specific vaccine and the related recommendations, can be accessed at any time on the WHO website at: <https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/position-papers>

Vaccine recommendations may be universal or conditional (e.g., in certain regions, in some high-risk populations or as part of immunization programmes with certain characteristics). Details are available in the relevant position papers, and in the Summary Tables of WHO Routine Immunization Recommendations available on the WHO website at: <https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/who-recommendations-for-routine-immunization---summary-tables>

Vaccines included on the Model Lists reflect the recommendations of SAGE, as per the available policy reflected in the WHO vaccine position papers.

Countries are encouraged to consider inclusion of specific vaccines into their national immunization schedule based on national priorities by carefully assessing various criteria such as local burden of disease and disease epidemiology, acceptability, cost, cost-effectiveness, programmatic feasibility, regulatory status, and availability of products.

All vaccines should comply with the WHO requirements for biological substances.

BCG vaccine	
cholera vaccine	
dengue vaccine	
diphtheria vaccine	
Ebola vaccine	
Haemophilus influenzae type b vaccine	
hepatitis A vaccine	
hepatitis B vaccine	
hepatitis E vaccine	
human papilloma virus (HPV) vaccine	
influenza vaccine (seasonal)	
Japanese encephalitis vaccine	
malaria vaccine [c]	
measles vaccine	
meningococcal meningitis vaccine	
mpox vaccine	
mumps vaccine	
pertussis vaccine	
pneumococcal vaccine	
poliomyelitis vaccine	
rabies vaccine	
respiratory syncytial virus vaccine	
rotavirus vaccine	
rubella vaccine	



## WHO Model List of Essential Medicines – 24th List (2025)

tetanus vaccine	
tick-borne encephalitis vaccine	
typhoid vaccine	
varicella vaccine	
yellow fever vaccine	
<b>20. MUSCLE RELAXANTS (PERIPHERALLY-ACTING) AND CHOLINESTERASE INHIBITORS</b>	
<input type="checkbox"/> atracurium Therapeutic alternatives to be reviewed	<b>Injection:</b> 10 mg/mL (besylate).
neostigmine	<b>Injection:</b> 500 micrograms/mL (methysulfate) in 1 mL ampoule; 2.5 mg/mL (methysulfate) in 1 mL ampoule. <b>Tablet:</b> 15 mg (bromide).
suxamethonium	<b>Injection:</b> 50 mg/mL (chloride) in 2 mL ampoule.
<input type="checkbox"/> vecuronium [c] Therapeutic alternatives: -atracurium	<b>Powder for injection:</b> 10 mg (bromide) in vial.
<i>Complementary List</i>	
<i>pyridostigmine</i>	<b>Injection:</b> 5 mg/mL (bromide) in ampoule or vial. <b>Tablet (scored):</b> 60 mg (bromide).
<input type="checkbox"/> vecuronium Therapeutic alternatives to be reviewed	<b>Powder for injection:</b> 10 mg (bromide) in vial.
<b>21. OPHTHALMOLOGICAL PREPARATIONS</b>	
<b>21.1 Anti-infective agents</b>	
aciclovir	<b>Ointment:</b> 3% w/w.
azithromycin	<b>Solution (eye drops):</b> 1.5%. – Trachoma
erythromycin	<b>Ointment:</b> 0.5% [c] – Infections due to <i>Chlamydia trachomatis</i> or <i>Neisseria gonorrhoea</i> .
<input type="checkbox"/> gentamicin Therapeutic alternatives: - amikacin - kanamycin - netilmicin - tobramycin	<b>Solution (eye drops):</b> 0.3% (sulfate). – Bacterial blepharitis – Bacterial conjunctivitis
natamycin	<b>Suspension (eye drops):</b> 5% – Fungal keratitis
<input type="checkbox"/> ofloxacin Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (S01AE Fluoroquinolones)	<b>Solution (eye drops):</b> 0.3%. – Bacterial conjunctivitis – Bacterial keratitis

## WHO Model List of Essential Medicines – 24th List (2025)

<input type="checkbox"/> tetracycline Therapeutic alternatives: - chlortetracycline - oxytetracycline	<b>Eye ointment:</b> 1% (hydrochloride). - <i>Bacterial blepharitis</i> - <i>Bacterial conjunctivitis</i> - <i>Bacterial keratitis</i> - <i>Trachoma</i>
<b>21.2 Anti-inflammatory agents</b>	
<input type="checkbox"/> prednisolone Therapeutic alternatives to be reviewed	<b>Solution (eye drops):</b> 0.5% (sodium phosphate).
<b>21.3 Local anaesthetics</b>	
<input type="checkbox"/> tetracaine <sup>a</sup> Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (S01HA Local anaesthetics) excluding cocaine and combinations	<b>Solution (eye drops):</b> 0.5% (hydrochloride). <sup>a</sup> Not in preterm neonates.
<b>21.4 Miotics and antiglaucoma medicines</b>	
acetazolamide	<b>Tablet:</b> 250 mg.
latanoprost	<b>Solution (eye drops):</b> 50 micrograms/mL.
<input type="checkbox"/> pilocarpine Therapeutic alternatives: - carbachol	<b>Solution (eye drops):</b> 2%; 4% (hydrochloride or nitrate).
<input type="checkbox"/> timolol Therapeutic alternatives: - 4 <sup>th</sup> level ATC chemical subgroup (S01ED Beta blocking agents) excluding combinations	<b>Solution (eye drops):</b> 0.25%; 0.5% (as hydrogen maleate).
<b>21.5 Mydriatics</b>	
<input type="checkbox"/> atropine <sup>a</sup> Therapeutic alternatives*: - cyclopentolate hydrochloride - homatropine hydrobromide <i>*EMLc only</i>	<b>Solution (eye drops):</b> 0.1%; 0.5%; 1% (sulfate). <sup>a</sup> > 3 months.
<b>Complementary List</b>	
<i>epinephrine (adrenaline)</i>	<b>Solution (eye drops):</b> 2% (as hydrochloride).
<b>21.6 Anti-vascular endothelial growth factor (VEGF) preparations</b>	
<b>Complementary List</b>	
<i>bevacizumab*</i> <i>*including quality-assured biosimilars</i>	<b>Injection:</b> 25 mg/mL.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>22. MEDICINES FOR REPRODUCTIVE HEALTH AND PERINATAL CARE</b>	
<b>22.1 Contraceptives</b>	
<b>22.1.1 Oral hormonal contraceptives</b>	
□ ethinylestradiol + □ levonorgestrel Therapeutic alternatives to be reviewed	<b>Tablet:</b> 30 micrograms + 150 micrograms.
□ ethinylestradiol + □ norethisterone Therapeutic alternatives to be reviewed	<b>Tablet:</b> 35 micrograms + 1 mg.
levonorgestrel	<b>Tablet:</b> 30 micrograms; 750 micrograms (pack of two); 1.5 mg.
ulipristal	<b>Tablet:</b> 30 mg (as acetate).
<b>22.1.2 Injectable hormonal contraceptives</b>	
estradiol cypionate + medroxyprogesterone acetate	<b>Injection:</b> 5 mg + 25 mg.
medroxyprogesterone acetate	<b>Injection (intramuscular):</b> 150 mg/mL in 1 mL vial. <b>Injection (subcutaneous):</b> 104 mg/0.65 mL in pre-filled syringe or single-dose injection delivery system.
norethisterone enantate	<b>Oily solution:</b> 200 mg/mL in 1 mL ampoule.
<b>22.1.3 Intrauterine devices</b>	
copper-containing device	
levonorgestrel-releasing intrauterine system	<b>Intrauterine system:</b> with reservoir containing 52 mg of levonorelrel
<b>22.1.4 Barrier methods</b>	
condoms	
diaphragms	
<b>22.1.5 Implantable contraceptives</b>	
etonogestrel-releasing implant	<b>Single-rod etonogestrel-releasing implant:</b> containing 68 mg of etonogestrel.
levonorgestrel-releasing implant	<b>Two-rod levonorgestrel-releasing implant:</b> each rod containing 75 mg of levonorgestrel (150 mg total).
<b>22.1.6 Intravaginal contraceptives</b>	
ethinylestradiol + etonogestrel	<b>Vaginal ring:</b> containing 2.7 mg + 11.7 mg.
progesterone vaginal ring*	<b>Progesterone-releasing vaginal ring:</b> containing 2.074 g of micronized progesterone.  *For use in women actively breastfeeding at least 4 times per day
<b>22.2 Ovulation inducers</b>	
<b>Complementary List</b>	
clomifene	<b>Tablet:</b> 50 mg (citrate).

## WHO Model List of Essential Medicines – 24th List (2025)

<input type="checkbox"/> <i>letrozole</i> <i>Therapeutic alternatives:</i> - <i>anastrozole</i>	<b>Solid oral dosage form:</b> 2.5 mg.
<b>22.3 Uterotonics</b>	
carbetocin	<b>Injection (heat stable):</b> 100 micrograms/mL.
<input type="checkbox"/> ergometrine <i>Therapeutic alternatives:</i> - methylexergometrine	<b>Injection:</b> 200 micrograms (hydrogen maleate) in 1 mL ampoule.
misoprostol	<b>Tablet:</b> 200 micrograms. – Prevention and treatment of postpartum haemorrhage where oxytocin is not available or cannot be safely used. <b>Vaginal tablet:</b> 25 micrograms.* *Only for use for induction of labour where appropriate facilities are available.
oxytocin	<b>Injection:</b> 10 IU in 1 mL.
<b>22.4 Medicines for medical abortion</b>	
mifepristone – misoprostol	<b>Tablet</b> 200 mg – <b>tablet</b> 200 micrograms. <b>Co-package containing:</b> mifepristone 200 mg tablet [1] and misoprostol 200 micrograms tablet [4] – Management of intrauterine fetal demise; – Management of induced abortion.
misoprostol	<b>Tablet:</b> 200 micrograms. – Management of incomplete abortion and miscarriage.
<b>22.5 Antioxytocics (tocolytics)</b>	
nifedipine	<b>Immediate-release capsule:</b> 10 mg.
<b>22.6 Other medicines administered to the mother</b>	
dexamethasone	<b>Injection:</b> 4 mg/mL dexamethasone phosphate (as sodium phosphate) (equivalent to 3.3 mg/mL dexamethasone base) in 1 mL ampoule.

## WHO Model List of Essential Medicines – 24th List (2025)

multiple micronutrient supplement*	<p><b>Tablet containing:</b></p> <table> <tr> <td>Vitamin A (retinol acetate)</td><td>800 micrograms retinol activity equivalent</td></tr> <tr> <td>Vitamin C (ascorbic acid)</td><td>70 mg</td></tr> <tr> <td>Vitamin D (cholecalciferol)</td><td>5 micrograms (200 IU)</td></tr> <tr> <td>Vitamin E (alpha tocopherol succinate)</td><td>10 mg alpha tocopherol equivalent</td></tr> <tr> <td>Vitamin B1 (thiamine mononitrate)</td><td>1.4 mg</td></tr> <tr> <td>Vitamin B2 (riboflavin)</td><td>1.4 mg</td></tr> <tr> <td>Vitamin B3 (niacinamide)</td><td>18 mg niacin equivalent</td></tr> <tr> <td>Vitamin B6 (pyridoxine hydrochloride)</td><td>1.9 mg</td></tr> <tr> <td>Folic acid (folic acid)</td><td>680 micrograms dietary folate equivalent (400 micrograms)</td></tr> <tr> <td>Vitamin B12 (cyanocobalamin)</td><td>2.6 micrograms</td></tr> <tr> <td>Iron (ferrous fumarate)</td><td>30 mg</td></tr> <tr> <td>Iodine (potassium iodide)</td><td>150 micrograms</td></tr> <tr> <td>Zinc (zinc oxide)</td><td>15 mg</td></tr> <tr> <td>Selenium (sodium selenite)</td><td>65 micrograms</td></tr> <tr> <td>Copper (cupric oxide)</td><td>2 mg</td></tr> </table> <p>*For use in specific contexts. Refer to current WHO recommendations.</p>	Vitamin A (retinol acetate)	800 micrograms retinol activity equivalent	Vitamin C (ascorbic acid)	70 mg	Vitamin D (cholecalciferol)	5 micrograms (200 IU)	Vitamin E (alpha tocopherol succinate)	10 mg alpha tocopherol equivalent	Vitamin B1 (thiamine mononitrate)	1.4 mg	Vitamin B2 (riboflavin)	1.4 mg	Vitamin B3 (niacinamide)	18 mg niacin equivalent	Vitamin B6 (pyridoxine hydrochloride)	1.9 mg	Folic acid (folic acid)	680 micrograms dietary folate equivalent (400 micrograms)	Vitamin B12 (cyanocobalamin)	2.6 micrograms	Iron (ferrous fumarate)	30 mg	Iodine (potassium iodide)	150 micrograms	Zinc (zinc oxide)	15 mg	Selenium (sodium selenite)	65 micrograms	Copper (cupric oxide)	2 mg
Vitamin A (retinol acetate)	800 micrograms retinol activity equivalent																														
Vitamin C (ascorbic acid)	70 mg																														
Vitamin D (cholecalciferol)	5 micrograms (200 IU)																														
Vitamin E (alpha tocopherol succinate)	10 mg alpha tocopherol equivalent																														
Vitamin B1 (thiamine mononitrate)	1.4 mg																														
Vitamin B2 (riboflavin)	1.4 mg																														
Vitamin B3 (niacinamide)	18 mg niacin equivalent																														
Vitamin B6 (pyridoxine hydrochloride)	1.9 mg																														
Folic acid (folic acid)	680 micrograms dietary folate equivalent (400 micrograms)																														
Vitamin B12 (cyanocobalamin)	2.6 micrograms																														
Iron (ferrous fumarate)	30 mg																														
Iodine (potassium iodide)	150 micrograms																														
Zinc (zinc oxide)	15 mg																														
Selenium (sodium selenite)	65 micrograms																														
Copper (cupric oxide)	2 mg																														
tranexamic acid	<b>Injection:</b> 100 mg/mL in 10 mL ampoule.																														
<b>22.7 Medicines administered to the neonate [c]</b>																															
caffeine citrate [c]	<p><b>Injection:</b> 20 mg/mL (equivalent to 10 mg caffeine base/mL).</p> <p><b>Oral liquid:</b> 20 mg/mL (equivalent to 10 mg caffeine base/mL).</p>																														
chlorhexidine [c]	<b>Solution or gel:</b> 7.1% (digluconate) delivering 4% chlorhexidine (for umbilical cord care).																														
<b>Complementary List</b>																															
<p><input type="checkbox"/> <i>alprostadil (prostaglandin E1) [c]</i></p> <p>Therapeutic alternatives:</p> <p>- <i>dinoprostone (prostaglandin E2)</i></p>	<b>Solution for injection:</b> 0.5 mg/mL in alcohol.																														
<i>beractant [c]</i>	<b>Suspension for intratracheal instillation:</b> 25 mg/mL.																														
<p><input type="checkbox"/> <i>ibuprofen [c]</i></p> <p>Therapeutic alternatives:</p> <p>- <i>indometacin</i></p>	<b>Solution for injection:</b> 5 mg/mL.																														
<i>poractant alfa [c]</i>	<b>Suspension for intratracheal instillation:</b> 80 mg/mL.																														
<b>23. PERITONEAL DIALYSIS SOLUTION</b>																															
<b>Complementary List</b>																															
<i>intraperitoneal dialysis solution</i>	<b>Solution:</b> of appropriate composition in accordance with local clinical guidelines.																														

## WHO Model List of Essential Medicines – 24th List (2025)

<b>24. MEDICINES FOR MENTAL AND BEHAVIOURAL DISORDERS</b>	
<b>24.1 Medicines for psychotic disorders</b>	
<input type="checkbox"/> fluphenazine Therapeutic alternatives: - haloperidol decanoate - zuclopenthixol decanoate	<b>Injection:</b> 25 mg (decanoate or enantate) in 1 mL ampoule.
<input type="checkbox"/> haloperidol Therapeutic alternatives: - chlorpromazine	<b>Tablet:</b> 2 mg; 5 mg.
haloperidol	<b>Injection:</b> 5 mg/mL in 1 mL ampoule.
olanzapine	<b>Powder for injection:</b> 10 mg in vial.
<input type="checkbox"/> paliperidone Therapeutic alternatives: - aripiprazole once-monthly injection - risperidone injection	<b>Injection (prolonged-release):</b> 25 mg; 50 mg; 75 mg; 100 mg; 150 mg (as palmitate) in pre-filled syringe.
<input type="checkbox"/> risperidone Therapeutic alternatives: - aripiprazole - olanzapine - paliperidone - quetiapine	<b>Solid oral dosage form:</b> 0.25 mg to 6.0 mg.
<b><i>Complementary List</i></b>	
<i>clozapine</i>	<b><i>Solid oral dosage form:</i></b> 25 mg to 200 mg.
<b>24.2 Medicines for mood disorders</b>	
<b><i>24.2.1 Medicines for depressive disorders</i></b>	
amitriptyline	<b>Tablet:</b> 25 mg; 75 mg (hydrochloride).
<input type="checkbox"/> fluoxetine Therapeutic alternatives: - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	<b>Solid oral dosage form:</b> 20 mg (as hydrochloride).
<b><i>24.2.2 Medicines for bipolar disorders</i></b>	
carbamazepine	<b>Tablet (scored):</b> 100 mg; 200 mg; 400 mg.
lithium carbonate	<b>Solid oral dosage form:</b> 300 mg.
<input type="checkbox"/> quetiapine Therapeutic alternatives: - aripiprazole - olanzapine - paliperidone	<b>Tablet (immediate-release):</b> 25 mg; 100 mg; 150 mg; 200 mg; 300 mg.  <b>Tablet (modified-release):</b> 50 mg; 150 mg; 200 mg; 300 mg; 400 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

valproic acid (sodium valproate)*  <i>*Valproic acid (sodium valproate) is not recommended in women and girls of childbearing potential owing to the high risk of birth defects and neurodevelopmental disorders in children exposed to valproic acid (sodium valproate) in the womb.</i>	Tablet (enteric-coated): 200 mg; 500 mg.
<b>24.3 Medicines for anxiety disorders</b>	
<input type="checkbox"/> diazepam*  Therapeutic alternatives: - lorazepam	Tablet (scored): 2 mg; 5 mg.  *For short-term emergency management of acute and severe anxiety symptoms only
<input type="checkbox"/> fluoxetine  Therapeutic alternatives:  - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	Solid oral dosage form: 20 mg (as hydrochloride).
<b>24.4 Medicines for obsessive compulsive disorders</b>	
clomipramine	Capsule: 10 mg; 25 mg (hydrochloride).
<input type="checkbox"/> fluoxetine  Therapeutic alternatives:  - citalopram - escitalopram - fluvoxamine - paroxetine - sertraline	Solid oral dosage form: 20 mg (as hydrochloride).
<b>24.5 Medicines for disorders due to psychoactive substance use</b>	
<b>24.5.1 Medicines for alcohol use disorders</b>	
acamprosate calcium	Tablet: 333 mg.
naltrexone	Injection suspension (extended-release): 380 mg in vial. Tablet: 50 mg.
<b>24.5.2 Medicines for nicotine use disorders</b>	
bupropion	Tablet (sustained-release): 150 mg (hydrochloride).
cytisine (cytisinicline)	Tablet: 1.5 mg.
nicotine replacement therapy (NRT)	Chewing gum: 2 mg; 4 mg (as polacrilex). Lozenge: 2 mg; 4 mg. Oral spray: 1 mg per actuation. Transdermal patch: 5 mg to 30 mg/16 hrs; 7 mg to 21 mg/24 hrs.
varenicline	Tablet: 0.5 mg, 1 mg.

## WHO Model List of Essential Medicines – 24th List (2025)

<b>24.5.3 Medicines for opioid use disorders</b>	
<b>Complementary List</b>	
<input type="checkbox"/> <b>methadone*</b> Therapeutic alternatives: - buprenorphine	<b>Concentrate for oral liquid:</b> 5 mg/mL; 10 mg/mL (hydrochloride). <b>Oral liquid:</b> 5 mg/5 mL; 10 mg/5 mL (hydrochloride). <i>*The medicines should only be used within an established support programme.</i>
<b>25. MEDICINES ACTING ON THE RESPIRATORY TRACT</b>	
<b>25.1 Antiasthmatic medicines and medicines for chronic obstructive pulmonary disease</b>	
<input type="checkbox"/> <b>budesonide</b> Therapeutic alternatives: - beclometasone - ciclesonide - fluticasone - mometasone	<b>Powder for inhalation:</b> 100 micrograms per actuation; 200 micrograms per actuation in dry powder inhaler. <b>Suspension for inhalation:</b> 100 micrograms per actuation; 200 micrograms per actuation in pressurized metered-dose inhaler.
<input type="checkbox"/> <b>budesonide + <input type="checkbox"/> formoterol</b> Therapeutic alternatives: - beclometasone + formoterol - budesonide + salmeterol - fluticasone + formoterol - fluticasone furoate + vilanterol - mometasone + formoterol	<b>Powder for inhalation:</b> 100 micrograms + 6 micrograms per actuation; 200 micrograms + 6 micrograms per actuation in dry powder inhaler.
epinephrine (adrenaline)	<b>Injection:</b> 1 mg/mL (as hydrochloride or hydrogen tartrate) in 1 mL ampoule.
ipratropium bromide	<b>Solution for inhalation:</b> 20 micrograms per actuation in pressurized metered-dose inhaler.
<input type="checkbox"/> <b>salbutamol</b> Therapeutic alternatives: - terbutaline	<b>Injection:</b> 500 micrograms/mL (as sulfate) in 1 mL, 5 mL ampoule. <b>Solution for inhalation:</b> 100 micrograms (as sulfate) per actuation in pressurized metered-dose inhaler; 2.5 mg/2.5 mL; 5 mg/2.5 mL (as sulfate) in 2.5 mL single-dose ampoules for use in nebulizers; 5 mg/mL (as sulfate) in multi-dose bottle for use in nebulizers.
<input type="checkbox"/> <b>tiotropium</b> Therapeutic alternatives: - aclidinium - glycopyrronium - umeclidinium	<b>Powder for inhalation:</b> 18 micrograms in capsule. <b>Solution for inhalation:</b> 1.25 micrograms; 2.5 micrograms per actuation in soft mist inhaler.
<b>26. SOLUTIONS CORRECTING WATER, ELECTROLYTE AND ACID–BASE DISTURBANCES</b>	
<b>26.1 Oral</b>	
oral rehydration salts	See section 17.5.1.
potassium chloride	<b>Powder for solution.</b>



## WHO Model List of Essential Medicines – 24th List (2025)

<b>26.2 Parenteral</b>	
glucose	<b>Injectable solution:</b> 5% (isotonic); 10% (hypertonic); 50% (hypertonic).
glucose with sodium chloride	<b>Injectable solution:</b> 4% glucose, 0.18% sodium chloride (equivalent to Na <sup>+</sup> 30 mmol/L, Cl <sup>-</sup> 30 mmol/L). <b>Injectable solution:</b> 5% glucose, 0.9% sodium chloride (equivalent to Na <sup>+</sup> 150 mmol/L and Cl <sup>-</sup> 150 mmol/L); 5% glucose, 0.45% sodium chloride (equivalent to Na <sup>+</sup> 75 mmol/L and Cl <sup>-</sup> 75 mmol/L) [c].
potassium chloride	<b>Solution:</b> 11.2% in 20 mL ampoule (equivalent to K <sup>+</sup> 1.5 mmol/mL, Cl <sup>-</sup> 1.5 mmol/mL). <b>Solution for dilution:</b> 7.5% (equivalent to K <sup>+</sup> 1 mmol/mL and Cl <sup>-</sup> 1 mmol/mL) [c]; 15% (equivalent to K <sup>+</sup> 2 mmol/mL and Cl <sup>-</sup> 2 mmol/mL) [c].
sodium chloride	<b>Injectable solution:</b> 0.9% isotonic (equivalent to Na <sup>+</sup> 154 mmol/L, Cl <sup>-</sup> 154 mmol/L).
sodium hydrogen carbonate	<b>Injectable solution:</b> 1.4% isotonic (equivalent to Na <sup>+</sup> 167 mmol/L, HCO <sub>3</sub> <sup>-</sup> 167 mmol/L). <b>Solution:</b> 8.4% in 10 mL ampoule (equivalent to Na <sup>+</sup> 1000 mmol/L, HCO <sub>3</sub> <sup>-</sup> 1000 mmol/L).
sodium lactate, compound solution	<b>Injectable solution.</b>
<b>26.3 Miscellaneous</b>	
water for injection	2 mL; 5 mL; 10 mL ampoules.
<b>27. VITAMINS AND MINERALS</b>	
ascorbic acid	<b>Tablet:</b> 50 mg.
calcium	<b>Tablet:</b> 500 mg (elemental).
<input type="checkbox"/> colecalciferol [c] Therapeutic alternatives: - ergocalciferol	<b>Oral liquid:</b> 400 IU/mL. <b>Solid oral dosage form:</b> 400 IU; 1000 IU.
<input type="checkbox"/> ergocalciferol Therapeutic alternatives: - colecalciferol	<b>Oral liquid:</b> 250 micrograms/mL (10 000 IU/mL). <b>Solid oral dosage form:</b> 1.25 mg (50 000 IU).
iodine	<b>Iodized oil:</b> 480 mg iodine/mL in 10 mL ampoule or vial (oral or injectable).
multiple micronutrient powder [c]	<b>Sachets containing:</b> - iron (elemental) 12.5 mg (as coated ferrous fumarate) - zinc (elemental) 5 mg - vitamin A 300 micrograms - with or without other micronutrients at recommended daily values
nicotinamide	<b>Tablet:</b> 50 mg.
pyridoxine	<b>Tablet:</b> 10 mg [c]; 25 mg (hydrochloride).

## WHO Model List of Essential Medicines – 24th List (2025)

retinol	<b>Soft capsule:</b> 50 000 IU; 100 000 IU; 200 000 IU (as acetate or palmitate). <b>Oral liquid:</b> 100 000 IU/mL (as palmitate). <b>Water-miscible injection:</b> 50 000 IU/mL (as palmitate) in 2 mL ampoule or vial.
riboflavin	<b>Tablet:</b> 5 mg.
thiamine	<b>Injection:</b> 50 mg/mL (hydrochloride) in ampoule or vial <b>[c]</b> . <b>Tablet:</b> 50 mg (hydrochloride).
<b>Complementary List</b>	
calcium gluconate	<b>Injection:</b> 100 mg/mL in 10 mL ampoule.
<b>28. EAR, NOSE AND THROAT MEDICINES</b>	
acetic acid <b>[c]</b>	<b>Solution (ear drops):</b> 2%.
<input type="checkbox"/> budesonide <b>[c]</b> Therapeutic alternatives to be reviewed	<b>Nasal spray:</b> 32 micrograms; 64 micrograms per actuation.
<input type="checkbox"/> ciprofloxacin <b>[c]</b> Therapeutic alternatives: - ofloxacin	<b>Solution (ear drops):</b> 0.3% (as hydrochloride).
<input type="checkbox"/> xylometazoline <b>[c]</b> Therapeutic alternatives to be reviewed	<b>Nasal drops:</b> 0.05%. <b>Nasal spray:</b> 0.05%.
<b>29. MEDICINES FOR DISEASES OF JOINTS</b>	
<b>29.1 Medicines used to treat gout</b>	
allopurinol	<b>Tablet:</b> 100 mg.
<b>29.2 Disease-modifying anti-rheumatic drugs (DMARDs)</b>	
chloroquine	<b>Tablet:</b> 100 mg; 150 mg (as phosphate or sulfate).
<b>Complementary List</b>	
azathioprine	<b>Tablet:</b> 50 mg.
hydroxychloroquine	<b>Solid oral dosage form:</b> 200 mg (as sulfate).
methotrexate	<b>Tablet:</b> 2.5 mg (as sodium).
penicillamine	<b>Solid oral dosage form:</b> 250 mg.
sulfasalazine	<b>Tablet:</b> 500 mg.
<b>29.3 Medicines for juvenile joint diseases</b>	
<b>Complementary List</b>	
acetylsalicylic acid*	<b>Suppository:</b> 150 mg; 300 mg. <b>Tablet:</b> 75 mg to 500 mg. <b>Tablet (dispersible):</b> 75 mg; 300 mg; 500 mg. <i>*For use for rheumatic fever, juvenile arthritis, Kawasaki disease.</i>

## WHO Model List of Essential Medicines – 24th List (2025)

<input type="checkbox"/> <i>adalimumab</i> * <i>Therapeutic alternatives</i> *: - certolizumab pegol - etanercept - golimumab - infliximab  <i>*including quality-assured biosimilars</i>	<b>Injection:</b> 10 mg/0.2 mL [ <b>c</b> ]; 20 mg/0.2 mL [ <b>c</b> ]; 20 mg/0.4 mL [ <b>c</b> ]; 40 mg/0.4 mL; 40 mg/0.8 mL; 80 mg/0.8 mL in pre-filled syringe or pre-filled pen.
<i>methotrexate</i>	<b>Tablet:</b> 2.5 mg (as sodium).
<input type="checkbox"/> <i>triamcinolone hexacetonide</i> <i>Therapeutic alternatives</i> : - triamcinolone acetonide	<b>Injection:</b> 20 mg/mL in vial.
<b>30. DENTAL MEDICINES AND PREPARATIONS</b>	
fluoride	<b>Gel:</b> containing 2500 to 12 500 ppm fluoride (any type). <b>Mouthrinse:</b> containing 230 to 900 ppm fluoride (any type). <b>Toothpaste, cream or gel:</b> containing 1000 to 1500 ppm fluoride (any type). <b>Varnish:</b> containing 22 500 ppm fluoride (any type).
glass ionomer cement	<b>Single-use capsules:</b> 0.4 g powder + 0.09 mL liquid. <b>Multi-use bottle:</b> powder + liquid. Powder (fluoro-alumino-silicate glass) contains: 25-50% silicate, 20-40% aluminium oxide, 1-20% fluoride, 15-40% metal oxide, 0-15% phosphate, remainder are polyacrylic acid powder and metals in minimal quantities. Liquid (aqueous) contains: 7-25% polybasic carboxylic acid, 45-60% polyacrylic acid.
resin-based composite (low-viscosity)*	<b>Single-use applicator or multi-use bottle</b> *of any type for use as dental sealant
resin-based composite (high-viscosity)*	<b>Single-use capsule or multi-use syringe</b> *of any type for use as dental filling material
silver diamine fluoride	<b>Solution:</b> 38% w/v.

Table 1: Explanation of dosage forms

A. Principal dosage forms used in EML – oral administration

Term	Definition
Solid oral dosage form	<p>Refers to tablets or capsules or other solid dosage forms such as 'melts' that are immediate-release preparations. It implies that there is no difference in clinical efficacy or safety between the available dosage forms, and countries should therefore choose the form(s) to be listed depending on quality and availability.</p> <p>The term 'solid oral dosage form' is <i>never</i> intended to allow any type of modified-release tablet.</p>
Tablets	<p>Refers to:</p> <ul style="list-style-type: none"> <li>• uncoated or coated (film-coated or sugar-coated) tablets that are intended to be swallowed whole;</li> <li>• unscored and scored*;</li> <li>• tablets that are intended to be chewed before being swallowed;</li> <li>• tablets that are intended to be dispersed or dissolved in water or another suitable liquid before being swallowed;</li> <li>• tablets that are intended to be crushed before being swallowed.</li> </ul> <p>The term 'tablet' without qualification is <i>never</i> intended to allow any type of modified-release tablet.</p>
Tablets (qualified)	<p>Refers to a specific type of tablet:</p> <p><b>chewable</b> - tablets that are intended to be chewed before being swallowed;</p> <p><b>dispersible</b> - tablets that are intended to be dispersed in water or another suitable liquid before being swallowed;</p> <p><b>soluble</b> - tablets that are intended to be dissolved in water or another suitable liquid before being swallowed;</p> <p><b>crushable</b> - tablets that are intended to be crushed before being swallowed;</p> <p><b>scored</b> - tablets bearing a break mark or marks where sub-division is intended in order to provide doses of less than one tablet;</p> <p><b>sublingual</b> - tablets that are intended to be placed beneath the tongue.</p> <p>The term 'tablet' is <i>always</i> qualified with an additional term (in parentheses) in entries where one of the following types of tablet is intended: <b>gastro-resistant</b> (such tablets may sometimes be described as enteric-coated or as delayed-release), <b>prolonged-release</b> or another modified-release form.</p>

---

\* Scored tablets may be divided for ease of swallowing, provided that dose is a whole number of tablets.

## WHO Model List of Essential Medicines – 24th List (2025)

Capsules	Refers to hard or soft capsules.  The term 'capsule' without qualification is <i>never</i> intended to allow any type of modified-release capsule.
Capsules (qualified)	The term 'capsule' with qualification refers to <b>gastro-resistant</b> (such capsules may sometimes be described as enteric-coated or as delayed-release), <b>prolonged-release</b> or another modified-release form.
Granules	Preparations that are issued to patient as granules to be swallowed without further preparation, to be chewed, or to be taken in or with water or another suitable liquid.  The term 'granules' without further qualification is <i>never</i> intended to allow any type of modified-release granules.
Oral powder	Preparations that are issued to patient as powder (usually as single-dose) to be taken in or with water or another suitable liquid.
Oral liquid	Liquid preparations intended to be <i>swallowed</i> i.e. oral solutions, suspensions, emulsions and oral drops, including those constituted from powders or granules, but <i>not</i> those preparations intended for <i>oromucosal administration</i> e.g. gargles and mouthwashes.  Oral liquids presented as powders or granules may offer benefits in the form of better stability and lower transport costs. If more than one type of oral liquid is available on the same market (e.g. solution, suspension, granules for reconstitution), they may be interchanged and in such cases should be bioequivalent. It is preferable that oral liquids do not contain sugar and that solutions for children do not contain alcohol.

### B. Principal dosage forms used in EML – parenteral administration

Term	Definition
Injection	Refers to solutions, suspensions and emulsions including those constituted from powders or concentrated solutions.
Injection (qualified)	Route of administration is indicated in parentheses where relevant.
Injection (oily)	The term 'injection' is qualified by '(oily)' in relevant entries.
Intravenous infusion	Refers to solutions and emulsions including those constituted from powders or concentrated solutions.

### C. Other dosage forms

Mode of administration	Term to be used
To the eye	Eye drops, eye ointments.
Topical	For liquids: lotions, paints. For semi-solids: cream, ointment.
Rectal	Suppositories, gel or solution.
Vaginal	Pessaries or vaginal tablets.
Inhalation	Powder for inhalation, solution for inhalation, suspension for inhalation.

# WHO Model List of Essential Medicines – 24th List (2025)

## Index

abacavir .....	21	benzoyl peroxide .....	44
abacavir + dolutegravir + lamivudine .....	22	benzyl benzoate .....	44
abacavir + lamivudine .....	22	benzylpenicillin .....	7, 12
<i>abiraterone</i> .....	35	<i>beractant</i> .....	56
acamprosate calcium .....	58	betamethasone .....	44
acetazolamide .....	53	<i>bevacizumab</i> .....	53
acetic acid .....	61	<i>bicalutamide</i> .....	35
acetylcysteine .....	4	biperiden .....	6
acetylsalicylic acid .....	2, 7, 42, 61	bisoprolol .....	39, 40, 41
acetylsalicylic acid + atorvastatin + ramipril .....	43	<i>bleomycin</i> .....	28
acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide .....	43	<i>blinatumomab</i> .....	34
aciclovir .....	8, 20, 52	<i>bortezomib</i> .....	33
<i>adalimumab</i> .....	27, 44, 62	budesonide .....	59, 61
albendazole .....	9	budesonide + formoterol .....	59
alcohol based hand rub .....	45	bupivacaine .....	1
<i>allopurinol</i> .....	36, 61	bupropion .....	58
<i>all-trans retinoid acid (ATRA)</i> .....	33	cabergoline .....	50
<i>alprostadil (prostaglandin E1)</i> .....	56	caffeine citrate .....	56
<i>alteplase</i> .....	42	calamine .....	44
amidotrizoate .....	45	calcipotriol .....	44
amikacin .....	11, 18	calcium .....	60
amiloride .....	46	<i>calcium folinate (leucovorin calcium)</i> .....	28
<i>amiodarone</i> .....	40	calcium gluconate .....	4, 61
amitriptyline .....	3, 57	<i>capecitabine</i> .....	28
amlodipine .....	40	carbamazepine .....	5, 57
amodiaquine – sulfadoxine + pyrimethamine .....	25	carbetocin .....	55
amoxicillin .....	7, 11	<i>carboplatin</i> .....	28
amoxicillin + clavulanic acid .....	11, 18	cefalexin .....	12
amphotericin B .....	19, 24	cefazolin .....	12
ampicillin .....	7, 12	<i>cefiderocol</i> .....	17
<i>anastrozole</i> .....	35	cefixime .....	15
ansuvimab .....	26	cefotaxime .....	8, 15
anti-D immunoglobulin .....	38	<i>ceftazidime</i> .....	17
anti-rabies immunoglobulin .....	38	<i>ceftazidime + avibactam</i> .....	17
anti-rabies virus monoclonal antibodies .....	50	<i>ceftolozane + tazobactam</i> .....	17
anti-tetanus immunoglobulin .....	38	ceftioxaone .....	8, 16
antivenom immunoglobulin .....	50	cefuroxime .....	16
<i>aprepitant</i> .....	47	charcoal, activated .....	4
<i>arsenic trioxide</i> .....	28	<i>chlorambucil</i> .....	28
artemether .....	24	chloramphenicol .....	8, 12
artemether + lumefantrine .....	24	chlorhexidine .....	45, 56
artesunate .....	24	chlorine base compound .....	45
artesunate – sulfadoxine + pyrimethamine .....	24	chloroquine .....	24, 25, 61
artesunate + amodiaquine .....	24	chloroxylenol .....	46
artesunate + mefloquine .....	24	cholera vaccine .....	51
artesunate + pyronaridine .....	24	<i>ciclosporin</i> .....	27
ascorbic acid .....	60	ciprofloxacin .....	16, 61
<i>asparaginase</i> .....	28	<i>cisplatin</i> .....	29
atazanavir + ritonavir .....	21	<i>cladribine</i> .....	6
atoltivimab + maftivimab + odesivimab .....	26	clarithromycin .....	16
atorvastatin + perindopril + amlodipine .....	43	clindamycin .....	13
atracurium .....	52	clofazimine .....	18
atropine .....	1, 4, 53	<i>clomifene</i> .....	54
<i>azathioprine</i> .....	27, 61	clomipramine .....	58
azithromycin .....	15, 52	clopidogrel .....	42
<i>baclofen</i> .....	6	clotrimazole .....	19
barium sulfate .....	45	cloxacillin .....	13
BCG vaccine .....	51	<i>clozapine</i> .....	57
bedaquiline .....	18	coagulation factor IX, plasma-derived .....	39
<i>bendamustine</i> .....	28	coagulation factor IX, recombinant .....	39
benzathine benzylpenicillin .....	12	coagulation factor VIII, plasma-derived .....	39
benznidazole .....	26	coagulation factor VIII, recombinant .....	39
		coal tar .....	44

## WHO Model List of Essential Medicines – 24th List (2025)

codeine .....	2	ethambutol + isoniazid + pyrazinamide + rifampicin .....	18
colecalfiferol .....	60	ethambutol + isoniazid + rifampicin.....	18
<i>colistin</i> .....	17	ethanol .....	45
condoms .....	54	ethinylestradiol + etonogestrel.....	54
copper-containing device .....	54	ethinylestradiol + levonorgestrel.....	54
cryoprecipitate, pathogen-reduced .....	38	ethinylestradiol + norethisterone .....	54
cyclizine .....	3	ethionamide.....	18
<i>cyclophosphamide</i> .....	29	<i>ethosuximide</i> .....	6
cycloserine .....	18	etonogestrel-releasing implant .....	54
<i>cytarabine</i> .....	29	<i>etoposide</i> .....	30
cytisine (cytisinicline) .....	58	<i>everolimus</i> .....	33
dabigatran .....	37	fentanyl .....	2
<i>dacarbazine</i> .....	29	ferrous salt .....	36
daclatasvir .....	23	ferrous salt + folic acid.....	36
daclatasvir + sofosbuvir .....	23	fexinidazole .....	26
<i>dactinomycin</i> .....	29	<i>filgrastim</i> .....	34
dapsone .....	18	fluconazole .....	19
darunavir .....	21	flucytosine .....	19
<i>dasatinib</i> .....	33	<i>fludarabine</i> .....	30
<i>daunorubicin</i> .....	29	fludrocortisone.....	48
deferasirox .....	38	fluorescein.....	45
<i>deferoxamine</i> .....	5, 38	fluoride .....	62
delamanid .....	18	<i>fluorouracil</i> .....	30, 44
dengue vaccine.....	51	fluoxetine.....	3, 57, 58
desmopressin .....	37	fluphenazine .....	57
dexamethasone.....	3, 4, 35, 47, 55	folic acid.....	36
dextran 70 .....	39	<i>fomepizole</i> .....	5
diaphragms .....	54	<i>fosfomycin</i> .....	17
diazepam .....	3, 5, 58	fresh-frozen plasma .....	38
<i>diazoxide</i> .....	49	furosemide .....	42, 46
diethylcarbamazine.....	9	<i>gemcitabine</i> .....	30
digoxin .....	39, 42	gentamicin.....	8, 13, 52
dihydroartemisinin + piperazine .....	24	glass ionomer cement.....	62
diloxanide .....	23	<i>glatiramer acetate</i> .....	6
<i>dimercaprol</i> .....	5	glecaprevir + pibrentasvir.....	23
diphtheria antitoxin .....	50	gliclazide .....	49
diphtheria vaccine .....	51	glucagon .....	49
<i>docetaxel</i> .....	30	glucose .....	60
docusate sodium .....	3	glucose with sodium chloride .....	60
dolutegravir .....	21	glutaral .....	46
dolutegravir + lamivudine + tenofovir.....	22	glycerol .....	45
<i>dopamine</i> .....	42	glyceryl trinitrate .....	39
<i>doxorubicin</i> .....	30	griseofulvin .....	20
<i>doxorubicin (as pegylated liposomal)</i> .....	30	Haemophilus influenzae type b vaccine.....	51
doxycycline .....	13, 25	haloperidol .....	3, 57
Ebola vaccine .....	51	heparin sodium.....	37
efavirenz.....	21	hepatitis A vaccine .....	51
efavirenz + emtricitabine + tenofovir .....	22	hepatitis B vaccine .....	51
efavirenz + lamivudine + tenofovir .....	22	hepatitis E vaccine.....	51
eflornithine.....	26	human papilloma virus (HPV) vaccine .....	51
elexacaftor + tezacaftor + ivacaftor .....	27	hydralazine.....	40
emicizumab.....	37	hydrochlorothiazide .....	40, 42, 46
empagliflozin .....	49	hydrocortisone .....	4, 35, 44, 47, 48
emtricitabine + tenofovir .....	22	hydroxocobalamin .....	37
enalapril .....	40, 42	<i>hydroxychloroquine</i> .....	61
enoxaparin .....	37	<i>hydroxyurea (hydroxycarbamide)</i> .....	31, 38
entecavir .....	23	hyoscine butylbromide .....	3
<i>ephedrine</i> .....	1	hyoscine hydrobromide.....	3
epinephrine (adrenaline).....	4, 39, 53, 59	hypochlorous acid .....	46
equine rabies immunoglobulin.....	50	<i>ibrutinib</i> .....	33
ergocalciferol.....	60	ibuprofen .....	2, 7, 56
ergometrine.....	55	<i>ifosfamide</i> .....	31
<i>erlotinib</i> .....	33	<i>imatinib</i> .....	33
erythromycin .....	52	influenza vaccine .....	51
<i>erythropoiesis-stimulating agents</i> .....	37	insulin (analogue, long-acting) .....	49
estradiol cypionate + medroxyprogesterone acetate ....	54	insulin (analogue, rapid-acting) .....	49
ethambutol .....	18	insulin (human, intermediate-acting) .....	49

# WHO Model List of Essential Medicines – 24th List (2025)

insulin (human, short-acting).....	49	methyl dopa .....	41
<i>intraperitoneal dialysis solution</i> .....	56	<i>methylprednisolone</i> .....	35
iodine .....	60	methylthioninium chloride (methylene blue) .....	4
<i>iodine + potassium iodide (Lugol's solution)</i> .....	50	metoclopramide.....	3, 47
iohexol .....	45	metronidazole.....	14, 23
ipratropium bromide .....	59	<i>micafungin</i> .....	20
<i>irinotecan</i> .....	31	miconazole.....	43
isoflurane.....	1	midazolam.....	1, 3, 5
isoniazid .....	18	mifepristone – misoprostol .....	55
isoniazid + pyrazinamide + rifampicin .....	19	miltefosine .....	24
isoniazid + pyridoxine + sulfamethoxazole + trimethoprim.....	22	misoprostol.....	55
isoniazid + rifampicin .....	19	morphine .....	1, 2
isoniazid + rifapentine .....	19	moxifloxacin .....	19
isosorbide dinitrate .....	39	mpox vaccine .....	51
itraconazole.....	20	multiple micronutrient powder .....	60
ivacaftor .....	27	multiple micronutrient supplement.....	56
ivermectin .....	9, 26	mumps vaccine .....	51
Japanese encephalitis vaccine.....	51	mupirocin .....	43
ketamine .....	1	naloxone .....	4
lactulose.....	3	naltrexone .....	58
lamivudine .....	21	natamycin.....	52
lamivudine + zidovudine.....	22	neostigmine.....	8, 52
lamotrigine .....	5	nevirapine.....	21
latanoprost .....	53	niclosamide .....	9
ledipasvir + sofosbuvir .....	23	nicotinamide.....	60
<i>lenalidomide</i> .....	34	nicotine replacement therapy (NRT).....	58
<i>letrozole</i> .....	55	nifedipine .....	55
<i>leuporelin</i> .....	35	nifurtimox .....	26
levamisole .....	9	<i>nilotinib</i> .....	33
levetiracetam.....	5, 6	nitrofurantoin .....	14
levodopa + carbidopa .....	6	nitrous oxide.....	1
levofloxacin .....	19	norethisterone enantate.....	54
levonorgestrel.....	54	<i>normal immunoglobulin</i> .....	8, 39
levonorgestrel-releasing implant.....	54	nystatin .....	20
levonorgestrel-releasing intrauterine system.....	54	<i>octreotide</i> .....	50
levothyroxine .....	50	ofloxacin.....	52
lidocaine.....	1, 39	olanzapine .....	57
lidocaine + epinephrine (adrenaline).....	1	omeprazole .....	46
<i>linezolid</i> .....	17, 19	ondansetron .....	4, 47
lisinopril + amlodipine.....	40	oral rehydration salts .....	48, 59
lisinopril + hydrochlorothiazide .....	40	oral rehydration salts – zinc sulfate.....	48
lithium carbonate .....	57	<i>oseltamivir</i> .....	22
loperamide .....	3	<i>oxaliplatin</i> .....	31
lopinavir + ritonavir .....	21	<i>oxamniquine</i> .....	9
loratadine .....	4	oxygen .....	1, 2
lorazepam .....	5	oxytocin.....	55
losartan .....	41, 42	<i>paclitaxel</i> .....	32
magnesium sulfate.....	5	paliperidone.....	57
malaria vaccine .....	51	p-aminosalicylate sodium.....	19
mannitol .....	46	<i>pancreatic enzymes</i> .....	27, 46
measles vaccine.....	51	paracetamol (acetaminophen) .....	2, 7
mebendazole.....	9	paromomycin .....	24
medroxyprogesterone acetate .....	48, 54	<i>pegaspargase</i> .....	32
mefloquine.....	25	<i>pegfilgrastim</i> .....	34
meglumine antimoniate.....	24	<i>pembrolizumab</i> .....	34
<i>meglumine iotroxate</i> .....	45	penicillamine.....	4, 61
melarsoprol .....	26	<i>pentamidine</i> .....	25, 26
<i>melfalan</i> .....	31	perindopril + amlodipine + indapamide .....	41
meningococcal meningitis vaccine.....	51	permethrin.....	44
<i>mercaptopurine</i> .....	31	pertussis vaccine.....	51
<i>meropenem</i> .....	8, 17, 19	phenobarbital .....	5
<i>meropenem + vaborbactam</i> .....	18	phenoxymethylpenicillin .....	14
<i>mesna</i> .....	36	phenytoin .....	6
metformin.....	49	phytomenadione .....	37
<i>methadone</i> .....	2, 59	pilocarpine.....	53
methimazole.....	50	piperacillin + tazobactam .....	17
<i>methotrexate</i> .....	31, 44, 61, 62	platelets .....	38



## WHO Model List of Essential Medicines – 24th List (2025)

<i>plazomicin</i> .....	18	sodium thiosulfate.....	4, 43
pneumococcal vaccine .....	51	sofosbuvir.....	23
podophyllum resin .....	44	sofosbuvir + velpatasvir .....	23
poliomyelitis vaccine .....	51	spectinomycin .....	14
<i>polymyxin B</i> .....	18	spironolactone.....	42, 46
<i>poractant alfa</i> .....	56	<i>streptokinase</i> .....	42
potassium chloride .....	59, 60	streptomycin.....	19
potassium ferric hexacyano-ferrate(II) -2H <sub>2</sub> O .....	4	<i>succimer</i> .....	5
<i>potassium iodide</i> .....	20, 50	sulfadiazine .....	25
potassium permanganate .....	43	sulfadoxine + pyrimethamine.....	25
povidone iodine .....	45	sulfamethoxazole + trimethoprim.....	14, 25
praziquantel.....	9	sulfasalazine .....	47, 61
prednisolone.....	4, 6, 7, 35, 47, 48, 53	sumatriptan .....	7
pretomanid.....	19	sunscreen, broad-spectrum .....	45
primaquine .....	24	suramin sodium .....	26
procaine benzylpenicillin .....	14	suxamethonium .....	52
<i>procarbazine</i> .....	32	<i>tacrolimus</i> .....	27
progesterone vaginal ring .....	54	<i>tamoxifen</i> .....	36
propofol.....	1	telmisartan + amlodipine .....	41
propranolol.....	7	telmisartan + hydrochlorothiazide.....	41
propylthiouracil .....	50	tenofovir disoproxil fumarate .....	21, 23
protamine sulfate .....	37	terbinafine .....	43
pyrantel .....	9	<i>testosterone</i> .....	48
pyrazinamide .....	19	tetanus vaccine .....	52
<i>pyridostigmine</i> .....	8, 52	tetracaine .....	53
pyridoxine.....	60	tetracycline.....	53
pyrimethamine.....	25	<i>thalidomide</i> .....	34
quetiapine .....	57	thiamine .....	61
quinine .....	25	tick-borne encephalitis vaccine .....	52
rabies vaccine .....	51	timolol .....	53
raltegravir.....	22	<i>tioguanine</i> .....	32
ranitidine .....	46	tiotropium .....	59
<i>rasburicase</i> .....	36	tranexamic acid.....	37, 56
ravidasvir.....	23	<i>trastuzumab</i> .....	33
ready-to-use therapeutic food .....	36	<i>triamcinolone hexacetonide</i> .....	62
<i>realgar-Indigo naturalis formulation</i> .....	32	triclabendazole .....	9
red blood cells.....	38	trimethoprim .....	15
resin-based composite (high-viscosity).....	62	tropicamide .....	45
resin-based composite (low-viscosity) .....	62	tuberculin, purified protein derivative (PPD).....	50
respiratory syncytial virus vaccine .....	51	typhoid vaccine .....	52
retinol .....	61	ulipristal.....	54
ribavirin .....	23	urea .....	44, 45
riboflavin.....	61	<i>ustekinumab</i> .....	44
rifabutin .....	19	valganciclovir.....	22
rifampicin .....	18, 19	valproic acid (sodium valproate) .....	6, 58
rifapentine .....	19	valsartan + amlodipine + hydrochlorothiazide .....	41
risperidone .....	57	vancomycin .....	17
ritonavir .....	21	varenicline .....	58
<i>rituximab</i> .....	6, 33	varicella vaccine .....	52
rotavirus vaccine .....	51	vecuronium .....	52
rubella vaccine .....	51	verapamil .....	7, 39, 40
salbutamol.....	59	<i>vinblastine</i> .....	32
salicylic acid .....	44	<i>vincristine</i> .....	32
selenium sulfide.....	43	<i>vinorelbine</i> .....	32
semaglutide.....	49	voriconazole .....	20
senna .....	4, 47	warfarin .....	37
sevoflurane.....	1	water for injection .....	60
silver diamine fluoride .....	62	whole blood .....	38
silver sulfadiazine.....	43	xylometazoline.....	61
simvastatin .....	43	yellow fever vaccine.....	52
<i>sodium calcium edetate</i> .....	5	zidovudine .....	21
sodium chloride .....	60	zinc sulfate .....	48
sodium hydrogen carbonate .....	60	<i>zoledronic acid</i> .....	36
sodium lactate.....	60		
sodium nitrite.....	4		
<i>sodium nitroprusside</i> .....	41		
sodium stibogluconate.....	24		

**Essential Medicines List Secretariat**

Department of Health Products Policy and Standards  
World Health Organization  
20, Avenue Appia  
1211 Geneva 27  
Switzerland  
Email: [emlsecretariat@who.int](mailto:emlsecretariat@who.int)