

# Part III – Introduction to Additive Combinatorics (Incomplete)

Based on lectures by Prof Julia Wolf  
Notes taken by Yaël Dillies

Lent 2024

## Contents

<b>1</b>	<b>Fourier-analytic techniques</b>	<b>2</b>
----------	------------------------------------	----------

# 1 Fourier-analytic techniques

## Lecture 1

Let  $G = \mathbb{F}_p^n$  where  $p$  is a small fixed prime and  $n$  is large.

**Notation.** Given a finite set  $B$  and any function  $f : B \rightarrow \mathbb{C}$ , write

$$\mathbb{E}_{x \in B} f(x) = \frac{1}{|B|} \sum_{x \in B} f(x)$$

Write  $\omega = e^{\frac{\pi i}{p}}$ . Note  $\sum_{a \in \mathbb{F}_p} \omega^a = 0$ .

**Definition 1.1.** Given  $f : \mathbb{F}_p^n \rightarrow \mathbb{C}$ , define its **Fourier transform**  $\hat{f} : \mathbb{F}_p^n \rightarrow \mathbb{C}$  by

$$\hat{f}(t) = \mathbb{E}_{x \in \mathbb{F}_p^n} f(x) \omega^{x \cdot t}$$

It is easy to verify the **inversion formula**

$$f(x) = \sum_{t \in \mathbb{F}_p^n} \hat{f}(t) \omega^{-x \cdot t}$$

Indeed,

$$\begin{aligned} \sum_{t \in \mathbb{F}_p^n} \hat{f}(t) \omega^{-x \cdot t} &= \sum_{t \in \mathbb{F}_p^n} (\mathbb{E}_y f(y) \omega^{y \cdot t}) \omega^{-x \cdot t} \\ &= \mathbb{E}_y f(y) \sum_t \omega^{(y-x) \cdot t} \\ &= \mathbb{E}_y f(y) 1_{y=x} p^n \\ &= f(x) \end{aligned}$$

**Notation.** Given a set  $A$  of a finite group  $G$ , write

- $1_A$  the *characteristic function* of  $A$ , ie

$$1_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

- $\mu_A$  the *characteristic measure* of  $A$ , ie

$$\mu_A = \alpha^{-1} 1_A$$

where  $\alpha = \frac{|A|}{|G|}$ .

- $f_A$  the *balanced function* of  $A$ , ie

$$f_A(x) = 1_A(x) - \alpha$$

Note  $\mathbb{E}_x f_A(x) = 0$ ,  $\mathbb{E}_x \mu_A(x) = 1$ ,  $\widehat{1_A}(0) = \mathbb{E}_x 1_A(x) = \alpha$ . Writing  $-A = \{-a | a \in A\}$ , we have

$$\begin{aligned} \widehat{1_{-A}}(t) &= \mathbb{E}_x 1_{-A}(x) \omega^{x \cdot t} \\ &= \mathbb{E}_x 1_A(-x) \omega^{x \cdot t} \\ &= \mathbb{E}_x 1_A(x) \omega^{-x \cdot t} \\ &= \overline{\widehat{1_A}(t)} \end{aligned}$$

**Example 1.2.** Let  $V \leq \mathbb{F}_p^n$ . Then

$$\widehat{1_V}(t) = \mathbb{E}_x 1_V(x) \omega^{x \cdot t} = \frac{|V|}{|G|} 1_{V^\perp}(t)$$

So

$$\widehat{\mu_V}(t) = 1_{V^\perp}(t)$$

**Example 1.3.** Let  $R \subseteq \mathbb{F}_p^n$  be such that each  $x$  is included with probability  $\frac{1}{2}$  independently. Then with high probability

$$\sup_{t \neq 0} |\widehat{1_R}(t)| = O\left(\sqrt{\frac{\log(p^n)}{p^n}}\right)$$

This is on Example Sheet 1 using a **Chernoff-type bound**: Given  $\mathbb{C}$ -valued independent random variables  $X_1, \dots, X_n$  with mean 0 and  $\theta \geq 0$ , we have

$$\mathbb{P}\left(\left|\sum_i X_i\right| \geq \theta \sqrt{\sum_i \|X_i\|_{L^\infty}^2}\right) \leq 4 \exp\left(-\frac{\theta^2}{4}\right)$$

**Example 1.4.** Let  $Q = \{x \in \mathbb{F}_p^n \mid x \cdot x = 0\}$ . Then  $|Q| = \left(\frac{1}{p} + O(p^{-n})\right) p^n$  and  $\sup_{t \neq 0} |\widehat{1_Q}(t)| = O(p^{-\frac{n}{2}})$ . See Example Sheet 1.

**Notation.** Given  $f, g : \mathbb{F}_p^n \rightarrow \mathbb{C}$ , write

$$\begin{aligned} \langle f, g \rangle &= \mathbb{E}_x f(x) \overline{g(x)} \\ \langle \hat{f}, \hat{g} \rangle &= \sum_t \hat{f}(t) \overline{\hat{g}(t)} \end{aligned}$$

Consequently,

$$\begin{aligned} \|f\|_2^2 &= \mathbb{E}_x |f(x)|^2 \\ \|\hat{f}\|_2^2 &= \sum_t |\hat{f}(t)|^2 \end{aligned}$$

**Lemma 1.5.** For all  $f, g : \mathbb{F}_p^n \rightarrow \mathbb{C}$ ,

$$\begin{aligned} \langle f, g \rangle &= \langle \hat{f}, \hat{g} \rangle && \text{(Plancherel)} \\ \|f\|_2 &= \|\hat{f}\|_2 && \text{(Parseval)} \end{aligned}$$

*Proof.* Exercise. □

**Definition 1.6.** Let  $\rho > 0$  and  $f : \mathbb{F}_p^n \rightarrow \mathbb{C}$ . Define the  $\rho$ -large spectrum of  $f$  to be

$$\text{Spec}_\rho(f) = \{t \mid |\hat{f}(t)| \geq \rho \|f\|_1\}$$

**Example 1.7.** By Example 1.2, if  $V \leq \mathbb{F}_p^n$ , then  $\text{Spec}_\rho(1_V) = V^\perp$  for all  $\rho > 0$ .

**Lemma 1.8.** For all  $\rho > 0$ ,  $|\text{Spec}_\rho(f)| \leq \rho^{-2} \frac{\|f\|_2^2}{\|f\|_1^2}$ .

*Proof.*

$$\|f\|_2^2 = \|\hat{f}\|_2^2 \geq \sum_{t \in \text{Spec}_\rho(f)} |\hat{f}(t)|^2 \geq |\text{Spec}_\rho(f)| (\rho \|f\|_1)^2$$

□

## Lecture 2

**Definition 1.9.** Given  $f, g : \mathbb{F}_p^n \rightarrow \mathbb{C}$ , define their **convolution**  $f * g : \mathbb{F}_p^n \rightarrow \mathbb{C}$  by

$$(f * g)(x) = \mathbb{E}_y f(y) g(x - y)$$

**Example 1.10.** Given  $A, B \subseteq \mathbb{F}_p^n$ ,

$$\begin{aligned} (1_A * 1_B)(x) &= \mathbb{E}_y 1_A(y) 1_B(x - y) \\ &= \frac{1}{p^n} |A \cap (x - B)| \\ &= \frac{\# \text{ ways to write } x = a + b, a \in A, b \in B}{p^n} \end{aligned}$$

In particular, the support of  $1_A * 1_B$  is the **sum set**

$$A + B = \{a + b \mid a \in A, b \in B\}$$

**Lemma 1.11.** Given  $f, g : \mathbb{F}_p^n \rightarrow \mathbb{C}$ ,

$$\widehat{f * g}(t) = \hat{f}(t) \hat{g}(t)$$

*Proof.*

$$\begin{aligned} \widehat{f * g}(t) &= \mathbb{E}_x (\mathbb{E}_y f(y) g(x - y)) \omega^{x \cdot t} \\ &= \mathbb{E}_y f(y) \mathbb{E}_u g(u) \omega^{(u+y) \cdot t} \\ &= \hat{f}(t) \hat{g}(t) \end{aligned}$$

□

**Example 1.12.**  $\|\hat{f}\|_4^4 = \mathbb{E}_{x+y=z+w} f(x) f(y) \overline{f(z)} \overline{f(w)}$ . See Example Sheet 1.

**Lemma 1.13** (Bogolyubov). If  $A \subseteq \mathbb{F}_p^n$  is of density  $\alpha > 0$ , then there exists a subspace  $V$  of codimension at most  $2\alpha^{-2}$  such that  $V \subseteq (A + A) - (A + A)$ .

*Proof.* Observe that  $(A + A) - (A + A) = \text{supp}(\underbrace{1_A * 1_A * 1_{-A} * 1_{-A}}_g)$ , so we wish to find

$V$  such that  $g(x) > 0$  for all  $x \in V$ . Let  $K = \text{Spec}_\rho(1_A)$  for some  $\rho > 0$  and define  $V = \langle K \rangle^\perp$ . By Lemma 1.8,  $\text{codim } V \leq |K| \leq \rho^{-2} \alpha^{-1}$ . We calculate

$$\begin{aligned} g(x) &= \sum_{t \in \mathbb{F}_p^n} 1_A * 1_A * \widehat{1_{-A}} * 1_{-A}(t) \omega^{-x \cdot t} \\ &= \sum_{t \in \mathbb{F}_p^n} |\widehat{1_A}(t)|^4 \omega^{-x \cdot t} \\ &= \underbrace{\alpha^4 + \sum_{t \in K \setminus \{0\}} |\widehat{1_A}(t)|^4 \omega^{-x \cdot t}}_{(1)} + \underbrace{\sum_{t \notin K} |\widehat{1_A}(t)|^4 \omega^{-x \cdot t}}_{(2)} \end{aligned}$$

We now see that

$$(1) = \sum_{t \in K \setminus \{0\}} \left| \widehat{1_A}(t) \right|^4 \geq 0$$

and

$$|(2)| \leq \sum_{t \notin K} \left| \widehat{1_A}(t) \right|^4 \leq \sup_{t \notin K} \left| \widehat{1_A}(t) \right|^2 \sum_{t \notin K} \left| \widehat{1_A}(t) \right|^2 \leq (\rho\alpha)^2 \|1_A\|_2^2 = \rho^2 \alpha^3$$

by Parseval. Picking  $\rho = \sqrt{\frac{\alpha}{2}}$ , we thus get  $\rho^2 \alpha^3 \leq \frac{\alpha^4}{2}$  and  $g(x) > 0$  whenever  $x \in V$ .  $\square$

**Example 1.14.** The set  $A = \{x \in \mathbb{F}_2^n \mid |x| \geq \frac{n}{2} + \frac{\sqrt{n}}{2}\}$  has density at least  $\frac{1}{4}$  but there is no coset  $C$  of any subspace of codimension  $\sqrt{n}$  such that  $C \subseteq A + A$ . See Example Sheet 1.

**Lemma 1.15.** Let  $A \subseteq \mathbb{F}_p^n$  of density  $\alpha$  be such that  $\text{Spec}_\rho(1_A)$  contains some  $t \neq 0$ . Then there exist  $V \leq \mathbb{F}_p^n$  of codimension 1 and  $x \in \mathbb{F}_p^n$  such that

$$|A \cap (x + V)| \geq \alpha \left(1 + \frac{\rho}{2}\right) |V|$$

*Proof.* Let  $t \neq 0$  be such that  $\left| \widehat{1_A}(t) \right| \geq \rho\alpha$  and let  $V = \langle t \rangle^\perp$ . For  $j = 1, \dots, p$ , write

$$v_j + V = \{x \in \mathbb{F}_p^n \mid x \cdot t = j\}$$

the cosets of  $V$ . Then

$$\begin{aligned} \widehat{1_A}(t) &= \widehat{f_A}(t) \\ &= \mathbb{E}_{x \in \mathbb{F}_p^n} (1_A(x)) - \alpha \omega^{x \cdot t} \\ &= \mathbb{E}_j \omega^j \mathbb{E}_{x \in v_j + V} (1_A(x) - \alpha) \\ &= \mathbb{E}_j a_j \omega^j \end{aligned}$$

where  $a_j = \frac{|A \cap (v_j + V)|}{|V|} - \alpha$ . Since  $\sum_j a_j = 0$ , we get

$$\rho\alpha \leq \left| \widehat{1_A}(t) \right| \leq \mathbb{E}_j |a_j| = \mathbb{E}_j (|a_j| + a_j)$$

So there is some  $j$  such that  $|a_j| + a_j \geq \rho\alpha$ . In particular, this  $a_j$  is positive, so

$$\frac{|A \cap (v_j + V)|}{|V|} \geq \alpha + \frac{\rho\alpha}{2}$$

as wanted.  $\square$

### Lecture 3

**Lemma 1.16.** Let  $p \geq 3$  and  $A \subseteq \mathbb{F}_p^n$  of density  $\alpha > 0$  be such that  $\sup_{t \neq 0} \left| \widehat{1_A}(t) \right| = o(1)$ . Then  $A$  contains  $(\alpha^3 + o(1)) |G|^2$  three terms arithmetic progressions (aka 3AP).

**Notation.** Given  $f, g, h : \mathbb{F}_p^n \rightarrow \mathbb{C}$ , write

$$T_3(f, g, h) = \mathbb{E}_x f(x) g(x + d) h(x + 2d)$$

Given  $A \subseteq \mathbb{F}_p^n$ , write  $2 \cdot A = \{2a \mid a \in A\}$ . This is distinct from  $2A = \{a + b \mid a, b \in A\}$ .

*Proof.* The number of 3AP (including the trivial ones of the form  $a, a, a$ ) in  $A$  is  $|G|^2$  times

$$\begin{aligned}
T_3(1_A, 1_A, 1_A) &= \mathbb{E}_{x,d} 1_A(x) 1_A(x+d) 1_A(x+2d) \\
&= \mathbb{E}_{x,y} 1_A(x) 1_A(y) 1_A(2y-x) \\
&= \mathbb{E}_y (1_A * 1_A)(2y) 1_A(y) \\
&= \langle 1_A * 1_A, 1_{2 \cdot A} \rangle \\
&= \langle \widehat{1_A}^2, \widehat{1_{2 \cdot A}} \rangle \\
&= \alpha^3 + \sum_{t \neq 0} \widehat{1_A}(t)^2 \overline{\widehat{1_{2 \cdot A}}(t)} \text{ by Plancherel}
\end{aligned}$$

In absolute value, the error term is at most

$$\sup_{t \neq 0} |\widehat{1_{2 \cdot A}}(t)| \sum_t |\widehat{1_A}(t)|^2 = \alpha \sup_{t \neq 0} |\widehat{1_A}(t)|$$

□

**Theorem 1.17** (Meshulam). Let  $p \geq 3$  and  $A \subseteq \mathbb{F}_p^n$  be a set containing only trivial 3APs. Then

$$|A| = O\left(\frac{p^n}{\log(p^n)}\right)$$

*Proof.* By assumption,  $T_3(1_A, 1_A, 1_A) = \frac{\alpha}{p^n}$ . But, as in Lemma 1.16,

$$|T_3(1_A, 1_A, 1_A) - \alpha^3| \leq \alpha \sup_{t \neq 0} |\widehat{1_A}(t)|$$

Hence, provided that  $2\alpha^{-2} \leq p^n$ , Lemma 1.15 gives us a subspace  $V \leq \mathbb{F}_p^n$  of codimension 1 and  $x \in \mathbb{F}_p^n$  such that

$$|A \cap (x + V)| \geq \alpha \left(1 + \frac{\alpha^2}{4}\right) |V|$$

We iterate this observation. Let  $A_0 = A, V_0 = \mathbb{F}_p^n$ . At step  $i$ , we are given a set  $A_i \subseteq V_i$  of density  $\alpha_i$  with only trivial 3APs. Provided that  $2\alpha_i^{-2} \leq p^{\dim V_i}$ , find  $V_{i+1} \leq V_i$  of codimension 1 and  $x \in V_i$  such that  $|A_i \cap (x + V_i)| \geq \left(\alpha_i + \frac{\alpha_i^2}{4}\right) |V_{i+1}|$  and set  $A_{i+1} = (A_i - x) \cap V_i$ . Note that  $\alpha_{i+1} \geq \alpha_i + \frac{\alpha_i^2}{4}$  and  $A_{i+1}$  only contains trivial 3APs (because, very importantly, 3AP are **translation-invariant**). Through this iteration, the density of  $A$  increases from  $\alpha$  to  $2\alpha$  in at most  $\lceil 4\alpha^{-1} \rceil$  steps, from  $2\alpha$  to  $4\alpha$  in at most  $\lceil 2\alpha^{-1} \rceil$  steps, etc... Since density can't increase past 1, it takes at most

$$\underbrace{\lceil 4\alpha^{-1} \rceil + \lceil 2\alpha^{-1} \rceil + \dots}_{\lceil \log \alpha^{-1} \rceil \text{ terms}} \leq (4\alpha^{-1} + 1) + (2\alpha^{-1} + 1) + \dots \leq 8\alpha^{-1} + \log \alpha^{-1} + 1 \leq 9\alpha^{-1}$$

steps to reach a point where the condition  $2\alpha_i^{-2} \leq p^{\dim V_i}$  is not respected anymore. Now either  $\alpha \leq \sqrt{2}p^{-\frac{n}{4}}$  (in which case the inequality is obvious) or  $\alpha \geq \sqrt{2}p^{-\frac{n}{4}}$  and

$$p^{n-9\alpha^{-1}} \leq p^{\dim V_i} \leq 2\alpha_i^{-2} \leq 2\alpha^{-2} \leq p^{\frac{n}{2}}$$

namely  $\alpha \leq \frac{18}{n}$ , as wanted. □

We have proved that if  $A \subseteq \mathbb{F}_3^n$  only contains trivial 3APs then  $|A| = O(\frac{3^n}{n})$ . The largest known set in  $\mathbb{F}_3^n$  with only trivial 3APs has size  $\geq 2.218^n$  (Tyrrell, 2022). We will return to this later.

From now on, let  $G$  be a finite abelian group.  $G$  comes equipped with a set of **characters**, ie group homomorphisms  $\gamma : G \rightarrow \mathbb{C}^\times$ . Characters themselves form a group denoted  $\hat{G}$  and called the **Pontryagin dual** (aka **dual group**) of  $G$ . It turns out that if  $G$  is finite abelian then  $\hat{\hat{G}} \cong G$  (but *non-canonically*). For instance,

- If  $G = \mathbb{F}_p^n$ , then  $\hat{G} = \{\gamma_t : x \mapsto \omega^{x \cdot t} \mid t \in G\}$
- If  $G = \mathbb{Z}/n\mathbb{Z}$ , then  $\hat{G} = \{\gamma_t : x \mapsto \omega^{xt} \mid t \in G\}$

The latter is a special case of the former, but again  $n$  should be thought of as an asymptotic variable.

**Definition 1.18.** Given  $f : G \rightarrow \mathbb{C}$ , define its **Fourier transform**  $\hat{f} : \hat{G} \rightarrow \mathbb{C}$  by

$$\hat{f}(\gamma) = \mathbb{E}_{x \in G} f(x) \gamma(x)$$

It is easy to verify that  $f(x) = \sum_{\gamma \in \hat{G}} \hat{f}(\gamma) \overline{\gamma(x)}$ . Similarly, Definitions 1.6, 1.9, Examples 1.3, 1.10 and Lemmas 1.5, 1.8, 1.11 go through in this more general context.

**Example 1.19.** Let  $p$  be a prime,  $L < p$  be even and  $J = [-\frac{L}{2}, \frac{L}{2}] \subseteq \mathbb{Z}/p\mathbb{Z}$ . Then for all  $t \neq 0$  we have

$$\widehat{1_J}(t) \leq \min\left(\frac{L+1}{p}, \frac{1}{2|t|}\right)$$

See Example Sheet 1.

**Theorem 1.20** (Roth). Let  $A \subseteq [N]$  be a set containing only trivial 3APs. Then  $|A| = O(\frac{N}{\log \log N})$ .

**Lemma 1.21.** Let  $A \subseteq [N]$  of density  $\alpha > 0$  containing only trivial 3APs and satisfying  $N > 50\alpha^{-2}$ . Let  $p$  be a prime in  $[\frac{N}{3}, \frac{2N}{3}]$  and write  $A' = A \cap [p] \subseteq \mathbb{Z}/p\mathbb{Z}$ . Then either

1.  $\sup_{t \neq 0} |\widehat{1_A}(t)| \geq \frac{\alpha^2}{10}$  (where the Fourier coefficients are computed in  $\mathbb{Z}/p\mathbb{Z}$ )
2. or there exists an interval  $J$  of length  $\geq \frac{N}{3}$  such that

$$|A \cap J| \geq \alpha \left(1 + \frac{\alpha}{400}\right) |J|$$

*Proof.* If  $|A'| \leq \alpha \left(1 - \frac{\alpha}{200}\right) p$ , then

$$|A \cap [p+1, N]| \geq \alpha(N-p) + \frac{\alpha^2 p}{200} \geq \alpha \left(1 + \frac{\alpha}{400}\right) (N-p)$$

and we are in Case 2 with  $J = [p+1, N]$ . Let  $A'' = A' \cap [\frac{p}{3}, \frac{2p}{3}]$ . Note that all 3APs of the form  $(x, x+d, x+2d) \in A' \times A'' \times A''$  are in fact 3APs in  $[N]$  (and in particular they are trivial).

If  $|A' \cap [\frac{p}{3}, \frac{2p}{3}]|$  or  $|A' \cap [\frac{2p}{3}, p]|$  were at least  $\frac{2}{5} |A'|$ , then we would again be in Case 2. We may therefore assume that  $|A''| \geq \frac{|A'|}{5}$ .

Now, as in Lemma 1.16 and Theorem 1.17 with  $\alpha' = \frac{|A'|}{p}$ ,  $\alpha'' = \frac{|A''|}{p}$ ,

$$\frac{\alpha''}{p} = T_3(1_{A'}, 1_{A''}, 1_{A''}) = \alpha' \alpha''^2 + \sum_{t \neq 0} \widehat{1_{A'}}(t) \widehat{1_{A''}}(t) \overline{\widehat{1_{2 \cdot A'}}(t)}$$

So, as before,  $\frac{\alpha'\alpha''}{2} \leq \alpha'' \sup_{t \neq 0} |\widehat{1_{A'}}(t)|$ , provided  $\frac{\alpha''}{p} \leq \frac{\alpha'\alpha''/2}{2}$ . This holds by assumption since  $p \geq \frac{N}{3}$ ,  $N \geq 50\alpha^{-2}$ ,  $\alpha' \geq \frac{199}{200}\alpha$ ,  $\alpha'' \geq \frac{\alpha'}{5}$ .  $\square$

## Lecture 5

We now want to convert the large Fourier coefficient into a density increment. This is harder now that the number of values of  $xt$  grows as  $n \rightarrow \infty$ . Compare this to the finite field case where  $x \cdot t$  only take  $p$  different values regardless of  $n$ . If we can't find a single big coefficient, then we might instead be able to find an interval of coefficients whose total contribution is big.

TODO: Insert picture

**Lemma 1.22.** Let  $m \in \mathbb{N}$  and  $\phi : [m] \rightarrow \mathbb{Z}/p\mathbb{Z}$  be multiplication by some fixed  $t \neq 0$ . Given  $\varepsilon > 0$ , there exists a partition of  $[m]$  into progressions  $P_i$  of length  $\in [\frac{\varepsilon\sqrt{m}}{2}, \varepsilon\sqrt{m}]$  such that  $\text{diam}(\phi(P_i)) \leq \varepsilon p$ .

*Proof.* Let  $u = \lfloor \sqrt{m} \rfloor$  and consider  $0, t, \dots, ut$ . By pigeonhole, find  $0 \leq v < w \leq u$  such that  $|wt - vt| \leq \frac{p}{u}$ . Set  $s = w - v \leq u$  so that  $|st| \leq \frac{p}{u}$ . Divide  $[m]$  into residue classes mod  $s$ . Each has size at least  $\lfloor \frac{m}{s} \rfloor \geq \lfloor \frac{m}{u} \rfloor$  and can be divided into progressions of the form  $a, a+s, \dots, a+ds$  with  $\frac{\varepsilon u}{2} < d \leq \varepsilon u$ . The diameter of each progression under  $\phi$  is  $|dst| \leq \varepsilon p$ .  $\square$

**Lemma 1.23.** Let  $A \subseteq [N]$  be of density  $\alpha > 0$ . Let  $p$  be a prime in  $[\frac{N}{3}, \frac{2N}{3}]$  and write  $A' = A \cap [p]$ . Suppose there exists  $t \neq 0$  such that  $|\widehat{1_{A'}}(t)| \geq \frac{\alpha^2}{10}$ . Then there exists a progression  $P$  of length at least  $\alpha^2 \frac{\sqrt{N}}{500}$  such that

$$|A \cap P| \geq \alpha \left(1 + \frac{\alpha}{50}\right) |P|$$

*Proof.* Let  $\varepsilon = \frac{\alpha^2}{40\pi}$  and use Lemma 1.22 to partition  $[p]$  into progressions  $P_i$  of length at least  $\frac{\varepsilon\sqrt{p}}{2} \geq \frac{\alpha^2}{80\pi} \sqrt{\frac{N}{3}} \geq \frac{\alpha^2\sqrt{N}}{500}$  and  $\text{diam} \phi(P_i) \leq \varepsilon p$ . Fix one  $x_i$  inside each  $P_i$ .

$$\begin{aligned} \frac{\alpha^2}{10} &\leq |\widehat{f_{A'}}(t)| \\ &= \frac{1}{p} \left| \sum_i \sum_{x \in P_i} f_{A'}(x) \omega^{xt} \right| \\ &= \frac{1}{p} \left| \sum_i \sum_{x \in P_i} f_{A'}(x) \omega^{x_i t} + \sum_i \sum_{x \in P_i} f_{A'}(x) (\omega^{xt} - \omega^{x_i t}) \right| \\ &\leq \frac{1}{p} \sum_i \left| \sum_{x \in P_i} f_{A'}(x) \omega^{x_i t} \right| + \frac{1}{p} \sum_i \sum_{x \in P_i} |f_{A'}(x)| 2\pi\varepsilon \\ &\leq \frac{1}{p} \sum_i \left| \sum_{x \in P_i} f_{A'}(x) \omega^{x_i t} \right| + \frac{\alpha^2}{20} \end{aligned}$$

So

$$\sum_i \left| \sum_{x \in P_i} f_{A'}(x) \right| \geq \frac{\alpha^2 p}{20}$$

Since  $f_{A'}$  has mean zero, there exists  $i$  such that  $\sum_{x \in P_i} f_{A'}(x) \geq \frac{\alpha^2 |P_i|}{40}$ .  $\square$



*Proof of Roth's theorem.* Put the ingredients together, Similarly to Meshulam. See Example Sheet 1 for details.  $\square$

**Example 1.24** (Behrend's construction). There exists a set  $A \subseteq [N]$  containing nontrivial 3APs of size at least  $e^{-O(\sqrt{\log n})}$ . See Example Sheet 1.

**Definition 1.25.** Let  $\Gamma \subseteq \hat{G}$ . The **Bohr set of frequencies**  $\Gamma$  and width  $\rho$  is

$$B(\Gamma, \rho) = \{x \in G \mid \forall \gamma \in \Gamma, |\gamma(x) - 1| \leq \rho\}$$

$|\Gamma|$  is the **rank** of the Bohr set.

**Example 1.26.** When  $G = \mathbb{F}_p^n$ ,  $B(\Gamma, \rho) = \langle \Gamma \rangle^\perp$  for all small enough  $\rho$  (depending only on  $p$ , not  $n$ ).

**Lemma 1.27.** Let  $B$  be a Bohr set of rank  $d$  and width  $\rho$ . Then  $|B| \geq \left(\frac{\rho}{2\pi}\right)^d |G|$ .

*Proof.* See Example Sheet 2.  $\square$

**Lemma 1.28** (Bogolyubov). Given  $A \subseteq \mathbb{Z}/p\mathbb{Z}$  of density  $\alpha > 0$ , there exists  $\Gamma \subseteq \widehat{\mathbb{Z}/p\mathbb{Z}}$  of size at most  $2\alpha^{-2}$  such that  $B(\Gamma, \frac{1}{2}) \subseteq (A + A) - (A + A)$ .