

CATHOLIC SECONDARY SCHOOLS ASSOCIATION OF NSW

	Centre Number					
			CCII		anioci	

DO NOT REMOVE PAPER FROM EXAM ROOM

2020
TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics Extension 2

Morning Session Thursday, 20 August 2020

General Instructions

- Reading time 10 minutes
- Working time 3 hours
- Write using black pen
- Calculators approved by NESA may be used
- · A Reference Sheet is provided
- In Questions 11 16, show relevant mathematical reasoning and/or calculations
- Write your Centre Number and Student Number at the top of this page

Total marks - 100

Section I Pages 2 - 5

10 marks

- Attempt Questions 1 10
- Allow 15 minutes for this section

Section II

Pages 6 - 11

90 marks

- Attempt Ouestions 11 16
- Allow about 2 hours and 45 minutes for this section

Disclaimer

Every effort has been made to prepare these "Tria" Higher Schoul Certificate Examinations in accordance with the NESA documents, Principles for Setting HSC Examinations in a Standards-Referenced Framework and Principles for Developing Marking Certificate Examinations in a Standards Referenced Framework. No gueranteer or warranty is made in middle faith to Trial. Examination parees mirror in every respect the actual RSC Examination question paper in any and all courses to be examined. These papers do not constitute advice inc can they be construed

Section I

10 marks Attempt Questions 1–10 Allow about 15 minutes for this section

Use the Multiple-Choice Answer Sheet for Questions 1-10.

- 1 Let $z = 1 + \sqrt{3}i$. What is z in exponential form?
 - (A) $e^{\frac{i\pi}{3}}$
 - (B) $e^{\frac{i\pi}{6}}$
 - (C) $2e^{\frac{it}{2}}$
 - (D) $2e^{\frac{i\pi}{6}}$
- What is the distance of the point (2, 3, 7) from the x-z plane?
 - (A) 2 units
 - (B) 3 units
 - (C) 7 units
 - (D) 9 units
- Consider the conditional statement for n > 2: If n is a prime number, then n is odd.

What is the contrapositive of the conditional statement?

- (A) If n is prime, then n is not odd.
- (B) If n is not prime, then n is not odd.
- (C) If n is odd, then n is not prime.
- (D) If n is not odd, then n is not prime.

- 4 Which of the following expressions is equal to $\int \frac{1}{x(\log_e x)^2}$?
 - (A) $\frac{1}{\log_e x} + c$
 - (B) $\frac{1}{\left(\log_e x\right)^3} + c$
 - (C) $\log_{e} \left(\frac{1}{x}\right) + c$
 - (D) $\frac{-1}{\log_{_{u}} x} + c$
- 5 Let $\alpha = 1 i$.

Which of the following is true about the value of α^{10} ?

- (A) It is purely real
- (B) It is purely imaginary
- (C) (
- (D) $32\left(\cos\frac{5\pi}{2} + i\sin\frac{5\pi}{2}\right)$
- 6 Which of the following statements does not have a counter-example?
 - (A) If a person does not own a pet, then they do not own a cat.
 - (B) A quadrilateral is formed by joining any four points in a plane.
 - (C) All primes are odd.
 - (D) If x is even, then x^2 is odd.

A particle of unit mass travels horizontally through a medium. When time t = 0, the particle is at point O with initial speed U. The resistance on the particle due to the medium is kv^2 , where v is the velocity of the particle at time t and k is a positive constant.


Which expression gives the correct velocity of the particle?

- (A) $\frac{1}{v} = kt + \frac{1}{U}$
- (B) $v = kt + \frac{1}{U}$
- (C) $\frac{1}{v} = kt$
- (D) v = kt
- 8 Which expression is equal to $\int \frac{dx}{\sqrt{8-2x-x^2}}?$
 - (A) $\sin^{-1}\left(\frac{1-x}{2\sqrt{2}}\right) + c$
 - (B) $\sin^{-1}\left(\frac{1-x}{3}\right) + c$
 - (C) $\sin^{-1}\left(\frac{1+x}{2\sqrt{2}}\right) + c$
 - (D) $\sin^{-1}\left(\frac{1+x}{3}\right) + c$

A particle is moving in simple harmonic motion about the origin according to the equation $x = 3\cos nt$, where x metres is its displacement after t seconds.

Given that the particle passes through the origin with a speed of $\sqrt{3}$ ms⁻¹, what is the period of the motion?

- (A) $\frac{2\sqrt{3}\pi}{3}$ seconds
- (B) $\frac{2\sqrt{3}}{3\pi}$ seconds
- (C) $\frac{6\pi}{\sqrt{3}}$ seconds
- (D) $\frac{6}{\sqrt{3}\pi}$ seconds
- The locus of z is displayed on the Argand diagram below.



Which of the following is the equation of the locus of z?

- (A) $\arg\left(\frac{z-i}{z-1}\right) = 0$
- (B) $\operatorname{arg}\left(\frac{z-i}{z-1}\right) = \pm \pi$
- (C) $\arg\left(\frac{z+i}{z+1}\right) = 0$
- (D) $\arg\left(\frac{z+i}{z+1}\right) = \pm \pi$

End of Section I

Section II

90 marks

Attempt Questions 11-16

Allow about 2 hours and 45 minutes for this section

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11 - 16, your responses should include relevant mathematical reasoning and/or calculations.

Question 11 (15 marks) Use a SEPARATE writing booklet.

(a) Let
$$z = 1 + i$$
 and $w = \cos \frac{\pi}{6} + i \sin \frac{\pi}{6}$.

- (i) Express $\frac{w}{z}$ in polar form.
- (ii) Hence or otherwise, express $(w\overline{z})^8$ in the form a+ib, where a and b are real.

(b) Let
$$\underline{a} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$$
 and $\underline{b} = \begin{pmatrix} -1 \\ 1 \\ 2 \end{pmatrix}$.

What is the angle between the vectors to the nearest degree?

(c) Let
$$P(z) = z^4 - 4z^3 - 3z^2 + 50z - 52$$
.

Solve P(z) = 0 if z = 3 - 2i is a root of the polynomial.

(d) A line passes through the points
$$A(1, 3, -2)$$
 and $B(2, -1, 5)$.

- (i) Show that the vector equation of the line AB is given by: $\underline{r} = (\underline{i} + 3j 2\underline{k}) + \lambda_1 (\underline{i} 4j + 7k) , \lambda_1 \in \mathbb{R}$
- (ii) Determine if the point C(3, 4, 9) lies on the line.
- (iii) Consider a line with parametric equations x=1-λ₂, y=2+3λ₂, z=-1+λ₂.
 Assuming this line is neither parallel nor perpendicular to AB, determine whether the lines intersect or are skew.

End of Question 11

2

2

1

Question 12 (15 marks) Use a SEPARATE writing booklet.

- (a) Consider the equation $z^2 2(1+2i)z + (1+i) = 0$.
 - (i) Show that $(z-(1+2i))^2 = -4+3i$.

1

(ii) Hence solve $z^2 - 2(1+2i)z + (1+i) = 0$.

2

3

(b) Sketch the intersection of the regions defined by

$$|z-2i| \le 1$$
 and $0 < Arg(z-2i) \le \frac{3\pi}{4}$.

- (c) Use the substitution $t = \tan \frac{x}{2}$ to show that $\int_0^{\frac{\pi}{3}} \frac{1}{8\sin x + 6\cos x 10} dx = \frac{1}{2} \left(\frac{1}{2 \sqrt{3}} \right).$
- (d) A function is defined by $f(x) = \frac{4x}{5-x}$. Solve the inequality $f(|x|) \le 2$.

3

(e) It is given that p and q are real numbers.

2

Consider the statement $\forall p \left(\forall q, \frac{1}{p^2} < \frac{1}{q^2} \right)$.

Either prove that the statement is true or provide a counter-example.

End of Question 12

Question 13 (15 marks) Use a SEPARATE writing booklet

(a) (i) Express
$$\frac{-x^2 + 2x + 5}{(x^2 + 2)(1 - x)}$$
 in the form $\frac{ax + b}{x^2 + 2} + \frac{c}{1 - x}$.

- (ii) Hence find $\int \frac{-x^2 + 2x + 5}{(x^2 + 2)(1 x)} dx$.
- (b) A particle of mass M kilograms is projected vertically upward with a velocity of 120 ms⁻¹. The air resistance acting on the particle is 3Mν newtons, where ν is the velocity of the particle.
 - (i) Show that if the acceleration due to gravity is 10 ms⁻², the equation of motion is given by $\ddot{x} = -(10 + 3v)$.
 - (ii) Find the maximum height reached by the particle, correct to the nearest metre. 3
 - (iii) Find the time at which the particle reaches its maximum height, correct to one decimal place.
- (c) A sphere S_1 with centre $\underline{c} = 2\underline{i} + 2\underline{j} + 2\underline{k}$ passes through $\underline{a} = 4\underline{i} + 4\underline{j} + 4\underline{k}$
 - (i) Find the Cartesian equation of S_1 .
 - (ii) A second sphere, S_2 , has equation $(x-2)^2 + (y-2)^2 + (z-5)^2 = 1$. Find the equation of the circle in which S_1 and S_2 intersect and state the centre and radius of this circle.

End of Question 13

Question 14 (15 marks) Use a SEPARATE writing booklet.

(a) Prove that
$$\frac{1+\sqrt{5}}{2}$$
 is irrational. 3

- (b) The price, p, of fuel rises and falls in simple harmonic motion according to the equation $p = 1.5 + \frac{1}{2} \sin \frac{\pi}{7} t + \frac{1}{2} \cos \frac{\pi}{7} t$, where the price is measured in dollars and t is the number of days after midnight Sunday.
 - (i) What is the amplitude and period of the fuel price?
 - (ii) What is the price of fuel, to the nearest cent, at 12 midday on Monday?
 - (iii) At what time and day, correct to the nearest hour, will the fuel price first be at a minimum?
- (c) A particle is moving such that $\ddot{x} = 2x^3 + 6x^2 + 4x$. Initially x = 1 and y = -3.
 - (i) Show that v = -x(x+2).
 - (ii) Find an expression for x in terms of t.
 - (iii) Hence, find the limiting position of the particle.

End of Question 14

3

3

Question 15 (15 marks) Use a SEPARATE writing booklet.

(a) Find $\int e^{-x} \sin(-x) dx$.

3

(b) A projectile is fired from ground level with an initial velocity u ms⁻¹ at an angle of θ to the horizontal. The air resistance is directly proportional to the velocity, with k the constant of proportionality.

Assume that the equations of motion are:

$$x = \frac{u\cos\theta}{k} \left(1 - e^{-kt}\right)$$

$$y = \frac{10 + ku\sin\theta}{k^2} \left(1 - e^{-kt} \right) - \frac{10t}{k}$$

where (x, y) are the coordinates of the projectile at time t seconds. Do NOT prove these equations.

The projectile is fired at an angle of 60°, with initial velocity $10\sqrt{3}$ ms⁻¹ and k = 0.4.

(i) Find the time when the projectile reaches its greatest height.

2

(ii) The projectile hits the ground when t≈ 2.6 seconds. Find the magnitude and direction of the velocity of the projectile when it hits the ground.

3

- (c) Consider the equation $z^5 = 1$.
 - (i) Write down, in polar form, the five roots of $z^5 = 1$.

2

(ii) Show that, for $z \neq 1$,

2

$$\frac{z^5 - 1}{z - 1} = \left(z^2 - 2z\cos\left(\frac{2\pi}{5}\right) + 1\right)\left(z^2 - 2z\cos\left(\frac{4\pi}{5}\right) + 1\right).$$

(iii) Deduce that $\cos\left(\frac{2\pi}{5}\right)$ and $\cos\left(\frac{4\pi}{5}\right)$ are roots of the equation $4x^2 + 2x - 1 = 0$.

End of Question 15

Question 16 (15 marks) Use a SEPARATE writing booklet.

- (a) A sequence is defined recursively as u₁ = 0, u₂ = 25 and u₂ = 10u₂ − 25u₂ , for n≥3.
 Using mathematical induction, prove that u₂ = (n-1)5ⁿ for n≥1.
- (b) Let $I_n = \int_0^{\frac{\pi}{2}} \cos^n x \sin^2 x \, dx$.

(i) Show that
$$I_n = \left(\frac{n-1}{n+2}\right)I_{n-2}$$
 for $n \ge 2$.

(ii) Hence show that
$$I_2 = \frac{\pi}{16}$$
.

(c) (i) Show that
$$0 < \frac{1}{x\sqrt{x}} < \frac{1}{x} < \frac{1}{\sqrt{x}}$$
 for $x > 1$.

(ii) Show that
$$0 < \frac{1}{\sqrt{x}} < \frac{1}{x} < \frac{1}{x\sqrt{x}}$$
 for $0 < x < 1$.

(iii) Show that, for
$$t \ge 1$$
, $\frac{2(\sqrt{t}-1)}{\sqrt{t}} \le \ln t \le 2(\sqrt{t}-1)$.

(iv) Find
$$\lim_{t\to 0} (t \ln t)$$
 and $\lim_{t\to \infty} \left(\frac{\ln t}{t}\right)$

End of paper