



Student Number: _____

Parramatta Marist High School

2020

YEAR 12 TRIAL 1

Mathematics Extension 2

General Instructions

- Reading time – 10 minutes
- Working time – 3 hours
- Write using black pen
- Calculators approved by NESA may be used
- A reference sheet is provided at the back of this paper
- Show relevant mathematical reasoning and/or calculations

Total Marks: 100

Section I – 10 marks (pages 2–4)

- Attempt Questions 1–10
- Allow about 15 minutes for this section

Section II – 90 marks (page 5–19)

- Attempt Questions 11–23
 - Allow about 2 hours and 45 minutes for this section
-

Section I

10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1–10.

- 1 What is the contrapositive of $P \implies \neg Q$?
- A. $Q \implies P$
 - B. $\neg Q \implies P$
 - C. $Q \implies \neg P$
 - D. $\neg Q \implies \neg P$
- 2 Let $z = 2 - 7i$ and $w = 5 + 3i$.
What is the value of $\bar{z} - 2w$?
- A. $-8 - 13i$
 - B. $-8 + i$
 - C. $12 - i$
 - D. $12 + 13i$
- 3 What is the Cartesian form of $z = i \sec \theta + j \tan \theta$?
- A. $x^2 - y^2 = 1$
 - B. $x^2 + y^2 = 1$
 - C. $y^2 - x^2 = 1$
 - D. $x^2 - y^2 = -1$
- 4 What are the roots of the polynomial $P(x) = x^3 + 3x^2 + 4x + 2$?
- A. $1, 1 + i, 1 - i$
 - B. $-1, 1 + i, 1 - i$
 - C. $1, -1 + i, -1 - i$
 - D. $-1, -1 + i, -1 - i$

- 5 What is the negation of $\exists x \in \mathbb{Z} : x^2 = -1$?
- $\exists x \notin \mathbb{Z} : x^2 = -1$
 - $\exists x \in \mathbb{Z} : x^2 \neq -1$
 - $\forall x \in \mathbb{Z} : x^2 = -1$
 - $\forall x \in \mathbb{Z} : x^2 \neq -1$
- 6 What is the angle between the vectors $u = \underline{i} + \underline{k}$ and $v = \underline{j} - \underline{k}$?
- $\frac{\pi}{3}$
 - $\frac{\pi}{2}$
 - $\frac{2\pi}{3}$
 - π
- 7 Let $z = 1 - i$.
What is z^3 in exponential form?
- $e^{\frac{3\pi i}{4}}$
 - $e^{-\frac{3\pi i}{4}}$
 - $2^{\frac{3}{2}} e^{\frac{3\pi i}{4}}$
 - $2^{\frac{3}{2}} e^{-\frac{3\pi i}{4}}$
- 8 The point $(0, 1, -1)$ lies on which line?
- $\underline{r} = \underline{i} + \underline{j} + \lambda(\underline{i} + \underline{k})$
 - $\underline{r} = \underline{i} + \underline{j} + \lambda(\underline{j} + \underline{k})$
 - $\underline{r} = \underline{i} + \underline{k} + \lambda(\underline{i} + \underline{j})$
 - $\underline{r} = \underline{i} + \underline{k} + \lambda(\underline{j} + \underline{k})$

- 9 The probability function $f(x) = \begin{cases} \pi x \sin \pi x, & x \in [0, 1] \\ 0, & x \notin [0, 1] \end{cases}$.

What is $P(x \leq \frac{1}{2})$?

- A. $\frac{1}{\pi^2}$
- B. $\frac{1}{\pi}$
- C. $\frac{1}{2}$
- D. $\frac{\pi}{2}$

- 10 A particle, initially at rest at the origin, moves with equation of motion $a = 1 + v^2$.

What is the equation of motion for v in terms of x ?

- A. $v = \tan x$
- B. $v = e^x - 1$
- C. $v = x + \frac{1}{3}x^3$
- D. $v = \sqrt{e^{2x} - 1}$

Section II

90 marks

Attempt Questions 11–23

Allow about 2 hours and 45 minutes for this section

Answer the questions in the spaces provided. Sufficient spaces are provided for typical responses.

Your responses should include relevant mathematical reasoning and/or calculations.

Extra writing space is provided at the back of this booklet. If you use this space, clearly indicate which question you are answering.

Question 11 (3 marks)

Prove that $2^{n+1} + 3^{2n-1}$ is divisible by 7 for $n \in \mathbb{Z}^+$. **3**

.....

.....

.....

.....

.....

.....

.....

.....

Question 12 (4 marks)

- (a) Express $1 + i\sqrt{3}$ in exponential form.

2

.....

.....

.....

.....

- (b) Hence find the two values of $\sqrt{1 + i\sqrt{3}}$ in Cartesian form.

2

.....


.....

.....

.....

Question 13 (4 marks)

In the box below, shade the region in the complex plane that simultaneously satisfies $|z| < |z - 2 - 2i|$ and $\frac{\pi}{12} \leq \arg z \leq \frac{5\pi}{12}$.



Question 14 (7 marks)

- (a) Let $u_1 = 1$, $u_n = u_{n-1} + n$ for $n \geq 2$. **3**

Prove that $u_n = \frac{1}{2}n(n+1)$ for $n \in \mathbb{Z}^+$.

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence, prove that $\sum_{k=0}^n k^3 = u_n^2$ for $n \in \mathbb{Z}^+$. **4**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Question 15 (15 marks)

(a) Find $\int_0^{\frac{\pi}{2}} \frac{dx}{1 + \cos x}$. **3**

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find $\int \frac{dx}{\sqrt{x^2 + 2x + 2}}$. **4**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Question 15 continues on page 9

Question 15 (continued)

(c) Find $\int_0^{\frac{\pi}{4}} \tan^3 x \sec^4 x \, dx$.

4

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Find $\int \frac{4 \, dx}{(x^2 + 1)(x - 1)}$.

4

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

End of Question 15

Question 16 (5 marks)

The displacement x at time t of a particle moving on the x -axis is given by

$$x = 3 + \sqrt{3} \sin 3t + \cos 3t.$$

- (a) Show that the motion of the particle is simple harmonic.

2

.....

.....

.....

.....

.....

.....

- (b) Find the amplitude and phase of the motion.

3

.....

.....

.....

.....

.....

.....


.....

.....

Question 17 (4 marks)

In the diagram below, B is the midpoint of AD and C is the midpoint of AE .

4



Using vectors, prove that \overrightarrow{BC} is half the magnitude of, and parallel to, \overrightarrow{DE} .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Questions 11–17 are worth 42 marks in total.

Question 18 (4 marks)

- (a) Prove that $\frac{x+y}{2} \geq \sqrt{xy}$ for $x, y \in \mathbb{R}^+$.

2

.....

.....

.....

.....

.....

- (b) Hence prove that $\frac{a}{b} + \frac{b}{a} \geq 2$ for $a, b \in \mathbb{R}^+$.

2

.....

.....

.....

.....

.....

Question 19 (8 marks)

- (a) Consider the sphere given by the Cartesian equation $x^2 + y^2 + z^2 + 2x - 4z - 4 = 0$. **2**

Show that the vector equation of the sphere is $\left| \underline{r} - \begin{bmatrix} -1 \\ 0 \\ 2 \end{bmatrix} \right| = 3$.

.....

.....

.....

.....

.....

- (b) Find the points of intersection between the sphere and the line $\underline{r} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$. **3**

.....

.....

.....

.....

.....

.....

.....

- (c) Show that the line $\underline{r} = -\underline{i} + \underline{j} - \underline{k} + \mu \underline{j}$ is tangent to the sphere. **3**

.....

.....

.....

.....

.....

.....

.....

Question 20 (7 marks)

(a) Let $I_n = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^n x \, dx$.

3

Show that for $n \geq 2$, $I_n = \frac{n-1}{n} I_{n-2}$.

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence find the volume of revolution for $y = \cos^3 x$ about the x -axis for $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

4

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Question 21 (7 marks)

- (a) Using De Moivre, show that $\cos 5x = 16\cos^5 x - 20\cos^3 x + 5\cos x$.

3

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence, show that $\cos 18^\circ = \frac{\sqrt{10+2\sqrt{5}}}{4}$.

4

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Question 22 (10 marks)

A projectile is launched from the ground with an initial velocity of u m/s at an angle of θ to the horizontal. The projectile experiences the effect of gravity, and a resistance proportional to its velocity in both the horizontal and vertical directions.

The equations of motion are given by $\underline{a} = -0.1\underline{v} - 10\underline{j}$, where \underline{a} is the acceleration vector.

- (a) Show that the velocity vector $\underline{v} = e^{-0.1t} (\underline{u} + \underline{j}) - \underline{j}$, where $\underline{u} = u \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$, satisfies the equations of motion and initial conditions. **3**

.....

.....

.....

.....

.....

.....

- (b) Show that the projectile reaches its peak (maximum height) after $10 \ln(u \sin \theta + 1)$ seconds. **3**

.....

.....

.....

.....

.....

.....

.....

.....

Question 22 continues on page 17

Question 22 (continued)

- (c) Let the speed of the projectile at its peak be w m/s.
Show that $\lim_{\mu \rightarrow \infty} w = \cot \theta$.

4

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

End of Question 22

Question 23 (12 marks)

Recall that $x \in \mathbb{Q} \iff \exists p \in \mathbb{Z}, q \in \mathbb{Z}^+ : x = \frac{p}{q}$.

- (a) Show that $\forall r \in \mathbb{Z}^+ \exists n \in \mathbb{Z}^+ : 0 < \int_0^1 x^n e^x \, dx < \frac{1}{r}$. **3**

.....

.....

.....

.....

.....

.....

.....

- (b) Prove, using induction, that for all integers $n \geq 0$, $\exists \alpha, \beta \in \mathbb{Z} : \int_0^1 x^n e^x \, dx = \alpha + \beta e$. **4**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Question 23 continues on page 19

Question 23 (continued)

- (c) Let $p \in \mathbb{Z}, q \in \mathbb{Z}^+, x = \frac{p}{q} \in \mathbb{Q}$.

Show that $\forall \alpha, \beta \in \mathbb{Z} : \left\{ \alpha + \beta x \neq 0 \implies |\alpha + \beta x| \geq \frac{1}{q} \right\}$.

2

.....

.....

.....

.....

.....

.....

- (d) Hence, prove that $e \notin \mathbb{Q}$.

3

.....

.....

.....

.....

.....

.....

.....

.....

End of Question 23

End of Paper

This image shows a full page of primary-ruled paper. It features a vertical margin line on the left side, creating a narrow left margin. The rest of the page is filled with horizontal dotted lines, which are evenly spaced and extend from the right edge to the margin line. There are no pre-written words or numbers on the page.

[illegible]

[illegible]



Mathematics Advanced

Mathematics Extension 1

Mathematics Extension 2

REFERENCE SHEET

Measurement

Length

$$l = \frac{\theta}{360} \times 2\pi r$$

Area

$$A = \frac{\theta}{360} \times \pi r^2$$

$$A = \frac{h}{2}(a + b)$$

Surface area

$$A = 2\pi r^2 + 2\pi rh$$

$$A = 4\pi r^2$$

Volume

$$V = \frac{1}{3}Ah$$

$$V = \frac{4}{3}\pi r^3$$

Functions

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\text{For } ax^3 + bx^2 + cx + d = 0:$$

$$\alpha + \beta + \gamma = -\frac{b}{a}$$

$$\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a}$$

$$\text{and } \alpha\beta\gamma = -\frac{d}{a}$$

Relations

$$(x - h)^2 + (y - k)^2 = r^2$$

Financial Mathematics

$$A = P(1 + r)^n$$

Sequences and series

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d] = \frac{n}{2}(a + l)$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} = \frac{a(r^n - 1)}{r - 1}, r \neq 1$$

$$S = \frac{a}{1 - r}, |r| < 1$$

Logarithmic and Exponential Functions

$$\log_a a^x = x = a^{\log_a x}$$

$$\log_a x = \frac{\log_b x}{\log_b a}$$

$$a^x = e^{x \ln a}$$

Trigonometric Functions

$$\sin A = \frac{\text{opp}}{\text{hyp}}, \quad \cos A = \frac{\text{adj}}{\text{hyp}}, \quad \tan A = \frac{\text{opp}}{\text{adj}}$$

$$A = \frac{1}{2} ab \sin C$$



$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$l = r\theta$$

$$A = \frac{1}{2} r^2 \theta$$



Trigonometric identities

$$\sec A = \frac{1}{\cos A}, \quad \cos A \neq 0$$

$$\operatorname{cosec} A = \frac{1}{\sin A}, \quad \sin A \neq 0$$

$$\cot A = \frac{\cos A}{\sin A}, \quad \sin A \neq 0$$

$$\cos^2 x + \sin^2 x = 1$$

Compound angles

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\text{If } t = \tan \frac{A}{2} \text{ then } \sin A = \frac{2t}{1+t^2}$$

$$\cos A = \frac{1-t^2}{1+t^2}$$

$$\tan A = \frac{2t}{1-t^2}$$

$$\cos A \cos B = \frac{1}{2} [\cos(A - B) + \cos(A + B)]$$

$$\sin A \sin B = \frac{1}{2} [\cos(A - B) - \cos(A + B)]$$

$$\sin A \cos B = \frac{1}{2} [\sin(A + B) + \sin(A - B)]$$

$$\cos A \sin B = \frac{1}{2} [\sin(A + B) - \sin(A - B)]$$

$$\sin^2 nx = \frac{1}{2} (1 - \cos 2nx)$$


$$\cos^2 nx = \frac{1}{2} (1 + \cos 2nx)$$

Statistical Analysis

$$z = \frac{x - \mu}{\sigma}$$

An outlier is a score
less than $Q_1 - 1.5 \times IQR$
or
more than $Q_3 + 1.5 \times IQR$

Normal distribution



- approximately 68% of scores have z -scores between -1 and 1
- approximately 95% of scores have z -scores between -2 and 2
- approximately 99.7% of scores have z -scores between -3 and 3

$$E(X) = \mu$$

$$\operatorname{Var}(X) = E[(X - \mu)^2] = E(X^2) - \mu^2$$

Probability

$$P(A \cap B) = P(A)P(B)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad P(B) \neq 0$$

Continuous random variables

$$P(X \leq r) = \int_a^r f(x) dx$$

$$P(a < X < b) = \int_a^b f(x) dx$$

Binomial distribution

$$P(X = r) = {}^nC_r p^r (1-p)^{n-r}$$

$$X \sim \operatorname{Bin}(n, p)$$

$$\Rightarrow P(X = x)$$

$$= \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$$

$$E(X) = np$$

$$\operatorname{Var}(X) = np(1-p)$$

Differential Calculus

Function

Derivative

$$y = f(x)^n$$

$$\frac{dy}{dx} = n f'(x) [f(x)]^{n-1}$$

$$y = uv$$

$$\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$y = g(u) \text{ where } u = f(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$y = \frac{u}{v}$$

$$\frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$y = \sin f(x)$$

$$\frac{dy}{dx} = f'(x) \cos f(x)$$

$$y = \cos f(x)$$

$$\frac{dy}{dx} = -f'(x) \sin f(x)$$

$$y = \tan f(x)$$

$$\frac{dy}{dx} = f'(x) \sec^2 f(x)$$

$$y = e^{f(x)}$$

$$\frac{dy}{dx} = f'(x) e^{f(x)}$$

$$y = \ln f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{f(x)}$$

$$y = a^{f(x)}$$

$$\frac{dy}{dx} = (\ln a) f'(x) a^{f(x)}$$

$$y = \log_a f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{(\ln a) f(x)}$$

$$y = \sin^{-1} f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{\sqrt{1 - [f(x)]^2}}$$

$$y = \cos^{-1} f(x)$$

$$\frac{dy}{dx} = -\frac{f'(x)}{\sqrt{1 - [f(x)]^2}}$$

$$y = \tan^{-1} f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{1 + [f(x)]^2}$$

Integral Calculus

$$\int f'(x) [f(x)]^n dx = \frac{1}{n+1} [f(x)]^{n+1} + c$$

where $n \neq -1$

$$\int f'(x) \sin f(x) dx = -\cos f(x) + c$$

$$\int f'(x) \cos f(x) dx = \sin f(x) + c$$

$$\int f'(x) \sec^2 f(x) dx = \tan f(x) + c$$

$$\int f'(x) e^{f(x)} dx = e^{f(x)} + c$$

$$\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c$$

$$\int f'(x) a^{f(x)} dx = \frac{a^{f(x)}}{\ln a} + c$$

$$\int \frac{f'(x)}{\sqrt{a^2 - [f(x)]^2}} dx = \sin^{-1} \frac{f(x)}{a} + c$$

$$\int \frac{f'(x)}{a^2 + [f(x)]^2} dx = \frac{1}{a} \tan^{-1} \frac{f(x)}{a} + c$$

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

$$\int_a^b f(x) dx$$

$$\approx \frac{b-a}{2n} \left\{ f(a) + f(b) + 2[f(x_1) + \dots + f(x_{n-1})] \right\}$$

where $a = x_0$ and $b = x_n$

Combinatorics

$${}^nP_r = \frac{n!}{(n-r)!}$$

$$\binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!}$$

$$(x+a)^n = x^n + \binom{n}{1}x^{n-1}a + \cdots + \binom{n}{r}x^{n-r}a^r + \cdots + a^n$$

Vectors

$$|\underline{u}| = |x_1\underline{i} + y_1\underline{j}| = \sqrt{x^2 + y^2}$$

$$\underline{u} \cdot \underline{v} = |\underline{u}| |\underline{v}| \cos \theta = x_1x_2 + y_1y_2,$$

$$\text{where } \underline{u} = x_1\underline{i} + y_1\underline{j}$$

$$\text{and } \underline{v} = x_2\underline{i} + y_2\underline{j}$$

$$\underline{r} = \underline{a} + \lambda \underline{b}$$

Complex Numbers

$$\begin{aligned} z = a + ib &= r(\cos \theta + i \sin \theta) \\ &= re^{i\theta} \end{aligned}$$

$$\begin{aligned} [r(\cos \theta + i \sin \theta)]^n &= r^n(\cos n\theta + i \sin n\theta) \\ &= r^ne^{in\theta} \end{aligned}$$

Mechanics

$$\frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx} \left(\frac{1}{2} v^2 \right)$$

$$x = a \cos(nt + \alpha) + c$$

$$x = a \sin(nt + \alpha) + c$$

$$\ddot{x} = -n^2(x - c)$$



Student Number: _____

Parramatta Marist High School

2020

YEAR 12 TRIAL 1

Mathematics Extension 2

General Instructions

- Reading time – 10 minutes
- Working time – 3 hours
- Write using black pen
- Calculators approved by NESA may be used
- A reference sheet is provided at the back of this paper
- Show relevant mathematical reasoning and/or calculations

**Total Marks:
100**

Section I – 10 marks (pages 2–4)

- Attempt Questions 1–10
- Allow about 15 minutes for this section

Section II – 90 marks (page 5–19)

- Attempt Questions 11–23
- Allow about 2 hours and 45 minutes for this section

Section I

10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1–10.

-
- 1 What is the contrapositive of $P \Rightarrow \neg Q$? $(A \Rightarrow B) \Leftrightarrow (\neg B \Rightarrow \neg A)$
- A. $Q \Rightarrow P$
B. $\neg Q \Rightarrow P$
☒ C. $Q \Rightarrow \neg P$
D. $\neg Q \Rightarrow \neg P$
- 2 Let $z = 2 - 7i$ and $w = 5 + 3i$. $z - 2w = 2 - 7i - 2(5 + 3i)$
 $= 2 - 10 - 6i = -8 - 6i$
 $\therefore B$
- What is the value of $\bar{z} - 2w$?
- A. $-8 - 13i$
☒ B. $-8 + i$
C. $12 - i$
D. $12 + 13i$
- 3 What is the Cartesian form of $r = i \sec \theta + j \tan \theta$? $1 + \tan^2 \theta = \sec^2 \theta$
 $\sec^2 \theta - \tan^2 \theta = 1$
- ☒ A. $x^2 - y^2 = 1$
B. $x^2 + y^2 = 1$
C. $y^2 - x^2 = 1$
D. $x^2 - y^2 = -1$
- 4 What are the roots of the polynomial $P(x) = x^3 + 3x^2 + 4x + 2$?
- A. $1, 1 + i, 1 - i$ $\text{sum} = -3, \therefore D.$
B. $-1, 1 + i, 1 - i$
C. $1, -1 + i, -1 - i$
☒ D. $-1, -1 + i, -1 - i$

- 5 What is the negation of $\exists x \in \mathbb{Z} : x^2 = -1$?

A. $\exists x \notin \mathbb{Z} : x^2 = -1$

B. $\exists x \in \mathbb{Z} : x^2 \neq -1$

C. $\forall x \in \mathbb{Z} : x^2 = -1$

☒ D. $\forall x \in \mathbb{Z} : x^2 \neq -1$

It's not the case that there exists $x \in \mathbb{Z}$ such that $x^2 = -1$.
 \therefore for any $x \in \mathbb{Z}$, $x^2 \neq -1$.


- 6 What is the angle between the vectors $u = \underline{i} + \underline{k}$ and $v = \underline{j} - \underline{k}$?

A. $\frac{\pi}{3}$

B. $\frac{\pi}{2}$

☒ C. $\frac{2\pi}{3}$

D. π



- 7 Let $z = 1 - i$.

What is z^3 in exponential form?

A. $e^{\frac{3\pi i}{4}}$

B. $e^{-\frac{3\pi i}{4}}$

C. $2^{\frac{3}{2}} e^{\frac{3\pi i}{4}}$

☒ D. $2^{\frac{3}{2}} e^{-\frac{3\pi i}{4}}$

$|z| = \sqrt{2}$
 $\text{Arg}(z^3) < 0$
 $\therefore D$

- 8 The point $(0, 1, -1)$ lies on which line?

☒ A. $r = \underline{i} + \underline{j} + \lambda(\underline{i} + \underline{k})$

B. $r = \underline{i} + \underline{j} + \lambda(\underline{j} + \underline{k})$

C. $r = \underline{i} + \underline{k} + \lambda(\underline{i} + \underline{j})$

D. $r = \underline{i} + \underline{k} + \lambda(\underline{j} + \underline{k})$

$x = 0 \Rightarrow A \text{ or } C$, with $\lambda = -1$.
 $y = 1$ when $\lambda = -1 \Rightarrow A$

- 9 The probability function $f(x) = \begin{cases} \pi x \sin \pi x, & x \in [0, 1] \\ 0, & x \notin [0, 1] \end{cases}$. What is $P(x \leq \frac{1}{2})$?
- A. $\frac{1}{\pi^2}$
 B. $\frac{1}{\pi}$
 C. $\frac{1}{2}$
 D. $\frac{\pi}{2}$
- oof!
- $$\int_0^{1/2} \pi x \sin \pi x \, dx$$
- $$= \left[-\frac{1}{\pi} x \cos \pi x \right]_0^{1/2} + \int_0^{1/2} \frac{1}{\pi} \cos \pi x \, dx$$
- $$= \left[-\frac{1}{2} \cos\left(\frac{\pi}{2}\right) - 0 \right] + \left[\frac{\sin \pi x}{\pi} \right]_0^{1/2}$$
- $$= 0 + \frac{\sin(\pi/2)}{\pi} - \frac{\sin(0)}{\pi}$$
- $$= \frac{1}{\pi}$$

- 10 A particle, initially at rest at the origin, moves with equation of motion $a = 1 + v^2$. What is the equation of motion for v in terms of x ?

- A. $v = \tan x$
 B. $v = e^x - 1$
 C. $v = x + \frac{1}{3}x^3$
 D. $v = \sqrt{e^{2x} - 1}$

As this is M.C.,
 looking at $\int \frac{v \, dv}{1+v^2}$
 should indicate the
 solution is the square root
 of an exponential,
 hence D.

$$v \frac{dv}{dx} = 1 + v^2$$

$$\frac{1}{2} \int \frac{2v \, dv}{1+v^2} = \int dx$$

$$\frac{1}{2} \ln(1+v^2) = x + C$$

when $t=0$, $x=0$, $v=0$:

$$\frac{1}{2} \ln(1+0^2) = 0 + C$$

$$\therefore C=0$$

$$\ln(1+v^2) = 2x$$

$$1+v^2 = e^{2x}$$

$$v^2 = e^{2x} - 1 \Rightarrow v = \sqrt{e^{2x} - 1}$$

Section II

90 marks

Attempt Questions 11–23

Allow about 2 hours and 45 minutes for this section

Answer the questions in the spaces provided. Sufficient spaces are provided for typical responses.

Your responses should include relevant mathematical reasoning and/or calculations.

Extra writing space is provided at the back of this booklet. If you use this space, clearly indicate which question you are answering.

Question 11 (3 marks)

Prove that $2^{n+1} + 3^{2n-1}$ is divisible by 7 for $n \in \mathbb{Z}^+$.

3

Base case ($n=1$): $2^{1+1} + 3^{2 \cdot 1 - 1} = 4 + 3$
 $= 7$ \therefore true for $n=1$.

Assume $\exists k \in \mathbb{Z}^+ : \exists p \in \mathbb{Z} : 2^{k+1} + 3^{2k-1} = 7p \Rightarrow 2^{k+1} = 7p - 3^{2k-1}$

Consider: $2^{k+2} + 3^{2k+1} = 2(2^{k+1}) + 3^{2k+1}$
 $= 2(7p - 3^{2k-1}) + 3^{2k+1}$
 $= 7(2p) + 3^{2k-1}(9 - 2)$
 $= 7(2p + 3^{2k-1})$
 $= 7Q$, for $Q = 2p + 3^{2k-1}$,
and Q clearly in \mathbb{Z} .

\therefore true by mathematical induction.

Question 12 (4 marks)

- (a) Express $1 + i\sqrt{3}$ in exponential form.

2

$\text{modulus} = 2$, $\text{argument} = \frac{\pi}{3}$
 $\therefore 1 + i\sqrt{3} = 2e^{i\pi/3}$

- (b) Hence find the two values of $\sqrt{1 + i\sqrt{3}}$ in Cartesian form.

2


roots are: $\sqrt[2]{2} e^{i\frac{\pi}{6}}$ or $-\sqrt[2]{2} e^{i\frac{\pi}{6}}$
 $= \sqrt{2} (\sqrt{3}/2 + i/2)$ or $\sqrt{2} (-\sqrt{3}/2 - i/2)$
 $= \pm (\frac{\sqrt{3}}{\sqrt{2}} + \frac{i}{\sqrt{2}})$

Question 13 (4 marks)

$|z - 0| < |z - (2 + 2i)| \rightarrow z$ is closer to 0 than $(2 + 2i)$

In the box below, shade the region in the complex plane that simultaneously satisfies

$|z| < |z - 2 - 2i|$ and $\frac{\pi}{12} \leq \arg z \leq \frac{5\pi}{12}$.



Question 14 (7 marks)

triangle numbers! Yay!

- (a) Let $u_1 = 1$, $u_n = u_{n-1} + n$ for $n \geq 2$.

3

Prove that $u_n = \frac{1}{2}n(n+1)$ for $n \in \mathbb{Z}^+$.

when $n=1$, $u_1 = \frac{1}{2} \times 1(1+1)$
 $= 1$ ✓

\therefore true for $n=1$

Assume $\exists k \in \mathbb{Z}^+ : u_k = \frac{k(k+1)}{2}$

consider $u_{k+1} = \frac{1}{2}k(k+1) + (k+1)$
 $= \frac{1}{2}k(k+1) + \frac{2}{2}(k+1)$
 $= \frac{1}{2}(k+1)[k+2]$
 $= \frac{1}{2}(k+1)(k+2)$

\therefore true by mathematical induction.

- (b) Hence, prove that $\sum_{k=0}^n k^3 = u_n^2$ for $n \in \mathbb{Z}^+$.

sum of cubes Yay!

4

Base case, $n=1$: $\sum_{k=0}^1 k^3 = 0^3 + 1^3$, $u_1^2 = 1^2$
 $= 1$ $= 1$ \therefore true ✓

Assume $\exists j \in \mathbb{Z}^+ : \sum_{k=0}^j k^3 = u_j^2$

consider $\sum_{k=0}^{j+1} k^3 = \sum_{k=0}^j k^3 + (j+1)^3$

$= \frac{1}{4}j^2(j+1)^2 + (j+1)^3 \times \frac{4}{4}$
 $= (j+1)^2[j^2 + 4j + 4] \quad 3 \times \frac{1}{4}$
 $= \frac{1}{4}(j+1)^2(j+2)^2$
 $= u_{j+1}^2$

\therefore true by mathematical induction.

Question 15 (15 marks)

Very tricky for 3 marks!

(a) Find $\int_0^{\pi/2} \frac{dx}{1 + \cos x} = I$

3

$$I = \int_0^{\pi/2} \frac{1 - \cos x}{\sin^2 x} dx$$

$$= \int_0^{\pi/2} \cos x^2 dx - \int_0^{\pi/2} (\sin x)^{-2} \cos x dx$$

$$= \left[-\sin x \right]_0^{\pi/2} - \left[\frac{1}{-1} (\sin x)^{-1} \right]_0^{\pi/2}$$

$$= \left[\frac{-\sin x}{\sin x} + 1 \right]_0^{\pi/2} \times \frac{(1 + \cos x)}{(1 + \cos x)}$$

$$= \left[\frac{1 - \cos x}{1 + \cos x} \right]_0^{\pi/2}$$

$$= \left[\frac{1 - \cos x}{1 + \cos x} \right]_0^{\pi/2}$$

$$= 1 - 0 = 1$$

(b) Find $\int \frac{dx}{\sqrt{x^2 + 2x + 2}}$

4

$$\int \frac{dx}{\sqrt{(x+1)^2 + 1}} = \ln(x+1 + \sqrt{(x+1)^2 + 1}) + C$$

$$= \ln(x^2 + 2x + 1 + \sqrt{x^2 + 2x + 2}) + C$$

What was the intuition here?

prove a standard integral?

Question 15 continues on page 9

Question 15 (continued)

$$1 + \tan^2 x = \sec^2 x$$

- (c) Find $\int_0^{\pi/4} \tan^3 x \sec^4 x \, dx$.

4

$$\begin{aligned} &= \int_0^{\pi/4} \sec^2 x [\tan^5 x + \tan^3 x] \, dx \\ &= \int_0^{\pi/4} \tan^5 x \sec^2 x \, dx + \int_0^{\pi/4} \tan^3 x \sec^2 x \, dx \\ &= \left[\frac{1}{6} \tan^6 x \right]_0^{\pi/4} + \left[\frac{1}{4} \tan^4 x \right]_0^{\pi/4} \\ &= \frac{1}{6} + \frac{1}{4} \\ &= \frac{5}{12} \end{aligned}$$

- (d) Find $\int \frac{4 \, dx}{(x^2+1)(x-1)}$.

4

$$\begin{aligned} &4 \equiv (Ax+B)(x-1) + (C)(x^2+1) \\ x=1 &\Rightarrow 4=2C \Rightarrow C=2 \\ x=i &\Rightarrow 4=-A-B, \quad 0=-A+B \Rightarrow B=A \\ &4=-2A \\ &A=-2=B \\ \therefore I &= -2 \int \frac{dx}{(x^2+1)} + 2 \int \frac{dx}{x-1} \\ &= -2 \tan^{-1} x + 2 \ln|x-1| + C \end{aligned}$$

End of Question 15

Question 16 (5 marks)

The displacement x at time t of a particle moving on the x -axis is given by

$$x = 3 + \sqrt{3} \sin 3t + \cos 3t.$$

- (a) Show that the motion of the particle is simple harmonic.

2

$$\begin{aligned} \ddot{x} &= 3\sqrt{3} \cos 3t - 3 \sin 3t \\ \ddot{x} &= -3^2 \sqrt{3} \sin 3t - 3^2 \cos 3t \\ &= -9 [\sqrt{3} \sin 3t + \cos 3t] \\ &= -9 [x - 3] \\ &\therefore \text{SHM} \end{aligned}$$

- (b) Find the amplitude and phase of the motion.




3

$$\begin{aligned} x &= 3 + 2 \left(\frac{\sqrt{3}}{2} \sin 3t + \frac{1}{2} \cos 3t \right) \\ &= 3 + 2 \left(\sin \frac{\pi}{4} \sin 3t + \cos \frac{\pi}{4} \cos 3t \right) \\ &= 3 + 2 \cos \left(3t - \frac{\pi}{4} \right) \\ &= 3 + 2 \cos \left(3 \left(t - \frac{\pi}{12} \right) \right) \\ \therefore \text{amplitude} &= 2, \quad \text{phase} = \frac{\pi}{12} \end{aligned}$$

Question 17 (4 marks)

In the diagram below, B is the midpoint of AD and C is the midpoint of AE .

4



Yay!
vectors!

Using vectors, prove that \vec{BC} is half the magnitude of, and parallel to, \vec{DE} .

$$\begin{aligned}
 \vec{DE} &= \vec{DA} + \vec{AE} \\
 &= \vec{DB} + \vec{BA} + \vec{AC} + \vec{CE} \\
 &= 2\vec{BA} + 2\vec{AC} \quad (\text{as } \vec{DB} = \vec{BA} \text{ and } \vec{AC} = \vec{CE}) \\
 &= 2(\vec{BC})
 \end{aligned}$$

Questions 11–17 are worth 42 marks in total.

Question 18 (4 marks)

- (a) Prove that $\frac{x+y}{2} \geq \sqrt{xy}$ for $x, y \in \mathbb{R}^+$.

2

$$\begin{aligned} \text{let } x, y \in \mathbb{R}^+ \quad (\sqrt{x} - \sqrt{y})^2 &\geq 0 \\ x + y - 2\sqrt{xy} &\geq 0 \\ x + y &\geq 2\sqrt{xy} \\ \frac{x+y}{2} &\geq \sqrt{xy} \end{aligned}$$

- (b) Hence prove that $\frac{a}{b} + \frac{b}{a} \geq 2$ for $a, b \in \mathbb{R}^+$.

2

$$\begin{aligned} \text{hence: let } x = \frac{a}{b}, y = \frac{b}{a} \quad & \text{otherwise: let } a, b \in \mathbb{R}^+ \\ \text{Then } \frac{x+y}{2} &\geq \sqrt{xy} \\ \Rightarrow \frac{\frac{a}{b} + \frac{b}{a}}{2} &\geq 2\sqrt{\frac{a}{b} \cdot \frac{b}{a}} \\ \frac{a}{b} + \frac{b}{a} &\geq 2 \end{aligned} \quad \left| \quad \begin{aligned} (\sqrt{\frac{a}{b}} - \sqrt{\frac{b}{a}})^2 &\geq 0 \\ \frac{a}{b} + \frac{b}{a} - 2 &\geq 0 \\ \frac{a}{b} + \frac{b}{a} &\geq 2 \end{aligned} \right.$$

Question 19 (8 marks)

- (a) Consider the sphere given by the Cartesian equation $x^2 + y^2 + z^2 + 2x - 4z - 4 = 0$. 2

Show that the vector equation of the sphere is $\left| r - \begin{bmatrix} -1 \\ 0 \\ 2 \end{bmatrix} \right| = 3$.

$C_1: (x^2 + 2x + 1) - 1 + y^2 + (z^2 - 4z + 4) - 4 - 4 = 0$
 $(x+1)^2 + y^2 + (z-2)^2 - 9 = 0$
 $\therefore (x+1)^2 + y^2 + (z-2)^2 = 3^2$
 and this is the equation of a sphere
 centered at $\begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix}$ with radius 3.

- (b) Find the points of intersection between the sphere and the line $r = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$. 3

$\left| \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} - \begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix} \right| = 3$
 $\begin{pmatrix} 1 \\ 1+\lambda \\ 1+2\lambda \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1+\lambda \\ 1+2\lambda \end{pmatrix} = 9$
 $(1+\lambda)^2 + (1+\lambda)^2 + (2\lambda-1)^2 = 9$
 $6\lambda^2 + 3 = 9$
 $6\lambda^2 = 6$
 $\lambda = \pm 1$
 $\therefore \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ AND $\begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ -1 \end{pmatrix}$

- (c) Show that the line $r = -i + j - k + \mu j$ is tangent to the sphere. 3

$\begin{pmatrix} -1 \\ 1+\mu \\ -1 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1+\mu \\ -1 \end{pmatrix} = 9$
 $(1+\mu)^2 + (-3)^2 = 9$
 $(1+\mu)^2 = 0$
 has one solution ($\mu=0$)
 \therefore this line is a tangent to the sphere.

Question 20 (7 marks)

(a) Let $I_n = \int_{-\pi/2}^{\pi/2} \cos^n x \, dx$.

3

Show that for $n \geq 2$, $I_n = \frac{n-1}{n} I_{n-2}$.

$$\begin{aligned}
 I_n &= \int_{-\pi/2}^{\pi/2} \cos^n x \, dx \\
 &= \left[\sin x \cos^{n-1} x \right]_{-\pi/2}^{\pi/2} + \int_{-\pi/2}^{\pi/2} \sin x \times (n-1) \sin x \cos^{n-2} x \, dx \\
 &= 0 + \int_{-\pi/2}^{\pi/2} (n-1)(1 - \cos^2 x) \cos^{n-2} x \, dx \\
 &= (n-1) I_{n-2} - (n-1) I_n \\
 \therefore I_n (1 + (n-1)) &= (n-1) I_{n-2} \\
 I_n &= \left(\frac{n-1}{n} \right) I_{n-2}
 \end{aligned}$$

(b) Hence find the volume of revolution for $y = \cos^3 x$ about the x -axis for $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

4

$$\begin{aligned}
 V &= \pi \int_{-\pi/2}^{\pi/2} y^2 \, dx \\
 &= \pi \int_{-\pi/2}^{\pi/2} \cos^6 x \, dx \\
 &= \pi \left[\frac{5}{6} I_4 \right] \\
 &= \pi \left(\frac{5}{6} \right) \left(\frac{3}{4} \right) I_2 \\
 &= \pi \left(\frac{5}{6} \right) \left(\frac{3}{4} \right) \left(\frac{2}{2} \right) \times \int_{-\pi/2}^{\pi/2} dx \\
 &= \pi^2 \times \frac{15}{48} \\
 &= \frac{15\pi^2}{48} \quad u^3
 \end{aligned}$$

Question 21 (7 marks)

- (a) Using De Moivre, show that $\cos 5x = 16\cos^5 x - 20\cos^3 x + 5\cos x$.

3

$$\begin{aligned} (\cos x + i\sin x)^5 &= \cos^5 x + 5i\cos^4 x \sin x - 10\cos^3 x \sin^2 x - 10i\cos^2 x \sin^3 x \\ &\quad + 5\cos x \sin^4 x + i\sin^5 x \end{aligned}$$

Equating real and imaginary:

$$\begin{aligned} \cos 5x &= \cos^5 x - 10\cos^3 x \sin^2 x + 5\cos x \sin^4 x \\ &= \cos^5 x - 10\cos^3 x (1 - \cos^2 x) + 5\cos x (1 - 2\cos^2 x + \cos^4 x) \\ &= \cos^5 x - 10\cos^3 x + 10\cos^5 x + 5\cos x - 10\cos^3 x + 5\cos^5 x \\ &= 16\cos^5 x - 20\cos^3 x + 5\cos x \end{aligned}$$

- (b) Hence, show that $\cos 18^\circ = \frac{\sqrt{10+2\sqrt{5}}}{4}$.

4

$$\cos(5 \times 18^\circ) = 16\cos^5 18^\circ - 20\cos^3 18^\circ + 5\cos 18^\circ$$

$$0 = 16\cos^5 18^\circ - 20\cos^3 18^\circ + 5\cos 18^\circ$$

Now $\cos 18^\circ \neq 0$, so

$$0 = 16\cos^4 18^\circ - 20\cos^2 18^\circ + 5$$

$$\cos^2 18^\circ = \frac{20 \pm \sqrt{400 - 320}}{32}$$

$$= \frac{20 + \sqrt{16 \cdot 5}}{32}$$

$$= \frac{10 + 2\sqrt{5}}{16}$$

$$\therefore \cos 18^\circ = \frac{\sqrt{10+2\sqrt{5}}}{4}$$

by calculator
methods, deduce to take
+ve square root.

now $\cos 18^\circ > 0$,
so, take positive option.

Question 22 (10 marks)

A projectile is launched from the ground with an initial velocity of u m/s at an angle of θ to the horizontal. The projectile experiences the effect of gravity, and a resistance proportional to its velocity in both the horizontal and vertical directions.

The equations of motion are given by $\mathbf{a} = -0.1\mathbf{i} - 10\mathbf{j}$, where \mathbf{a} is the acceleration vector.



- (a) Show that the velocity vector $\mathbf{v} = e^{-0.1t}(\mathbf{u} + \mathbf{j}) - \mathbf{j}$, where $\mathbf{u} = u \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$, satisfies the equations of motion and initial conditions. 3

When $t = 0$,

$$\mathbf{v} = u \cos \theta \mathbf{i} + u (\sin \theta + 1) \mathbf{j}$$

$$= u \cos \theta \mathbf{i} + u \sin \theta \mathbf{j}$$

which satisfies the initial condition on \mathbf{v} .

$$\mathbf{a} = \frac{d\mathbf{v}}{dt} = -0.1 e^{-0.1t} (\mathbf{u} + \mathbf{j})$$

$$= -0.1 [e^{-0.1t} (\mathbf{u} + \mathbf{j}) - \mathbf{j}]$$

$$\Rightarrow \mathbf{a} = -0.1 \mathbf{i} - 0.1 \mathbf{j}$$

either I misread or misinterpreted the question, or this should say $(-10\mathbf{j})$

- (b) Show that the projectile reaches its peak (maximum height) after $10 \ln(u \sin \theta + 1)$ seconds. 3

$$\text{Peak reached when } v_y = e^{-0.1t} (u \sin \theta + 1) - 1 = 0$$

$$\text{consider } v_y \text{ when } t = 10 \ln(u \sin \theta + 1):$$

$$v_y = (u \sin \theta + 1)^{-10} (u \sin \theta + 1) - 1$$

$$= (u \sin \theta + 1)^0 - 1$$

$$= 0$$

\therefore peak after $10 \ln(u \sin \theta + 1)$ seconds.

Question 22 continues on page 17

Question 22 (continued)

- (c) Let the speed of the projectile at its peak be w m/s.

4

Show that $\lim_{u \rightarrow \infty} w = \cot \theta$.

$$v_x = u \cos \theta e^{-0.1t}$$

$$\text{When } t = 10 \ln(u \sin \theta + 1);$$

$$v_x = u \cos \theta (u \sin \theta + 1)^{-1}$$

$$= \frac{u \cos \theta}{u \sin \theta + 1} = w$$

$$\lim_{u \rightarrow \infty} w = \lim_{u \rightarrow \infty} \frac{\cos \theta}{\sin \theta + 1/u}$$


$$= \frac{\cos \theta}{\sin \theta}$$

$$= \cot \theta.$$

End of Question 22

Question 23 (12 marks)

Recall that $x \in \mathbb{Q} \iff \exists p \in \mathbb{Z}, q \in \mathbb{Z}^+ : x = \frac{p}{q}$.



- (a) Show that $\forall r \in \mathbb{Z}^+ \exists n \in \mathbb{Z}^+ : 0 < \int_0^1 x^n e^x dx < \frac{1}{r}$.

3

$x^n e^x$ is concave up, monotonic increasing in $x \in [0, 1]$.

clearly $\int_0^1 x^n e^x dx > 0$ as $x^n e^x > 0$ in $x \in [0, 1]$

as $e < 3$, $e^x < 3$ for $x \in [0, 1]$

hence: $0 < \int_0^1 x^n e^x dx < \int_0^1 3 x^n dx$

$$0 < \int_0^1 x^n e^x dx < \frac{3}{n+1}$$

now choose n such that $\frac{3}{n+1} < \frac{1}{r} : n > \frac{n+1}{3}$

$$n < 3r - 1$$

Hence $\forall r \in \mathbb{Z}^+ 0 < \int_0^1 x^n e^x dx < \frac{1}{r}$.

- (b) Prove, using induction, that for all integers $n \geq 0$, $\exists \alpha, \beta \in \mathbb{Z} : \int_0^1 x^n e^x dx = \alpha + \beta e$.

4

Base case $n=0$: $I_0 = \int_0^1 e^x dx = e - 1 \Rightarrow \alpha = -1, \beta = 1$ ✓

Assume $\exists k \in \mathbb{N} : \exists \alpha, \beta \in \mathbb{Z}^+ : I_k = \int_0^1 x^k e^x dx = \alpha + \beta e$.

consider $I_{k+1} = \int_0^1 x^{k+1} e^x dx$

$$= [x^{k+1} e^x]_0^1 - \int_0^1 (k+1) x^k e^x dx$$

$$= e - (k+1) I_k$$

$$= e - (k+1)(\alpha + \beta e)$$

$$= -(k+1)\alpha + (1 - (k+1)\beta)e$$

Now $-(k+1)\alpha \in \mathbb{Z}$ as $k, \alpha \in \mathbb{Z}$

and $(1 - (k+1)\beta) \in \mathbb{Z}$ as $k, \beta \in \mathbb{Z}$.

\therefore true by mathematical induction

Question 23 continues on page 19

Question 23 (continued)

- (c) Let $p \in \mathbb{Z}$, $q \in \mathbb{Z}^+$, $x = \frac{p}{q} \in \mathbb{Q}$.

Show that $\forall \alpha, \beta \in \mathbb{Z} : \left\{ \alpha + \beta x \neq 0 \implies |\alpha + \beta x| \geq \frac{1}{q} \right\}$.

2

Consider $q^2(\alpha + \beta x)^2$. Note $q \neq 0$. Suppose $\alpha + \beta x \neq 0$. Then $q^2(\alpha + \beta x)^2 \neq 0$. Now $q^2(\alpha + \beta x)^2 = (\alpha q + \beta p)^2 \in \mathbb{Z}^+$, as $\alpha, \beta, p, q \in \mathbb{Z}$, and $q^2(\alpha + \beta x)^2 \neq 0$. So $q^2(\alpha + \beta x)^2 \geq 1 \implies (\alpha + \beta x)^2 \geq \frac{1}{q^2} \implies |\alpha + \beta x| \geq \frac{1}{q}$ as required.

- (d) Hence, prove that $e \notin \mathbb{Q}$.

3

from a) $0 < \int_0^1 x^n e^x dx < \frac{1}{n}$
 from b) $\int_0^1 x^n e^x dx = \alpha + \beta e$ for some $\alpha, \beta \in \mathbb{Z}$.
 from c), suppose e is rational; $e = \frac{p}{q}$, as defined previously.
 then $|\alpha + \beta e| \geq \frac{1}{q}$

Using a) and b): $0 < \alpha + \beta e < \frac{1}{q}$
 which contradicts c): $\alpha + \beta e \geq \frac{1}{q}$ or $\alpha + \beta e \leq -\frac{1}{q}$.
 $\therefore e$ is not rational, i.e. $e \notin \mathbb{Q}$.

End of Question 23

End of Paper

A large rectangular box containing 25 horizontal dotted lines for writing.

A large rectangular box containing 25 horizontal dotted lines for writing.

A large rectangular box containing 25 horizontal dotted lines for writing.

A large rectangular box containing 25 horizontal dotted lines for writing.

A large rectangular box containing 25 horizontal dotted lines for writing.

**Mathematics Advanced**
Mathematics Extension 1
Mathematics Extension 2**REFERENCE SHEET****Measurement****Length**

$$l = \frac{\theta}{360} \times 2\pi r$$

Area

$$A = \frac{\theta}{360} \times \pi r^2$$

$$A = \frac{h}{2}(a + b)$$

Surface area

$$A = 2\pi r^2 + 2\pi rh$$

$$A = 4\pi r^2$$

Volume

$$V = \frac{1}{3}Ah$$

$$V = \frac{4}{3}\pi r^3$$

Functions

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\text{For } ax^3 + bx^2 + cx + d = 0:$$

$$\alpha + \beta + \gamma = -\frac{b}{a}$$

$$\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a}$$

$$\text{and } \alpha\beta\gamma = -\frac{d}{a}$$

Relations

$$(x - h)^2 + (y - k)^2 = r^2$$

Financial Mathematics

$$A = P(1 + r)^n$$

Sequences and series

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d] = \frac{n}{2}(a + l)$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} = \frac{a(r^n - 1)}{r - 1}, r \neq 1$$

$$S = \frac{a}{1 - r}, |r| < 1$$

Logarithmic and Exponential Functions

$$\log_a a^x = x = a^{\log_a x}$$

$$\log_a x = \frac{\log_b x}{\log_b a}$$

$$a^x = e^{x \ln a}$$

Trigonometric Functions

$$\sin A = \frac{\text{opp}}{\text{hyp}}, \quad \cos A = \frac{\text{adj}}{\text{hyp}}, \quad \tan A = \frac{\text{opp}}{\text{adj}}$$

$$A = \frac{1}{2} ab \sin C$$



$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$l = r\theta$$

$$A = \frac{1}{2} r^2 \theta$$



Trigonometric identities

$$\sec A = \frac{1}{\cos A}, \quad \cos A \neq 0$$

$$\operatorname{cosec} A = \frac{1}{\sin A}, \quad \sin A \neq 0$$

$$\cot A = \frac{\cos A}{\sin A}, \quad \sin A \neq 0$$

$$\cos^2 x + \sin^2 x = 1$$

Compound angles

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\text{If } t = \tan \frac{A}{2} \text{ then } \sin A = \frac{2t}{1 + t^2}$$

$$\cos A = \frac{1 - t^2}{1 + t^2}$$

$$\tan A = \frac{2t}{1 - t^2}$$

$$\cos A \cos B = \frac{1}{2} [\cos(A - B) + \cos(A + B)]$$

$$\sin A \sin B = \frac{1}{2} [\cos(A - B) - \cos(A + B)]$$

$$\sin A \cos B = \frac{1}{2} [\sin(A + B) + \sin(A - B)]$$

$$\cos A \sin B = \frac{1}{2} [\sin(A + B) - \sin(A - B)]$$

$$\sin^2 nx = \frac{1}{2} (1 - \cos 2nx)$$


$$\cos^2 nx = \frac{1}{2} (1 + \cos 2nx)$$

Statistical Analysis

$$z = \frac{x - \mu}{\sigma}$$

An outlier is a score
less than $Q_1 - 1.5 \times IQR$
or
more than $Q_3 + 1.5 \times IQR$

Normal distribution



- approximately 68% of scores have z-scores between -1 and 1
- approximately 95% of scores have z-scores between -2 and 2
- approximately 99.7% of scores have z-scores between -3 and 3

$$E(X) = \mu$$

$$\operatorname{Var}(X) = E[(X - \mu)^2] = E(X^2) - \mu^2$$

Probability

$$P(A \cap B) = P(A)P(B)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad P(B) \neq 0$$

Continuous random variables

$$P(X \leq r) = \int_a^r f(x) dx$$

$$P(a < X < b) = \int_a^b f(x) dx$$

Binomial distribution

$$P(X = r) = {}^n C_r p^r (1 - p)^{n-r}$$

$$X \sim \operatorname{Bin}(n, p)$$

$$\Rightarrow P(X = x)$$

$$= \binom{n}{x} p^x (1 - p)^{n-x}, \quad x = 0, 1, \dots, n$$

$$E(X) = np$$

$$\operatorname{Var}(X) = np(1 - p)$$

Differential Calculus

Function

Derivative

$$y = f(x)^n$$

$$\frac{dy}{dx} = n f'(x) [f(x)]^{n-1}$$

$$y = uv$$

$$\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$y = g(u) \text{ where } u = f(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$y = \frac{u}{v}$$

$$\frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$y = \sin f(x)$$

$$\frac{dy}{dx} = f'(x) \cos f(x)$$

$$y = \cos f(x)$$

$$\frac{dy}{dx} = -f'(x) \sin f(x)$$

$$y = \tan f(x)$$

$$\frac{dy}{dx} = f'(x) \sec^2 f(x)$$

$$y = e^{f(x)}$$

$$\frac{dy}{dx} = f'(x) e^{f(x)}$$

$$y = \ln f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{f(x)}$$

$$y = a^{f(x)}$$

$$\frac{dy}{dx} = (\ln a) f'(x) a^{f(x)}$$

$$y = \log_a f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{(\ln a) f(x)}$$

$$y = \sin^{-1} f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{\sqrt{1 - [f(x)]^2}}$$

$$y = \cos^{-1} f(x)$$

$$\frac{dy}{dx} = -\frac{f'(x)}{\sqrt{1 - [f(x)]^2}}$$

$$y = \tan^{-1} f(x)$$

$$\frac{dy}{dx} = \frac{f'(x)}{1 + [f(x)]^2}$$

Integral Calculus

$$\int f'(x) [f(x)]^n dx = \frac{1}{n+1} [f(x)]^{n+1} + c$$

where $n \neq -1$

$$\int f'(x) \sin f(x) dx = -\cos f(x) + c$$

$$\int f'(x) \cos f(x) dx = \sin f(x) + c$$

$$\int f'(x) \sec^2 f(x) dx = \tan f(x) + c$$

$$\int f'(x) e^{f(x)} dx = e^{f(x)} + c$$

$$\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c$$

$$\int f'(x) a^{f(x)} dx = \frac{a^{f(x)}}{\ln a} + c$$

$$\int \frac{f'(x)}{\sqrt{a^2 - [f(x)]^2}} dx = \sin^{-1} \frac{f(x)}{a} + c$$

$$\int \frac{f'(x)}{a^2 + [f(x)]^2} dx = \frac{1}{a} \tan^{-1} \frac{f(x)}{a} + c$$

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

$$\int_a^b f(x) dx$$

$$\approx \frac{b-a}{2n} \left\{ f(a) + f(b) + 2[f(x_1) + \dots + f(x_{n-1})] \right\}$$

where $a = x_0$ and $b = x_n$

Combinatorics

$${}^nP_r = \frac{n!}{(n-r)!}$$

$$\binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!}$$

$$(x+a)^n = x^n + \binom{n}{1}x^{n-1}a + \cdots + \binom{n}{r}x^{n-r}a^r + \cdots + a^n$$

Vectors

$$|\underline{u}| = |x\underline{i} + y\underline{j}| = \sqrt{x^2 + y^2}$$

$$\underline{u} \cdot \underline{v} = |\underline{u}| |\underline{v}| \cos \theta = x_1x_2 + y_1y_2,$$

$$\text{where } \underline{u} = x_1\underline{i} + y_1\underline{j}$$

$$\text{and } \underline{v} = x_2\underline{i} + y_2\underline{j}$$

$$\underline{r} = \underline{a} + \lambda \underline{b}$$

Complex Numbers

$$\begin{aligned} z &= a + ib = r(\cos \theta + i \sin \theta) \\ &= re^{i\theta} \end{aligned}$$

$$\begin{aligned} [r(\cos \theta + i \sin \theta)]^n &= r^n(\cos n\theta + i \sin n\theta) \\ &= r^n e^{in\theta} \end{aligned}$$

Mechanics

$$\frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx} \left(\frac{1}{2} v^2 \right)$$

$$x = a \cos(nt + \alpha) + c$$

$$x = a \sin(nt + \alpha) + c$$

$$\ddot{x} = -n^2(x - c)$$