



2023

## TRIAL HIGHER SCHOOL CERTIFICATE EXAMINATION

DO NOT REMOVE PAPER FROM EXAMINATION ROOM

--	--	--	--	--

Centre Number

--	--	--	--	--	--	--	--

Student Number

# Mathematics Extension 2

Morning Session  
Monday, 7 August 2023

---

**General Instructions**

- Reading time – 10 minutes
- Working time – 3 hours
- Write using black pen
- Calculators approved by NESA may be used
- A reference sheet is provided
- Use the Multiple-Choice Answer Sheet provided
- For questions in Section II, show relevant mathematical reasoning and/or calculations
- Write your Centre Number and Student Number at the top of this page

---

**Total marks:**  
100

**Section I – 10 marks (pages 2–6)**

- Attempt Questions 1–10
- Allow about 15 minutes for this section

**Section II – 90 marks (pages 7–13)**

- Attempt Questions 11–16
- Allow about 2 hours and 45 minutes for this section

**Disclaimer**

These "Trial" Higher School Certificate Examinations have been prepared by CSSA, a division of Catholic Schools NSW Limited. Every effort has been made to prepare these "Trial" Higher School Certificate Examinations in accordance with the NSW Education Standards Authority (NESA) documents, Principles for Setting HSC Examinations in a Standards Referenced Framework and Principles for Developing Marking Guidelines Examinations in a Standards Referenced Framework. No guarantee or warranty is made or implied that the "Trial" HSC Examination papers mirror in every respect the actual HSC Examination papers in any or all courses to be examined. These papers do not constitute "advice" nor can they be construed as authoritative interpretations of NESA intentions. Catholic Schools NSW Limited accepts no liability for any reliance, use or purpose related to these "Trial" HSC Examination papers. Advice on HSC examination issues is only to be obtained from the NESA.

## Section I

10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the Multiple-Choice Answer Sheet for Questions 1–10

---

- 1** A composite number is a positive integer that can be formed by multiplying two smaller integers, both larger than 1. Which set of positive integers  $n$  is this statement true for?

"The number  $n^2 - 1$  is composite."

- A.  $n \geq 1$
- B.  $n \geq 2$
- C.  $n \geq 3$
- D.  $n \geq 4$

- 2** Consider the following two statements.

Statement  $X$ : The sum of two rational numbers is rational.

Statement  $Y$ : The sum of two irrational numbers is irrational.

Which of the statements is true?

- A.  $X$  and  $Y$  are both true.
- B.  $X$  is true and  $Y$  is false.
- C.  $X$  is false and  $Y$  is true.
- D.  $X$  and  $Y$  are both false.

- 3 Maeve is asked to prove or disprove the following statement for  $x$  and  $y$  positive real numbers:  $\frac{1}{x^2} + \frac{1}{y^2} \geq \frac{2}{xy}$ . Her working is shown below.

$$\text{We know: } \frac{1}{(x-y)^2} \geq 0 \Leftrightarrow \frac{1}{x^2 - 2xy + y^2} \geq 0$$

$$\Leftrightarrow \frac{1}{x^2 + y^2} \geq \frac{1}{2xy} \quad \text{line A}$$

$$\Leftrightarrow x^2 + y^2 \leq 2xy \quad \text{line B}$$

$$\Leftrightarrow \frac{x^2 + y^2}{x^2 y^2} \leq \frac{2xy}{x^2 y^2} \quad \text{line C}$$

$$\Leftrightarrow \frac{1}{x^2} + \frac{1}{y^2} \leq \frac{2}{xy} \quad \text{line D}$$

So the statement is false.

On which line of working has Maeve made a mistake?

- A. line A
- B. line B
- C. line C
- D. line D

- 4 The amount of apples, bananas and oranges sold by a fruit seller over a year is shown in the table below.

Fruit	Amount Sold (tonnes)	Profit (\$/tonne)
Apples	25	530
Bananas	55	380
Oranges	10	410

Let  $a = \begin{pmatrix} 25 \\ 55 \\ 10 \end{pmatrix}$ ,  $b = \begin{pmatrix} 530 \\ 380 \\ 410 \end{pmatrix}$  and  $c = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ . Which of the following expressions calculates the average profit in dollars per tonne of fruit sold over the year?

- A.  $\frac{a \cdot c}{b \cdot c}$
- B.  $\frac{b \cdot c}{a \cdot c}$
- C.  $\frac{a \cdot b}{b \cdot c}$
- D.  $\frac{a \cdot b}{a \cdot c}$


5 Which of the following expressions is equivalent to  $\int \ln(x^2 + 1) dx$ ?

- A.  $x \ln(x^2 + 1) - 2x + 2 \tan^{-1} x + c$
- B.  $x \ln(x^2 + 1) - \ln(x^2 + 1) + c$
- C.  $\ln(x^2 + 1) - 2x + 2 \tan^{-1} x + c$
- D.  $\ln(x^2 + 1) - x \ln(x^2 + 1) + c$

6 Consider the complex, non-real cube roots of unity  $\omega$  and  $\omega^2$ . What is the value of  $(1 - \omega + \omega^2)(1 + \omega - \omega^2)$ ?

- A. 0
- B. 1
- C. 2
- D. 4

- 7 The complex numbers  $a$ ,  $b$ ,  $c$  and  $d$  and  $z$  are solutions to  $z^7 = 1$  as shown in the Argand diagram below.



Which of the following is a cube root of  $z$ ?

- A.  $a$
- B.  $b$
- C.  $c$
- D.  $d$

- 8 Consider the transitive property of implication:

$$(a \Rightarrow b \text{ AND } b \Rightarrow c) \Rightarrow (a \Rightarrow c).$$

Which of the following is the contrapositive of the transitive property of implication?

- A.  $\neg(a \Rightarrow c) \Rightarrow (\neg(a \Rightarrow b) \text{ OR } \neg(b \Rightarrow c))$
- B.  $\neg(a \Rightarrow c) \Rightarrow (\neg(a \Rightarrow b) \text{ AND } \neg(b \Rightarrow c))$
- C.  $\neg(a \Rightarrow c) \Rightarrow \neg(\neg(a \Rightarrow b) \text{ OR } \neg(b \Rightarrow c))$
- D.  $\neg(a \Rightarrow c) \Rightarrow \neg(\neg(a \Rightarrow b) \text{ AND } \neg(b \Rightarrow c))$

- 9** An object of unit mass falls from rest with downward velocity of  $v \text{ ms}^{-1}$ . The object undergoes acceleration due to gravity of  $g \text{ ms}^{-2}$ . It also experiences air resistance of magnitude  $kv \text{ ms}^{-2}$ , in the opposite direction to the velocity, where  $k$  is some positive constant.

Which is the correct expression for the distance  $x \text{ m}$  it has fallen in terms of its downward velocity  $v \text{ ms}^{-1}$ ?

A.  $x = -\frac{v}{k} - \frac{g}{k^2} \ln \left| \frac{g}{g - kv} \right|$

B.  $x = -\frac{v}{k} + \frac{g}{k^2} \ln \left| \frac{g}{g - kv} \right|$

C.  $x = \frac{v}{k} - \frac{g}{k^2} \ln \left| \frac{g - kv}{g} \right|$

D.  $x = \frac{v}{k} + \frac{g}{k^2} \ln \left| \frac{g - kv}{g} \right|$

- 10** The function  $F(x)$  is a primitive of  $f(x)$ , that is  $F'(x) = f(x)$ .  
Which of the following is true?

A.  $\int \left( \frac{d}{dx} \int_a^b f(x) dx \right) dx = f(b) - f(a)$

B.  $\int_a^b \left( \frac{d}{dx} \int f(x) dx \right) dx = f(b) - f(a)$

C.  $\frac{d}{dx} \int \left( \int_a^b f(x) dx \right) dx = F(b) - F(a)$

D.  $\frac{d}{dx} \int_a^b \left( \int f(x) dx \right) dx = F(b) - F(a)$

## Section II

90 marks

Attempt Questions 11–16

Allow about 2 hours and 45 minutes for this section


Answer each question in a separate writing booklet. Extra writing booklets are available.

Your responses for Questions 11–16 should include relevant mathematical reasoning and/or calculations.

---

### Question 11 (15 marks)

- (a) Express  $2\sqrt{2}e^{-\frac{3\pi}{4}i}$  in the form  $x + iy$ . 2
- (b) Consider the two points  $A(2, 2, 2)$  and  $B(2, -2, 2)$ .
- (i) Find  $\vec{AB}$ . 1
  - (ii) Find  $|\vec{AB}|$ . 1
  - (iii) Find  $\angle AOB$ . Give your answer correct to the nearest degree. 2
- (c) Consider the complex number  $z = 2e^{0.4i}$ , as sketched below. 3



Copy and clearly label this Argand diagram, and sketch the four points represented by  $z$ ,  $\bar{z}$ ,  $-\bar{z}$ , and  $z - \bar{z}$ .

- (d) A particle starts at rest from the origin and moves in a straight line such that its displacement  $x$  metres at time  $t$  seconds is determined by the equation  $\ddot{x} = -16x$ . How long will it take for the particle to first return to the origin? 3
- (e) Show for any complex numbers  $z$  and  $w$ , that  $\overline{zw} = \bar{z} \times \bar{w}$ . 3

**Question 12** (15 marks)

- (a) Consider the two lines

$$L_1 : \begin{pmatrix} 7 \\ -1 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 3 \\ -5 \end{pmatrix} \quad \text{and} \quad L_2 : \begin{pmatrix} -1 \\ 2 \\ -6 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix}.$$

(i) Find the value of  $k$  given  $B(9, k, 24)$  lies on  $L_2$ . 2

(ii) Find the point of intersection of  $L_1$  and  $L_2$ . 3

(b) Find  $\int_0^{\frac{1}{\sqrt{2}}} \frac{x^3}{\sqrt{1-x^2}} dx$ . 4

(c) Solve  $z^2 + (7-i)z + 16 + 4i = 0$ . 3


(d) Find  $\int \frac{x^3 - 2}{x^3 + x} dx$ . 3

**Question 13** (15 marks)

- (a) Prove that  $\left(a + \frac{1}{b}\right)\left(b + \frac{1}{a}\right) \geq 4$  where  $a, b > 0$ . 2
- (b) The graph of a polynomial function  $f(x) = (x+3)(x-2)(x^2+bx+c)$  has a  $y$ -intercept of  $-6$  and passes through the point  $(1, -4)$ .
- (i) Find the two complex roots of the equation  $f(x) = 0$ . 2
  - (ii) Express these two complex roots in the form  $r(\cos \theta + i \sin \theta)$ . 2
  - (iii) Plot all four solutions to  $f(x) = 0$  on an Argand diagram. 2
  - (iv) Write down the name of the quadrilateral formed by these four points. 1
- (c) Consider the sphere with vector equation  $\left| \underline{x} - \begin{pmatrix} 3 \\ -12 \\ 4 \end{pmatrix} \right| = 3$ .
- (i) Show the point  $(5, -10, 3)$  lies on the sphere. 1
  - (ii) Find the point on the sphere farthest from the origin. 2
- (d) Prove by mathematical induction that  $3n^2 - 3n \leq 2^n - 1$  for  $n \geq 7$ . 3

**Question 14** (15 marks)

- (a) The number  $z$  is a fifth root of unity where  $z \neq 1$ , that is  $z^5 = 1$ .
- (i) Show that  $(z + z^{-1})^2 + (z + z^{-1}) - 1 = 0$ . 2
- (ii) If  $z = e^{i\theta}$ , show  $\cos \theta = \frac{z + z^{-1}}{2}$ . 1
- (iii) Hence show  $\cos \frac{2\pi}{5} = \frac{\sqrt{5} - 1}{4}$ . 2
- (iv) Consider the five fifth roots of unity  $z_1, z_2, z_3, z_4$  and  $z_5$  as shown in the diagram below. 3



$$\text{Show that } \left| \frac{z_3 - z_1}{z_2 - z_1} \right| = \frac{1 + \sqrt{5}}{2}.$$

- (b) The position of an object after  $t$  seconds is given by the vector equation
- $$\underline{r} = \cos \frac{\pi t}{4} \underline{i} + \left( \cos \frac{\pi t}{4} + \sin \frac{\pi t}{4} \right) \underline{j} + \sin \frac{\pi t}{4} \underline{k}.$$
- (i) What is the position of the object after 3 seconds? 1
- (ii) Find the vector equation of the tangent to the path taken by the object after 3 seconds. 3
- (c) Evaluate  $\int_0^9 \frac{1}{\sqrt{1+\sqrt{x}}} dx$ . 3


**Question 15** (15 marks)

- (a) Consider the function  $f(x) = e^{-x} \cos x$ , with domain  $x \in \left[0, \frac{3\pi}{2}\right]$ . 4

Show that the ratio of the area above the  $x$ -axis to the area below the  $x$ -axis is

$$\frac{e^\pi \left(e^{\frac{\pi}{2}} + 1\right)}{e^\pi + 1}.$$

- (b) An emergency ramp is a steep ramp designed for cars on a highway to safely stop if their brakes have failed. A car rolls onto an emergency ramp inclined at  $30^\circ$  to the horizontal at  $144 \text{ km/h}$  without its brakes on as shown in the diagram below. 4



The car experiences resistance due to friction of magnitude  $0.05mv$  newtons where  $m \text{ kg}$  is the car's mass and  $v \text{ ms}^{-1}$  is the car's velocity at time  $t$  seconds after entering the ramp.

Assuming the acceleration due to gravity is  $10 \text{ ms}^{-2}$ , calculate how far the car travels up the ramp before stopping. Give your answer correct to the nearest metre.

- (c) Let  $I_n = \int \frac{\cos nx}{\sin x} dx$ .

(i) Show that  $\cos((n-2)x) - \cos nx = 2 \sin((n-1)x) \sin x$ . 2

(ii) Show that  $I_n - I_{n-2} = \frac{2 \cos((n-1)x)}{n-1} + c$ . 2

(iii) Hence, or otherwise, evaluate  $\int_0^{\frac{\pi}{3}} \frac{\cos 2x - \cos 6x}{\sin x} dx$ . 3

**Question 16** (15 marks)


- (a) Consider the tile below consisting of three equilateral triangles of side length 1 unit.

3

Prove the following result for positive integers  $n$ , using mathematical induction:

An equilateral triangle of side length  $2^n$  units may be covered by the tiles above (in any orientation) such that a single equilateral triangle of side length 1 unit is left over at one of the vertices. The tiles may not overlap.

- (b) An ant follows a spiral path up a cylindrical column from
- $P(0, 7, 0)$
- to the point
- $Q$
- as shown in the diagram below.

The ant's position  $\underline{r}$  in centimetres after  $t$  seconds is given by the vector equation below.

$$\underline{r}(t) = 7 \sin \frac{\pi t}{32} \mathbf{i} + 7 \cos \frac{\pi t}{32} \mathbf{j} + \frac{t}{15} \mathbf{k}$$

- (i) Find the coordinates of the point  $Q$  if  $|\overrightarrow{OQ}| = 25$ . 3
- (ii) How many times has the ant crossed the line  $x = 7$  on its journey to  $Q$ ? 2
- (iii) The ant crawls back from  $Q$  to  $P$  along the shortest path possible. How far did it crawl on this leg of its journey? Give your answer correct to one decimal place. 2

**Question 16 continues on page 13**

Question 16 (continued)

- (c) A projectile of unit mass is launched vertically upwards from a horizontal plane with initial speed  $v_0 \text{ ms}^{-1}$ . The projectile experiences a resistive force that causes an acceleration with a magnitude of  $0.01v^2 \text{ ms}^{-2}$ , where  $v \text{ ms}^{-1}$  is the velocity of the projectile. The acceleration due to gravity is  $10 \text{ ms}^{-2}$ .

5

The projectile lands on the horizontal plane at  $\frac{1}{7}$  the speed that it was launched.

Find the value of  $v_0$ , correct to 1 decimal place.

---

End of Examination

BLANK PAGE

BLANK PAGE

## EXAMINERS

David Houghton (Convenor)  
Geoff Carroll  
Stephen Ewington  
Rebekah Johnson  
Svetlana Onisczenko

Oxley College, Burradoo  
Sydney Grammar School, Darlinghurst  
Ascham School, Edgecliff  
Loreto Kirribilli, Kirribilli  
Meriden School, Strathfield

### Additional Disclaimer

Users of CSSA Trial HSC Examinations are advised that due to changing NESA examination policies, it cannot be assumed that CSSA Trial HSC Examinations and NESA Examinations will, from year to year, always fully align with respect to either or both the format and content of examination questions. Candidates for HSC examinations and their teachers should always anticipate a dynamic assessment environment.

### Copyright Notice

1. The copyright in this examination paper is that of Catholic Schools NSW Limited ACN 619 593 369 trading as CSSA - ©2023 Catholic Schools NSW.
2. This examination paper may only be used in accordance with the CSSA Trial HSC Examination Terms & Conditions (Terms). Those Terms should be read in full. The Terms contain a number of conditions including those relating to:
  - a) how this examination paper may be used for trial HSC examinations and/or as a practice paper;
  - b) when copies may be made of this examination paper;
  - c) the security and storage requirements for this examination paper; and
  - d) guidelines for the conduct of trial HSC examinations.
3. Some of the material in this examination paper may have been copied and communicated to you in accordance with the statutory licence in section 113P of the Copyright Act 1988 (Cth) (Act). Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.
4. Do not remove this notice.

# Mathematics Extension 2

**Section I  
10 marks**
**Multiple Choice Answer Key**

Question	Answer	Outcomes Assessed	Targeted Performance Bands
1	C	MEX12-2	E1-E2
2	B	MEX12-2	E1-E2
3	A	MEX12-2	E1
4	D	MEX12-3	E2
5	A	MEX12-5	E2-E3
6	D	MEX12-4	E2
7	C	MEX12-4	E3-E4
8	A	MEX12-1	E3-E4
9	B	MEX12-6	E3-E4
10	C	MEX12-5	E4

**Question 1 (1 mark)**
*Outcomes Assessed: MEX12-2*
*Targeted Performance Bands: E1-E2*

Solution	Mark
$n^2 - 1 = (n+1)(n-1)$ . If $n+1 \geq 2$ , and $n-1 \geq 2$ , the definition is satisfied. Therefore $n \geq 3$ satisfies the definition.	1
Hence C	

**Question 2 (1 mark)**
*Outcomes Assessed: MEX12-2*
*Targeted Performance Bands: E1-E2*

Solution	Mark
The sum of two rational numbers is rational, hence statement X is true. However, the sum of two irrational numbers could be either rational or irrational. e.g. $(1 + \sqrt{2}) + (1 - \sqrt{2}) = 2 \in \mathbb{Q}$ whereas $\sqrt{2} + \sqrt{3} \notin \mathbb{Q}$ . Therefore Statement Y is false. Hence B	1

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific Trial HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

**Question 3 (1 mark)**

Outcomes Assessed: MEX12-2

Targeted Performance Bands: E1

Solution	Mark
$\frac{1}{x^2 - 2xy + y^2} \neq \frac{1}{x^2 + y^2} - \frac{1}{2xy}$ , hence the error is at line A.  Hence A	1

**Question 4 (1 mark)**

Outcomes Assessed: MEX12-3

Targeted Performance Bands: E2

Solution	Mark
$\begin{aligned} \frac{\text{\$ profit}}{\text{tonnes sold}} &= \frac{25 \times 530 + 55 \times 380 + 10 \times 410}{25 + 55 + 10} \\ &= \frac{\begin{pmatrix} 25 \\ 55 \\ 10 \end{pmatrix} \cdot \begin{pmatrix} 530 \\ 380 \\ 410 \end{pmatrix}}{\begin{pmatrix} 25 \\ 55 \\ 10 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}} = \frac{a \cdot b}{a \cdot c} \end{aligned}$  Hence D	1

**Question 5 (1 mark)**

Outcomes Assessed: MEX12-5

Targeted Performance Bands: E2-E3

Solution	Mark
$I = \int \ln(x^2 + 1) dx$  $\begin{aligned} I &= x \ln(x^2 + 1) - \int \frac{2x^2}{x^2 + 1} dx \\ &= x \ln(x^2 + 1) - 2 \int 1 - \frac{1}{x^2 + 1} dx \\ &= x \ln(x^2 + 1) - 2x + 2 \tan^{-1} x + c \end{aligned}$  Let $u = \ln(x^2 + 1)$ , $v = x$ $\text{So } du = \frac{2x}{x^2 + 1} , dv = dx$  Hence A	1

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

**Question 6 (1 mark)***Outcomes Assessed: MEX12-4**Targeted Performance Bands: E2*

Solution	Mark
$\omega^3 - 1 = 0$ $(\omega - 1)(\omega^2 + \omega + 1) = 0$ , and since $\omega \notin \mathbb{R}$ $\omega^2 = -(\omega + 1)$ Therefore $(1 - \omega + \omega^2)(1 + \omega - \omega^2) = (1 - \omega - \omega - 1)(-\omega^2 - \omega^2)$ $= -2\omega \times -2\omega^2$ $= 4\omega^3 = 4$ , since $\omega^3 = 1$ .  Hence D	1

**Question 7 (1 mark)***Outcomes Assessed: MEX12-4**Targeted Performance Bands: E3-E4*

Solution	Mark
$\arg z = -\frac{4\pi}{7}$ $\arg a = 0$ , hence $\arg(a^3) = 0 \neq \arg z$ . $\arg b = \frac{4\pi}{7}$ , hence $\arg(b^3) = \frac{12\pi}{7}$ , equivalent to $-\frac{2\pi}{7} \neq \arg z$ . $\arg c = -\frac{6\pi}{7}$ , hence $\arg(c^3) = -\frac{18\pi}{7}$ , equivalent to $-\frac{4\pi}{7} = \arg z$ . $\arg d = -\frac{2\pi}{7}$ , hence $\arg(d^3) = -\frac{6\pi}{7} \neq \arg z$ .  Hence C	1

**Question 8 (1 mark)***Outcomes Assessed: MEX12-1**Targeted Performance Bands: E3-E4*

Solution	Mark
Contrapositive is $\neg(a \Rightarrow c) \Rightarrow \neg(a \Rightarrow b \text{ AND } b \Rightarrow c)$ . Now, $\neg(p \text{ AND } q)$ is logically equivalent to $\neg p \text{ OR } \neg q$ , hence $\neg(a \Rightarrow c) \Rightarrow (\neg(a \Rightarrow b) \text{ OR } \neg(b \Rightarrow c))$ . Hence A	1

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of the NSW Education Standards Authority to provide marking criteria for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

**Question 9 (1 mark)**

Outcomes Assessed: MEX12-6

Targeted Performance Bands: E3-E4

Solution	Mark
<p>With downwards being positive, the acceleration due to resistance <math>kv</math> acts upwards, so:</p> $\ddot{v} = g - kv$ $v \frac{dv}{dx} = g - kv$ $\frac{dx}{dv} = \frac{v}{g - kv}$ $= -\frac{1}{k} \times \frac{-kv}{g - kv}$ $= -\frac{1}{k} \left( \frac{g - kv}{g - kv} - \frac{g}{g - kv} \right)$ $\frac{dx}{dv} = -\frac{1}{k} - \frac{g}{k^2} \times \frac{-k}{g - kv}$ $x = -\frac{v}{k} - \frac{g}{k^2} \ln g - kv  + c$ <p>Now, when <math>x = 0, v = 0</math>, so</p> $0 = 0 - \frac{g}{k^2} \ln g + c$ $x = -\frac{v}{k} - \frac{g}{k^2} \ln g - kv  + \frac{g}{k^2} \ln g$ $x = -\frac{v}{k} + \frac{g}{k^2} \ln \left  \frac{g}{g - kv} \right $	1

Hence B

**Question 10 (1 mark)**

Outcomes Assessed: MEX12-5

Targeted Performance Bands: E4

Solution	Mark
$\text{LHS of A} = \int \left( \frac{d}{dx} \int_a^b f(x) dx \right) dx = \int \left( \frac{d}{dx} [F(b) - F(a)] \right) dx$ $= \int (0) dx = c \neq \text{RHS}$	
$\text{LHS of B} = \int_a^b \left( \frac{d}{dx} \int f(x) dx \right) dx = \int_a^b \left( \frac{d}{dx} (F(x) + c) \right) dx$ $= \int_a^b (f(x)) dx = F(b) - F(a) \neq \text{RHS}$	1
$\text{LHS of C} = \frac{d}{dx} \int \left( \int_a^b f(x) dx \right) dx = \frac{d}{dx} \int (F(b) - F(a)) dx$ $= \frac{d}{dx} [xF(b) - xF(a) + c] = F(b) - F(a) = \text{RHS}$	
$\text{LHS of D} = \frac{d}{dx} \int_a^b \left( \int f(x) dx \right) dx = \frac{d}{dx} \int_a^b (F(x) + c) dx$ $= \frac{d}{dx} [\text{some real number}] = 0 \neq \text{RHS}$	

Hence C

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

## Section II

90 marks

### Question 11 (15 marks)

Question 11(a) (2 marks)

*Outcomes Assessed:* MEX12-4

*Targeted Performance Bands:* E1

Criteria	Marks
• correct solution	2
• progress towards correct solution	1

*Sample Answer:*

$$2\sqrt{2}e^{-\frac{3\pi}{4}i} = 2\sqrt{2}\left(-\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}\right) = -2 - 2i$$

Question 11(b) (i) (1 mark)

*Outcomes Assessed:* MEX12-3

*Targeted Performance Bands:* E1

Criteria	Mark
• correct solution	1

*Sample Answer:*

$$\vec{AB} = 0\hat{i} - 4\hat{j} + 0\hat{k}$$

Question 11(b) (ii) (1 mark)

*Outcomes Assessed:* MEX12-3

*Targeted Performance Bands:* E1

Criteria	Mark
• correct solution	1

*Sample Answer:*

$$|\vec{AB}| = \sqrt{4^2} = 4$$

Question 11(b) (iii) (2 marks)

*Outcomes Assessed:* MEX12-3

*Targeted Performance Bands:* E2

Criteria	Marks
• correct solution with justification (dot product or cosine rule)	2
• progress towards correct solution	1

*Sample Answer:*

$$\cos \angle AOB = \frac{\vec{OA} \cdot \vec{OB}}{|\vec{OA}| \times |\vec{OB}|} = \frac{4 - 4 + 4}{(\sqrt{2^2 + 2^2 + 2^2})^2} = \frac{1}{3}. \text{ Hence } \angle AOB = \cos^{-1}\left(\frac{1}{3}\right) \approx 71^\circ.$$

**Disclaimer**  
The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific answers to all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 11(c) (3 marks)


*Outcomes Assessed:* MEX12-4

*Targeted Performance Bands:* E2

Criteria	Marks
• correct sketch including the point $z - \bar{z}$ , not just the vector from $\bar{z}$ to $z$	3
• two correct points, probably $z$ and $\bar{z}$ as reflections over $x$ -axis	2
• at least one correct point, representing modulus and argument	1

*Sample Answer:*

The image to the right displays the four points. Note that the question is not asking for the vector from  $\bar{z}$  to  $z$ , but rather the number  $z - \bar{z}$ , which sits between  $i$  and  $2i$ .



Question 11(d) (3 marks)

*Outcomes Assessed:* MEX12-6

*Targeted Performance Bands:* E2

Criteria	Marks
• correct solution	3
• evaluates period	2
• evaluates $n$	1

*Sample Answer:*

The particle is moving in simple harmonic motion, so  $a = -4^2x \therefore n = 4$ .

So period  $= \frac{2\pi}{4} = \frac{\pi}{2}$ . The particle returns to the origin in half a period, hence in  $\frac{\pi}{4}$  s.

Question 11(e) (3 marks)

*Outcomes Assessed:* MEX12-1, MEX12-4

*Targeted Performance Bands:* E2-E3

Criteria	Marks
• equates LHS and RHS to complete the proof	3
• expands LHS or RHS correctly	2
• correctly sets up real and imaginary parts of both numbers and substitutes	1

*Sample Answer:*

Let  $z = a + ib$  and  $w = c + id$  where  $a, b, c, d \in \mathbb{R}$ .

$$\begin{aligned} \text{LHS} &= \bar{z}w \\ &= \overline{(a+ib)(c+id)} \\ &= \overline{(ac-bd)+i(ad+bc)} \\ &= (ac-bd)-i(ad+bc) \end{aligned} \quad \begin{aligned} \text{RHS} &= \bar{z} \times \bar{w} \\ &= (a-ib)(c-id) \\ &= (ac-bd)-i(ad+bc) = \text{LHS} \end{aligned}$$

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

**Question 12 (15 marks)**

Question 12(a) (i) (2 marks)

*Outcomes Assessed: MEX12-3**Targeted Performance Bands: E2*

Criteria	Marks
• correct solution	2
• correctly substitutes in at least one dimension	1

*Sample Answer:*

$$\begin{pmatrix} 9 \\ k \\ 24 \end{pmatrix} = \begin{pmatrix} -1 \\ 2 \\ -6 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix}$$

$$\text{So, } 9 = -1 + 2\mu \Rightarrow \mu = 5.$$

$$\text{Further, } k = 2 - 3\mu \Rightarrow k = -13.$$

Question 12(a) (ii) (3 marks)

*Outcomes Assessed: MEX12-3**Targeted Performance Bands: E2*

Criteria	Marks
• correct solution	3
• finds $\lambda$ in terms of $\mu$ or similar progress	2
• correctly substitutes in at least one dimension	1

*Sample Answer:*

$$\text{From } j, 7 + \lambda = -1 + 2\mu \Rightarrow \lambda = -8 + 2\mu, \quad \text{and from } j, \quad -1 + 3\lambda = 2 - 3\mu$$

$$\text{Substituting the first into the second gives: } -1 + 3(-8 + 2\mu) = 2 - 3\mu$$

$$9\mu = 2 + 24 + 1$$

Which gives  $\mu = 3$  and  $\lambda = -2$ . Substituting in  $L_1$  or  $L_2$  gives the point  $(5, -7, 12)$ .

Question 12(b) (4 marks)

*Outcomes Assessed: MEX12-5**Targeted Performance Bands: E2*

Criteria	Marks
• correct solution	4
• correctly integrates and correctly substitutes limits	3
• correctly integrates or correctly substitutes limits	2
• devises appropriate substitution or similar progress	1

*Sample Answer:*If  $x = \sin \theta$  then  $dx = \cos \theta d\theta$ .When  $x = 0$ ,  $\theta = 0$ , when  $x = \frac{1}{\sqrt{2}}$ ,  $\theta = \frac{\pi}{4}$ .

$$\int_0^{\frac{1}{\sqrt{2}}} \frac{x^3}{\sqrt{1-x^2}} dx = \int_0^{\frac{\pi}{4}} \frac{\sin^3 \theta}{\sqrt{\cos^2 \theta}} \cos \theta d\theta$$

$$= \int_0^{\frac{\pi}{4}} \sin \theta (1 - \cos^2 \theta) d\theta$$

$$= [-\cos \theta + \frac{1}{3} \cos^3 \theta]_0^{\frac{\pi}{4}}$$

$$= \left( -\frac{1}{\sqrt{2}} + \frac{1}{3} \left( \frac{1}{\sqrt{2}} \right)^3 \right) - \left( -1 + \frac{1}{3} \right)$$

$$= \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{12} \right) + \frac{2}{3}$$

$$= \frac{8 - 5\sqrt{2}}{12}$$

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 12(c) (3 marks)

Outcomes Assessed: MEX12-4

Targeted Performance Bands: E2

Criteria	Marks
• correct solution	3
• finds square root of discriminant	2
• finds discriminant or attempts quadratic formula	1

*Sample Answer:*

Let  $a$  and  $b$  be real numbers such that  $(a+ib)^2 = \Delta = (7-i)^2 - 4(16+4i) = -16 - 30i$ .

Hence  $a^2 - b^2 = -16$  while  $2ab = -30$ . By inspection,  $a = 3$  and  $b = -5$ .

$$\text{Therefore } z = \frac{-(7-i) \pm (3-5i)}{2} = \frac{-4-4i}{2} \text{ or } \frac{-10+6i}{2} = -2-2i \text{ or } -5+3i.$$

Question 12(d) (3 marks)

Outcomes Assessed: MEX12-5

Targeted Performance Bands: E2-E3

Criteria	Marks
• correct solution	3
• finds correct partial fraction expression to integrate	2
• attempts partial fractions or similar progress	1

*Sample Answer:*

$$\int \frac{x^3-2}{x^3+x} dx = \int \left( \frac{x^3+x}{x^3+x} - \frac{x+2}{x(x^2+1)} \right) dx \quad \begin{aligned} &\text{Then } (A+B)x^2 + Cx + A \equiv x + 2 \\ &\text{Hence } A = 2, B = -2, C = 1. \end{aligned}$$

Now, if  $\frac{x+2}{x(x^2+1)} \equiv \frac{A}{x} + \frac{Bx+C}{x^2+1}$

$$\begin{aligned} \text{And further, } \int \frac{x^3-2}{x^3+x} dx &= \int \left( 1 - \frac{2}{x} + \frac{2x-1}{x^2+1} \right) dx \\ &= x - 2 \ln|x| + \int \frac{2x}{x^2+1} dx - \int \frac{1}{x^2+1} dx \\ &= x - 2 \ln|x| + \ln|x^2+1| - \tan^{-1}x + c \end{aligned}$$

**Question 13 (15 marks)**

Question 13(a) (2 marks)

Outcomes Assessed: MEX12-2

Targeted Performance Bands: E2-E3

Criteria	Marks
• completed proof	2
• significant progress towards proof	1

*Sample Answer:*

$$\begin{aligned} a, b > 0 \Rightarrow \left( \sqrt{ab} - \frac{1}{\sqrt{ab}} \right)^2 \geq 0 &\Rightarrow ab + 2 + \frac{1}{ab} \geq 4 \\ \Rightarrow ab - 2 + \frac{1}{ab} \geq 0 &\Rightarrow (a + \frac{1}{b})(b + \frac{1}{a}) \geq 4 \end{aligned}$$

*Disclaimer*

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSEA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather, it is intended to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 13(b) (i) (2 marks)

Outcomes Assessed: MEX12-4

Targeted Performance Bands: E2

Criteria	Marks
• correct solution	2
• evaluates $b$ or $c$ or work of equivalent merit	1

Sample Answer:

$$f(x) = (x+3)(x-2)(x^2 + bx + c)$$

$$f(0) = -6 = 3 \times -2 \times c \Rightarrow c = 1$$

$$f(1) = -4 = 4 \times -1 \times (1+b+1) \Rightarrow b+2=1 \Rightarrow b=-1$$

Now the two non-real roots of the equation are given by:

$$x^2 - x + 1 = 0$$

$$x^2 - x + \frac{1}{4} = -\frac{3}{4}$$

$$\left| \begin{aligned} & \left( x - \frac{1}{2} \right)^2 = \left( i \frac{\sqrt{3}}{2} \right)^2 \\ & x = \frac{1}{2} \pm \frac{\sqrt{3}}{2}i \end{aligned} \right.$$

Question 13(b) (ii) (2 marks)

Outcomes Assessed: MEX12-4

Targeted Performance Bands: E2

Criteria	Marks
• correct solution	2
• one correct root or both correct moduluses or both correct arguments	1

Sample Answer:

$$x = 1 \left( \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right) \text{ or } 1 \left( \cos \left( -\frac{\pi}{3} \right) + i \sin \left( -\frac{\pi}{3} \right) \right)$$


Question 13(b) (iii) (2 marks)

Outcomes Assessed: MEX12-4

Targeted Performance Bands: E2-E3

Criteria	Marks
• correct solution	2
• at least two correct points	1

Sample Answer:



#### Disclaimer

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 13(b) (iv) (1 mark)

*Outcomes Assessed:* MEX12-4

*Targeted Performance Bands:* E1

Criteria	Mark
• names the kite	1

*Sample Answer:*

Two pairs of adjacent sides are equal, hence they form a kite.

Question 13(c) (i) (1 mark)

*Outcomes Assessed:* MEX12-3

*Targeted Performance Bands:* E2

Criteria	Mark
• correct solution	1

*Sample Answer:*

$$\text{LHS} = \left| \begin{pmatrix} 5 \\ -10 \\ 3 \end{pmatrix} - \begin{pmatrix} 3 \\ -12 \\ 4 \end{pmatrix} \right| = \sqrt{2^2 + 2^2 + (-1)^2} = 3 = \text{RHS}$$

Question 13(c) (ii) (2 marks)

*Outcomes Assessed:* MEX12-3, MEX12-7

*Targeted Performance Bands:* E3

Criteria	Marks
• correct solution	2
• some progress towards correct solution	1

*Sample Answer:*

Centre of sphere is  $(3, -12, 4)$ , and the point on the sphere farthest from the origin would be 3 units (one radius) farther along the vector  $\vec{OC}$ .

Now,  $|\vec{OC}| = \sqrt{3^2 + (-12)^2 + 4^2} = 13$ , so the point  $P$  on the surface of the sphere farthest

from  $O$  will be given by:  $\vec{OP} = \frac{13+3}{13} \begin{pmatrix} 3 \\ -12 \\ 4 \end{pmatrix}$ . Hence the point  $P = \left( \frac{48}{13}, -\frac{192}{13}, \frac{64}{13} \right)$ .

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of the CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

**Question 13(d) (3 marks)****Outcomes Assessed:** MEX12-2, MEX12-8**Targeted Performance Bands:** E3

Criteria	Marks
• Completes the proof correctly	3
• Substitutes the correct expression for $n = k$ and $n = k + 1$	2
• Demonstrates the result true for $n = 7$	1

**Sample Answer:****RTP:**  $3n^2 - 3n \leq 2^n - 1$  for  $n \geq 7$ **PROOF:** if  $n = 7$ , LHS =  $21 \times 6 = 126$ RHS =  $2^7 - 1 = 127 \geq \text{LHS}$ . Hence the result is true for  $n = 7$ .Let's assume the result is true for some integer  $k$ .That is, assume  $3k(k-1) \leq 2^k - 1$  and then attempt to prove that  $3(k+1)k \leq 2^{k+1} - 1$ .So, IF  $3k(k-1) \leq 2^k - 1$ THEN  $6k(k-1) \leq 2^{k+1} - 2$ THEN  $6k^2 - 6k + 1 \leq 2^{k+1} - 1$ THEN  $3k^2 + 3k + (3k^2 - 9k + 1) \leq 2^{k+1} - 1$ Now to arrive at the result, we now need  $3k^2 - 9k + 1 \geq 0$ , which is true for  $k \geq \frac{9+\sqrt{81-12}}{6}$ .Further  $\frac{9+\sqrt{69}}{6} \approx 2.9$ , so for the values of  $k \geq 7$  in this result,  $3k^2 - 9k + 1 > 0$ .Putting this together,  $3(k+1)k < 3(k+1)k + (3k^2 - 9k + 1) \leq 2^{k+1} - 1$ Therefore,  $3(k+1)^2 - 3(k+1) \leq 2^{k+1} - 1$ .Hence if the result is true for  $n = k$ , the result will be true for  $n = k + 1$ .By the principle of Mathematical Induction, the result is true for  $n \geq 7$ .**Question 14 (15 marks)**

Question 14(a) (i) (2 marks)

**Outcomes Assessed:** MEX12-4**Targeted Performance Bands:** E2-E3

Criteria	Marks
• correct solution	2
• expands correctly or recognises sum of roots = 0 or equivalent merit	1

**Sample Answer:**

$$\begin{aligned} \text{LHS} &= (z + z^{-1})^2 + (z + z^{-1}) - 1 \\ &= \frac{1}{z^2} (z^4 + 2z^2 + 1 + z^3 + z - z^2) \\ &= \frac{1}{z^2} (z^4 + z^3 + z^2 + z + 1) \end{aligned}$$

Now given  $z^5 = 1$   
 $(z - 1)(z^4 + z^3 + z^2 + z + 1) = 0$   
and since  $z \neq 1, z^4 + z^3 + z^2 + z + 1 = 0$   
 $\text{LHS} = 0 = \text{RHS}$

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of the NSW Education Standards Authority to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 14(a) (ii) (1 mark)

*Outcomes Assessed:* MEX12-4

*Targeted Performance Bands:* E2

Criteria	Mark
• correct solution	1

*Sample Answer:*

$$\text{RHS} = \frac{1}{2}(\cos \theta + i \sin \theta + \cos(-\theta) + i \sin(-\theta)), \text{ and since cos is even fn and sin is odd}$$
$$= \frac{1}{2}(\cos \theta + i \sin \theta + \cos(\theta) - i \sin(\theta)) = \frac{1}{2} \times 2 \cos \theta = \text{LHS}$$

Question 14(a) (iii) (2 marks)

*Outcomes Assessed:* MEX12-4

*Targeted Performance Bands:* E3

Criteria	Marks
• correct solution	2
• some progress towards correct solution	1

*Sample Answer:*

$z$  is a non-real fifth root of unity so  
 $z = \text{cis } \theta$  where  $\theta = \pm \frac{2\pi}{5}$  or  $\pm \frac{4\pi}{5}$

From (i) and (ii):

$$(2 \cos \theta)^2 + (2 \cos \theta) - 1 = 0$$
$$4 \cos^2 \theta + 2 \cos \theta - 1 = 0$$

$$\cos \theta = \frac{-2 \pm \sqrt{4+16}}{8} = \frac{-1 \pm \sqrt{5}}{4}$$

Now,  $\cos \frac{2\pi}{5} > 0$ .

$$\text{Therefore } \cos \frac{2\pi}{5} = \frac{-1 + \sqrt{5}}{4}.$$

Question 14(a) (iv) (3 marks)

*Outcomes Assessed:* MEX12-4

*Targeted Performance Bands:* E3-E4

Criteria	Marks
• correct solution	3
• evaluates $ z_2 - z_1 $ and $ z_3 - z_1 $ correctly	2
• evaluates $ z_2 - z_1 $ or $ z_3 - z_1 $ correctly or significant progress towards both	1

*Sample Answer:*

From (iii),  $\cos \frac{2\pi}{5} = \frac{-1 + \sqrt{5}}{4}$ . Whereas  $\cos \frac{4\pi}{5} < 0$  and hence  $\cos \frac{4\pi}{5} = \frac{-1 - \sqrt{5}}{4}$ .

$$|z_3 - z_1| = |\text{cis } \frac{4\pi}{5} - 1| \quad \text{Also } |z_2 - z_1| = |\text{cis } \frac{2\pi}{5} - 1|$$
$$= \sqrt{(\cos \frac{4\pi}{5} - 1)^2 + (\sin \frac{4\pi}{5})^2} \quad = \sqrt{2 - 2 \cos \frac{2\pi}{5}}$$
$$= \sqrt{2 - 2 \cos \frac{4\pi}{5}} = \sqrt{2 + \frac{1}{2} + \frac{\sqrt{5}}{2}} \quad = \sqrt{2 + \frac{1}{2} - \frac{\sqrt{5}}{2}}$$
$$= \frac{\sqrt{5+\sqrt{5}}}{\sqrt{2}} \quad = \frac{\sqrt{5-\sqrt{5}}}{\sqrt{2}}$$

$$\text{Hence } \left| \frac{z_3 - z_1}{z_2 - z_1} \right| = \frac{\sqrt{5+\sqrt{5}}}{\sqrt{5-\sqrt{5}}} = \sqrt{\frac{\sqrt{5}+1}{\sqrt{5}-1}} = \sqrt{\frac{(\sqrt{5}+1)^2}{4}} = \frac{\sqrt{5}+1}{2}, \text{ as required.}$$

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific Trial HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 14(b) (i) (1 mark)

*Outcomes Assessed:* MEX12-3

*Targeted Performance Bands:* E2

Criteria	Mark
• correct solution	1

*Sample Answer:*

$$\text{When } t = 3, \underline{r} = -\frac{1}{\sqrt{2}}\underline{i} + 0\underline{j} + \frac{1}{\sqrt{2}}\underline{k}$$

Question 14(b) (ii) (3 marks)

*Outcomes Assessed:* MEX12-3

*Targeted Performance Bands:* E3

Criteria	Marks
• correct solution, NB there are many correct vector equations for the line	3
• evaluates direction vector $\underline{v}(3)$ or a scalar multiple of the direction vector	2
• differentiates correctly to find $\underline{v}(t)$	1

*Sample Answer:*

$$\underline{v}(t) = \frac{\pi}{4} \left[ -\sin\left(\frac{\pi}{4}t\right)\underline{i} + \left(-\sin\left(\frac{\pi}{4}t\right) + \cos\left(\frac{\pi}{4}t\right)\right)\underline{j} + \cos\left(\frac{\pi}{4}t\right)\underline{k} \right]$$

$$\underline{v}(3) = \frac{\pi}{4} \left[ -\frac{1}{\sqrt{2}}\underline{i} + \left(-\frac{1}{\sqrt{2}} + -\frac{1}{\sqrt{2}}\right)\underline{j} + \frac{1}{\sqrt{2}}\underline{k} \right] = \frac{\pi}{4\sqrt{2}} \begin{pmatrix} -1 \\ -2 \\ 1 \end{pmatrix}$$

$$\text{Therefore the tangent will be given by: } \underline{r} = \begin{pmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ -2 \\ 1 \end{pmatrix}, \text{ for } \lambda \in \mathbb{R}.$$

Question 14(c) (3 marks)

*Outcomes Assessed:* MEX12-5

*Targeted Performance Bands:* E2-E3

Criteria	Marks
• correct solution	3
• correctly completes two substitutions or one substitution with correct limits	2
• correctly completes at least one relevant substitution with or without limits	1

*Sample Answer:*

Attempting with a single substitution:

$$\text{Let } u^2 = 1 + \sqrt{x}$$

$$2udu = \frac{1}{2\sqrt{x}}dx$$

$$4u\sqrt{x}du = dx$$

$$4u(u^2 - 1) du = dx$$

$$\begin{aligned} \int_0^9 \frac{1}{\sqrt{1+\sqrt{x}}} dx &= \int_1^2 \frac{1}{u} \times 4u(u^2 - 1) du \\ &= 4 \int_1^2 (u^2 - 1) du \\ &= 4 \left[ \frac{u^3}{3} - u \right]_1^2 \\ &= 4 \left[ \left( \frac{8}{3} - 2 \right) - \left( \frac{1}{3} - 1 \right) \right] = \frac{16}{3} \end{aligned}$$

#### Disclaimer

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking guidelines for individual CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

### Question 15 (15 marks)

Question 15(a) (4 marks)

*Outcomes Assessed: MEX12-5*

*Targeted Performance Bands: E3-E4*

Criteria	Marks
• correct solution	4
• two correct applications of integration by parts to arrive at $\int f(x) dx$	3
• one correct application of integration by parts	2
• correct x-intercepts	1

*Sample Answer:*

$f(x) = e^{-x} \cos x$ ,  $x \in [0, \frac{3\pi}{2}]$ . Now, the x-intercepts at  $e^{-x} \cos x = 0$ , hence at  $x = \frac{\pi}{2}$  then  $\frac{3\pi}{2}$ , and since  $e^{-x} > 0$ , the sign of the cosine function will be the sign of the function, so the curve is above the x-axis for  $0 < x < \frac{\pi}{2}$  and below for  $\frac{\pi}{2} < x < \frac{3\pi}{2}$ .

So we're looking to find the ratio of integrals  $\int_0^{\frac{\pi}{2}} f(x) dx : -\int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} f(x) dx$ .

$$\int e^{-x} \cos x dx \quad dv = e^{-x} dx \quad \text{and} \quad u = \cos x \\ v = -e^{-x} \quad du = -\sin x dx$$

$$\text{Hence, } \int e^{-x} \cos x dx = [-e^{-x} \cos x] - \int -e^{-x} \times -\sin x dx$$

$$= -e^{-x} \cos x - \int e^{-x} \times \sin x dx \\ dv = e^{-x} dx \quad \text{and} \quad u = \sin x \\ v = -e^{-x} \quad du = \cos x dx$$

$$\text{Hence, } \int e^{-x} \cos x dx = -e^{-x} \cos x - \left( -e^{-x} \sin x - \int -e^{-x} \times \cos x dx \right)$$

$$\text{So, } 2 \int e^{-x} \cos x dx = -e^{-x} \cos x + e^{-x} \sin x + c_1$$

$$\text{Finally, } \int e^{-x} \cos x dx = \frac{1}{2} e^{-x} (\sin x - \cos x) + c$$

Ratio of areas will be given by

$$\begin{aligned} \int_0^{\frac{\pi}{2}} f(x) dx : -\int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} f(x) dx &= \left[ \frac{1}{2} e^{-x} (\sin x - \cos x) \right]_0^{\frac{\pi}{2}} : -\left[ \frac{1}{2} e^{-x} (\sin x - \cos x) \right]_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \\ &= \left[ e^{-\frac{\pi}{2}} (1-0) - e^0 (0-1) \right] : -\left[ e^{-\frac{3\pi}{2}} (-1-0) - e^{-\frac{\pi}{2}} (1-0) \right] \\ &= \left( e^{-\frac{\pi}{2}} + 1 \right) : \left( e^{-\frac{3\pi}{2}} + e^{-\frac{\pi}{2}} \right), \text{ now multiply both terms by } e^{\frac{3\pi}{2}} \\ &= \left( e^{\pi} + e^{\frac{3\pi}{2}} \right) : (1+e^{\pi}) = \frac{e^{\pi} (e^{\frac{\pi}{2}} + 1)}{e^{\pi} + 1}, \text{ as required.} \end{aligned}$$

#### Disclaimer

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

**Question 15(b) (4 marks)**

**Outcomes Assessed:** MEX12-6, MEX12-7

**Targeted Performance Bands:** E3-E4

Criteria	Marks
• correct solution	4
• uses initial conditions to find $x$ as a function of $v$	3
• integrates to find an expression for $x$ wrt $v$ , even without constant	2
• correct $\dot{v}$	1

**Sample Answer:**

Gravity acts in the direction of the slope with magnitude  $10\cos 60^\circ = 5 \text{ ms}^{-2}$ .

Also since  $F = ma$ , the acceleration due to friction is  $\frac{v}{20}$ .

$$\begin{aligned}\dot{v} &= -5 - \frac{v}{20} \\ v \frac{dv}{dx} &= -\frac{100 + v}{20} \\ \frac{dx}{dv} &= -\frac{20v}{100 + v} \\ x &= -\int \frac{20v}{100 + v} dv \\ -x &= 20 \int \left( \frac{v+100}{v+100} - \frac{100}{v+100} \right) dv \\ -x &= 20(v - 100 \ln|v+100|) + c\end{aligned}$$

When  $x = 0, v = 144 \text{ km/h} = 40 \text{ m/s}$ , hence

$$\begin{aligned}0 &= 20(40 - 100 \ln 140) + c \\ c &= 20(100 \ln 140 - 40) \\ -x &= 20(v - 100 \ln|v+100| + 100 \ln 140 - 40) \\ x &= 20 \left( 40 - v + 100 \ln \left| \frac{v+100}{140} \right| \right)\end{aligned}$$

$$\text{When } v = 0, x = 20 \left( 40 + 100 \ln \frac{5}{7} \right) \approx 127 \text{ metres.}$$

**Question 15(c) (i) (2 marks)**

**Outcomes Assessed:** MEX12-7

**Targeted Performance Bands:** E2-E3

Criteria	Marks
• correct solution	2
• significant progress towards correct solution	1

**Sample Answer:**

From the Reference Sheet:  $\sin A \sin B = \frac{1}{2} [\cos(A-B) - \cos(A+B)]$

$$\begin{aligned}\text{So, RHS} &= 2 \sin((n-1)x) \sin x \\ &= \cos((n-1)x - x) - \cos(((n-1)x + x)) \\ &= \cos((n-2)x) - \cos(nx) = \text{LHS}\end{aligned}$$

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific Trial HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 15(c) (ii) (2 marks)

*Outcomes Assessed:* MEX12-7

*Targeted Performance Bands:* E3-E4

Criteria	Marks
• correct solution	2
• uses identity in (i) to find simple sine integrand or similar progress	1

*Sample Answer:* LHS =  $I_n - I_{n-2}$

$$\begin{aligned}
 &= \int \frac{\cos nx}{\sin x} dx - \int \frac{\cos(n-2)x}{\sin x} dx \\
 &= - \int \frac{\cos((n-2)x) - \cos(nx)}{\sin x} dx, \quad \text{and from (i):} \\
 &= 2 \int -\sin((n-1)x) dx \\
 &= 2 \left[ \frac{1}{n-1} \times \cos((n-1)x) \right] + c \\
 &= \text{RHS}
 \end{aligned}$$

Question 15(c) (iii) (3 marks)

*Outcomes Assessed:* MEX12-5

*Targeted Performance Bands:* E3-E4

Criteria	Marks
• correct solution	3
• uses (ii) to find an expression in $x$ for the integral	2
• expresses integral in terms of $I_n$	1

*Sample Answer:*

$$\begin{aligned}
 \int_0^{\frac{\pi}{3}} \frac{\cos 2x - \cos 6x}{\sin x} dx &= \left[ I_2 - I_6 \right]_0^{\frac{\pi}{3}} \\
 &= \left[ -(I_6 - I_4) - (I_4 - I_2) \right]_0^{\frac{\pi}{3}}, \quad \text{and from (ii):} \\
 &= - \left[ \frac{2\cos(5x)}{5} + \frac{2\cos(3x)}{3} \right]_0^{\frac{\pi}{3}} \\
 &= - \left[ \frac{2}{5} \left( \frac{1}{2} - 1 \right) + \frac{2}{3} (-1 - 1) \right] \\
 &= \frac{1}{5} + \frac{4}{3} = \frac{23}{15}
 \end{aligned}$$

#### Disclaimer

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the intention is to provide a guide to assist teachers to better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific "Trial" HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

## Question 16 (15 marks)

Question 16(a) (3 marks)


*Outcomes Assessed: MEX12-7, MEX12-8*

*Targeted Performance Bands: E3-E4*


Criteria	Marks
• Completes the proof correctly	3
• Correctly illustrates the figure for $n = k$	2
• Demonstrates the result true for $n = 1$	1

*Sample Answer:*

In the case of  $n = 1$ , the triangle of side length  $2^1 = 2$  has one triangle left over, as shaded.



Not to contribute to the proof, but the case of  $n = 2$  is pictured, with total side length of  $2^2 = 4$  and 1 missing triangle, as shaded.




Hence the result is true for  $n = 1$ .

Assume the result is true for some integer  $n = k$ , which means it is possible to create a triangle of side length  $2^k$  with one triangle of side length 1 missing in the corner.

Now take four of these triangles of side length  $2^k$  and arrange them as in the diagram on the right. Observe that one of the tiles can fill the gap left by three of the missing corner triangles on the right.

Thus an equilateral triangle with sides  $2^k + 2^k = 2^{k+1}$  with one triangle of side length 1 missing in the corner.



Hence the result is true by the Principle of Mathematical Induction.

### Disclaimer

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific Trial HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 16(b) (i) (3 marks)

Outcomes Assessed: MEXI2-3

Targeted Performance Bands: E3-E4

Criteria	Marks
• correct solution	3
• evaluates $t$ for $Q$	2
• substitutes $ OQ  = 25$ into $ \underline{r} $	1

Sample Answer:

$$\begin{aligned} |OQ|^2 &= 25^2 = 49 \sin^2\left(\frac{\pi t}{32}\right) + 49 \cos^2\left(\frac{\pi t}{32}\right) + \left(\frac{t}{15}\right)^2 \\ 625 &= 49 + \frac{t^2}{15^2} \\ t^2 &= 576 \times 15^2 \\ t &= 24 \times 15 = 360 \text{ seconds} \end{aligned}$$

So, if  $t = 360$ , then  $\frac{\pi t}{32} = \frac{45\pi}{4}$

$$\begin{aligned} \underline{r}(t) &= 7 \sin \frac{45\pi}{4} \underline{i} + 7 \cos \frac{45\pi}{4} \underline{j} + \frac{360}{15} \underline{k} \\ &= 7 \sin \frac{-3\pi}{4} \underline{i} + 7 \cos \frac{-3\pi}{4} \underline{j} + 24 \underline{k} \end{aligned}$$

$$\text{Therefore } Q = \left( \frac{-7}{\sqrt{2}}, \frac{-7}{\sqrt{2}}, 24 \right)$$

Question 16(b) (ii) (2 marks)

Outcomes Assessed: MEXI2-3

Targeted Performance Bands: E3-E4

Criteria	Marks
• correct solution	2
• attempting to solve for $x = 7$ or work of similar merit	1

Sample Answer:

The line  $x = 7$  is crossed when the  $i$  component is equal to 7.

$$\begin{aligned} 7 \sin \frac{\pi t}{32} &= 7 \\ \frac{\pi t}{32} &= \frac{\pi}{2}, \frac{5\pi}{2}, \frac{9\pi}{2}, \dots \\ t &= 16, 80, 144, 208, 272, 336, 400, \dots \end{aligned}$$

Hence line  $x = 7$  is crossed 6 times in the 360 seconds it takes the ant to crawl to  $Q$ .

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific "Trial" HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 16(b) (iii) (2 marks)

**Outcomes Assessed:** MEXI2-3

**Targeted Performance Bands:** E4

Criteria	Marks
• correct solution	2
• significant progress towards correct solution, for example finding the arc length $QP'$ , or conceiving of the ant's path as a rectangle	1


**Sample Answer:**

As illustrated in the diagrams on the right, the most direct path from  $Q$  to  $P$  can be found by considering the cylindrical column as a rectangle with  $Q$  and  $P$  as opposite vertices.

Let the point  $P'$  be at the same height of  $Q$  and directly above  $P$ . Hence  $P' = (0, 7, 24)$ , and the points  $Q$  and  $P'$  lie on the plane  $z = 24$ , as in the first diagram on the right.

The arc length from  $Q$  to  $P'$  will be  $7 \times \frac{3\pi}{4}$ . This is also represented in the second diagram, and this gives:

$$\text{The ant's path } QP = \sqrt{24^2 + \left(\frac{21\pi}{4}\right)^2} \approx 29.1 \text{ cm}$$



**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements. It is established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.

Question 16(c) (5 marks)

*Outcomes Assessed:* MEX12-6, MEX12-7

*Targeted Performance Bands:* E3-E4

Criteria	Marks
• correct solution	5
• forming an equation with only $v_0$	4
• one correct integration on the downward journey	3
• correct substitution of initial conditions	2
• one correct integration on the upward journey	1

*Sample Answer:*

Upwards is positive, and  
when  $x = 0, v = v_0$ .

$$\begin{aligned}\dot{v} &= v \frac{dv}{dx} = -10 - 0.01v^2 \\ \frac{dv}{dx} &= -0.01 \left( \frac{1000}{v} + v \right) \\ \frac{dx}{dv} &= -100 \left( \frac{v}{1000+v^2} \right) \\ x &= -50 \int \frac{2v}{1000+v^2} dv \\ &= -50 \ln |1000+v^2| + c_1\end{aligned}$$

when  $x = 0, v = v_0$ , so

$$\begin{aligned}0 &= -50 \ln |1000+v_0^2| + c_1 \\ x &= 50 \ln \left| \frac{1000+v_0^2}{1000+v^2} \right|\end{aligned}$$

Reaches max height,  $x = H$  when  $v = 0$

$$H = 50 \ln \left| \frac{1000+v_0^2}{1000} \right|$$

On the downward journey  
let downwards be positive, and  
when  $x = 0, v = 0$ .

$$\begin{aligned}\dot{v} &= v \frac{dv}{dx} = 10 - 0.01v^2 \\ \frac{dv}{dx} &= 0.01 \left( \frac{1000}{v} - v \right) \\ \frac{dx}{dv} &= 100 \left( \frac{v}{1000-v^2} \right) \\ x &= -50 \int \frac{-2v}{1000-v^2} dv \\ &= -50 \ln |1000-v^2| + c_2\end{aligned}$$

when  $x = 0, v = 0$ , so

$$\begin{aligned}0 &= -50 \ln |1000| + c_2 \\ x &= 50 \ln \left| \frac{1000}{1000-v^2} \right|\end{aligned}$$

On impact,  $x = H$ , and crucially  $v = \frac{v_0}{7}$

$$50 \ln \left| \frac{1000+v_0^2}{1000} \right| = 50 \ln \left| \frac{1000}{1000-(v_0/7)^2} \right|$$

$$\begin{aligned}(1000+v_0^2) \left( 1000 - \left( \frac{v_0}{7} \right)^2 \right) &= 1000^2 \\ 1000v_0^2 - \frac{1000}{49}v_0^2 - \frac{1}{49}v_0^4 &= 0 \\ v_0^2 (v_0^2 + 1000 - 49000) &= 0\end{aligned}$$

Hence  $v_0 = 0$  or  $\sqrt{48000} \approx 219.1 \text{ ms}^{-1}$ .

**Disclaimer**

The information contained in this document is intended for the professional assistance of only teaching staff. It does not constitute advice to students. Further it is not the intention of CSSA to provide specific marking outcomes for all possible CSSA Trial HSC Examination answers. Rather the purpose is to provide teachers with information so that they can better explore, understand and apply HSC marking requirements, as established by the NSW Education Standards Authority. No guarantee or warranty is made or implied with respect to the application or use of CSSA Trial HSC Examination Marking Guidelines in relation to any specific 'Trial' HSC Examination question or answer. The CSSA assumes no liability or responsibility for the accuracy, completeness or usefulness of any Marking Guidelines provided for the CSSA Trial HSC Examination papers.