



coastal soil nonfarmed	irrigated sand mulched soil
92	30
irrigated tephra mulched soil	natural soil nonfarmed I
80	36
natural soil nonfarmed II	natural soil nonfarmed III
31	53
natural soil nonterraced	rainfed sand mulched soil
50	30
rainfed tephra mulched soil	runoff capture plots
80	83
soil terraced	
50	


Barplot of CLASS



# CLASS according to location and Agricultural System



# CLASS according to location and Soil Farming



[illegible]

This scatter plot displays the spatial distribution of various land use classes across a grid defined by XUTM (horizontal axis, ranging from 610,000 to 650,000) and YUTM (vertical axis, ranging from 6,000 to 14,000). The data points are categorized by 'Soil Irrigation' status and 'CLASS'.

**Soil Irrigation Legend:**

- Irrigated (Red circle)
- No Irrigation non-farmed (Green circle)
- Rainfed (Blue circle)
- Runoff (Purple circle)

**CLASS Legend:**

- coastal soil nonf (Black circle)
- irrigated sand m (Black triangle)
- irrigated tephra (Black square)
- natural soil nonf (Black plus)
- natural soil nonf (Black cross)
- natural soil nonf (Black asterisk)
- natural soil nonf (Black dot)
- rainfed sand mu (Black diamond)
- rainfed tephra m (Black square)
- runoff capture p (Black circle)
- soil terraced (Black cross)

The plot shows a high density of 'Irrigated' (red) and 'No Irrigation non-farmed' (green) points in the central and right-hand areas, particularly between YUTM 10,000 and 12,000. 'Rainfed' (blue) and 'Runoff' (purple) points are less frequent and appear more scattered. The 'CLASS' legend indicates that the symbols used in the plot correspond to specific land use categories, such as 'coastal soil nonf', 'irrigated sand m', 'irrigated tephra', etc.

