

Papers to read/cite for Str2LeadTrans Paper

February 12, 2021

1

In [Luque and Ebert, 2010],

- (a) They write, “As discussed in the previous section, the densities of charged particles increase like N so an exponentially increasing number of electrons is liberated around the head. These electrons drift upwards, where the ion density is basically frozen (neglecting small variations due to attachment). **Hence at some point the electron density surpasses the ion density and creates a net negative charge that is responsible for the increase in the electric field, eventually to values above E_k ; then a second ionization wave sets in.**” The question, I pose is, does a second-ionization wave occur if heating is neglected?
- (b) They also write, “Other observations in high-speed sprite imaging also suggest a negative charging of the streamer channel. The first is the emergence of negative (upward-propagating) streamers, always reported to occur from a previous channel and some milliseconds after the passage of a positive streamer head. In some observations [Stanley et al., 1999; Cummer et al., 2006; Stenbaek-Nielsen and McHarg, 2008] the emergence of negative streamers coincides with the lower edge of the trailing emissions.”

From Liu and Pasko [2004]:

- (a) “These processes, however, are known to be important for the dynamics of long streamers developing in point-to-plane discharge gaps in low electric fields ($<E_k$) Morrow and Lowke [1997].”

From Shi et al. [2016]:

- (a) “On the other hand, the exponential growth of a streamer may be a property particularly important for lightning initiation, since the current flowing in the channel also exponentially increase Liu [2010], potentially accelerating the heating processes in the discharge channel. It is thus necessary to investigate whether the streamer initiated from a hydrometeor will exponentially grow over a long distance as it was assumed in Liu et al. [2012] and Sadighi [2015] and to obtain its propagation characteristics.”

References

- N. Y. Liu. Model of sprite luminous trail caused by increasing streamer current. *Geophys. Res. Lett.*, 37:L04102, February 2010. 10.1029/2009GL042214.
- N. Y. Liu and V. P. Pasko. Interaction of sprite streamers with the lower ionosphere. *Abstracts*, page 245, 2004. National Radio Science Meeting, Boulder, Colorado, January 5-8.
- N. Y. Liu, J. R. Dwyer, and H. K. Rassoul. Effects of pressure and humidity on positive corona inception from thundercloud hydrometeors. *J. Atmos. Solar Terr. Phys.*, 80:179–186, May 2012. 10.1016/j.jastp.2012.01.012.
- A. Luque and U. Ebert. Sprites in varying air density: Charge conservation, glowing negative trails and changing velocity. *Geophys. Res. Lett.*, 37:L06806, Mar 2010. 10.1029/2009gl041982.
- R. Morrow and J. J. Lowke. Streamer propagation in air. *J. Phys. D: Appl. Phys.*, 30:614–627, 1997.
- Samaneh Sadighi. Initiation of streamers from thundercloud hydrometeors and implications to lightning initiation. 2015. Ph.D. thesis, Florida Tech, Melbourne, FL.
- F. Shi, N. Y. Liu, and H. K. Rassoul. Properties of relatively long streamers initiated from an isolated hydrometeor. *J. Geophys. Res.*, 121(12):7284–7295, 2016. 10.1002/2015JD024580.