

Residual Spatial Autocorrelation Test with Moran's I

Outline:

- What is spatial autocorrelation
- Definition of the link matrix
- Why autocorrelation in regression residuals?
- Motivating example: Italian Fertility
- General structure of test statistic and its distribution
- Simplifications under the assumption of spatial independence
- Maximum likelihood estimation

What is Spatial Autocorrelation

- The concept of an ***internal spatial relationship*** (order) is trickier than in the time-series situation, where the order comes naturally (the future depends on the past)
- In spatial analysis, the relationships among the observations are [a] ***multidirectional***, [b] ***multilateral*** and [c] ***not equally spaced***.


There are also more observations at the ***edge of the study area*** and factors like ***spatial extend of the regions*** and ***underlying densities*** become important.

- Geographical theories provide many ***concepts of spatial relationships*** within a single variable. For instance:
 - simple distances between points or representative points of areas
 - **neighborhood** relationships between areas (rook's or queen's specification in square tessellations, higher order neighborhood relationship of regions several neighbors apart)
 - traffic flows or migration patterns
 - spatial hierarchies (hub and spokes)
 - diffusion processes etc.

The Spatial Link Matrix

- The spatial **connectivity matrix** operationalizes the underlying structure of the potential spatial relationships among the observations
- For potential distance relationships we have the distance matrix (known from road atlases, perhaps using spherical distances)
- For potential neighborhood relationships we must use a binary spatial connectivity matrix

Example: Encoding a spatial tessellation as a binary connectivity matrix



Spatial tessellation of 9 hexagonal cells with underlying connectivity structure



symmetric

	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>	<i>g</i>	<i>h</i>	<i>i</i>
<i>a</i>	0	1	0	1	0	0	0	0	0
<i>b</i>	1	0	1	1	1	0	0	0	0
<i>c</i>	0	1	0	0	1	1	0	0	0
<i>d</i>	1	1	0	0	1	0	1	0	0
<i>e</i>	0	1	1	1	0	1	1	1	0
<i>f</i>	0	0	1	0	1	0	0	1	1
<i>g</i>	0	0	0	1	1	0	0	1	0
<i>h</i>	0	0	0	0	1	1	1	0	1
<i>i</i>	0	0	0	0	0	1	0	1	0

Binary 9×9 spatial connectivity matrix

B

- An element $b_{ij} = 1$ denotes that the tiles i and j are **adjacent** and an element $b_{ij} = 0$ signifies that the tiles i and j are not common neighbors
- The spatial connectivity matrix **B** is **symmetric**
- A tile is not connected to itself. Thus all **diagonal elements are zero**
- For study areas with a **large number n** of individual regions the generation of the connectivity matrix **B** (or distance matrix) must be left to a GIS program. The connectivity matrix has $n \times n$ elements
- Problems occur if we have island and holes in our study area. Usually machine generated connectivity matrices must be polished manually.
- Most of the elements are zero. There are efficient storage modes for sparse matrices. For **empirical** map patterns an area in the interior has on **average 6 neighbors**.

- The binary spatial link matrix \mathbf{B} is usually **coded**: For instance, if each row is divided by its row sum we get the **row-sum standardized** link matrix $\mathbf{V} = \underset{\text{row-sum}}{\text{diag}(\mathbf{B} \cdot \mathbf{1})}^{-1} \cdot \mathbf{B}$.

Each row of \mathbf{V} sums to one and the overall sum over all elements of \mathbf{V} is n .

- Other coding scheme are possible and they follow the generic equations:

$$\mathbf{V} = \frac{n}{\underbrace{\mathbf{1}^T \cdot \text{diag}(\mathbf{d}^q) \cdot \mathbf{B} \cdot \mathbf{1}}_{\text{sum of transformed component values}}} \cdot \text{diag}(\mathbf{d}^q) \cdot \mathbf{B} \text{ with } \mathbf{d} = \mathbf{B} \cdot \mathbf{1}$$

For the **row sum standardized coding scheme** the transformation parameter is $q = -1$.

- Spatial econometrics** prefers the row-sum standardized coding scheme and **denotes the standardized link matrix by \mathbf{W}** . In contrast, geo-statistics usually works with standardized interpoint distance matrices.

Why autocorrelation in regression residuals?

- Regression residuals \mathbf{e} capture the unexplained part of the regression model
- They are usually assumed to relate to population disturbances that are $\boldsymbol{\varepsilon} \sim N(0, \sigma^2 \cdot \mathbf{I})$
- Whereas the residual are distributed as $\mathbf{e} \sim N(0, \sigma^2 \cdot \mathbf{M})$ with $\mathbf{M} = \mathbf{I} - \mathbf{X} \cdot (\mathbf{X}^T \cdot \mathbf{X})^{-1} \cdot \mathbf{X}^T$ where \mathbf{X} is the matrix of independent variables including the intercept constant vector $\mathbf{1}$.
- (1) **Misspecification Rational**: if we are **missing relevant variables in \mathbf{X}** that exhibit a spatial pattern, their spatial pattern will **spill over into the regression residuals \mathbf{e}** .
- (2) **Spatial Process Rational**: The spatial object exhibit some ***spatial exchange relationships***, e.g.,
 - interaction flows,

caused by unknown variable which has a spatial pattern.

- competition effects or **price and customers**
- agglomerative advantages

then they will become spatially autocorrelated.

These exchange relationships cannot be captured by the regression matrix **X** but manifest in a **covariance** matrix $\Omega(\rho)$ with $\epsilon \sim N(0, \sigma^2 \cdot \Omega(\rho))$

where ρ is the **autocorrelation level** and measuring the strength of the spatial process. For

$$\rho = \begin{cases} > 0 & \text{positive autocorrelation} \\ 0 & \text{spatial independence} \\ < 0 & \text{negative autocorrelation} \end{cases}$$

Moran's I is tailored to work best for **Gaussian spatial processes** with

$$\Omega(\rho) = \begin{cases} \text{simultaneous autoregressive spatial process} \\ \text{conditional autoregressive spatial process} \\ \text{moving average spatial process} \end{cases}$$


All these processes dependent on the **coded** spatial link matrix **V** among the spatial objects, which reflects the spatial relationships and that is derived from the binary links **B**.

- (3) Spatial Aggregation Rational: If areal objects are split into parts and these split parts are merged with adjacent areal objects then these aggregated objects share parts of the information that they inherited from the split objects.

This induces spatial autocorrelation among adjacent spatial objects.
Implication: One needs to get the spatial scale of analysis right.

Motivating Example: **Fertility** in Italy


- Several spatial link matrices are conceivable:



- The spatial relationships can be captured in the spatial link matrix \mathbf{V}
- The map pattern of the observed dependent variable **Total Period Fertility** and the autocorrelation of the regression residuals around the mean is $e_i = y_i - \bar{y}$:

color not good. should use bipolar

Provincial Total Fertility Rate 1994



Global Moran's I for regression residuals

```
model: lm(formula = TOTFERTRAT ~ 1, data = prov.df)
weights: nb2listw(prov.nb, style = "S")
```

Moran I statistic **standard deviate** = **12.7804**, p-value < 2.2e-16

alternative hypothesis: greater

sample estimates:

Observed Moran's I	Expectation	Variance
0.853201213	-0.010638298	0.004568551

- The applied *regression model* to explain the Total Fertility Rate is

```
lm(formula = TOTFERTRAT ~ FEMMARAGE9 + DIVORCERAT + log(ILLITERRAT) +
    TELEPERFAM, data = prov.df)
```

Residuals:

Min	1Q	Median	3Q	Max
-0.19958	-0.05474	-0.01284	0.05272	0.42922

Coefficients:



	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	4.78139	0.48606	9.837	6.23e-16	***
FEMMARAGE9	-0.09647	0.02050	-4.706	9.11e-06	***
DIVORCERAT	-0.11839	0.05772	-2.051	0.0431	*

```

log(ILLITERAT)  0.03072      0.01707    1.799    0.0753 .
TELEPERFAM     -1.28499     0.18078   -7.108  2.69e-10 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


```

Residual standard error: 0.1047 on 90 degrees of freedom
 Multiple R-squared: 0.8051, Adjusted R-squared: 0.7965
 F-statistic: 92.96 on 4 and 90 DF, p-value: < 2.2e-16




outlier ?
would focus on it

- The map pattern of the residuals $e_i = y_i - \hat{y}_i$ and their autocorrelation level are



- Notes:
 - Accounting** for the **exogenous variables** has substantially reduced the autocorrelation level (\Rightarrow misspecification perspective).
 - The **expectation** and **variance** of Moran's I dependent on the regression matrix **X**.
 - Residuals are best mapped by a **bipolar** map theme.

- Most *extreme possible spatial map patterns* and their autocorrelation levels



General structure of the test statistic and its distribution

The observed value of Moran's I

- Moran's I is defined to measure the strength of spatial autocorrelation in regression residuals

$\mathbf{e} = \mathbf{M} \cdot \mathbf{y}$ with

$$\mathbf{e} = \begin{cases} \mathbf{y} - \mathbf{1} \cdot \bar{y} & \text{if } [\mathbf{I} - \mathbf{1} \cdot (\mathbf{1}^T \cdot \mathbf{1})^{-1} \cdot \mathbf{1}^T] \cdot \mathbf{y} \\ \mathbf{y} - \hat{\mathbf{y}} & \text{if } [\mathbf{I} - \mathbf{X} \cdot (\mathbf{X}^T \cdot \mathbf{X})^{-1} \cdot \mathbf{X}^T] \cdot \mathbf{y} \end{cases}$$

- Its observed value I^{obs} has the formal structure of a **ratio of quadratic forms** in the random variable \mathbf{y}

$$I^{obs} = \frac{\mathbf{y}^T \cdot \mathbf{M} \cdot \frac{1}{2} \cdot (\mathbf{V} + \mathbf{V}^T) \cdot \mathbf{M} \cdot \mathbf{y}}{\mathbf{y}^T \cdot \mathbf{M} \cdot \mathbf{y}} = \frac{\mathbf{e}^T \cdot \frac{1}{2} \cdot (\mathbf{V} + \mathbf{V}^T) \cdot \mathbf{e}}{\mathbf{e}^T \cdot \mathbf{e}}$$

- Besides the spatial autocorrelation pattern in the dependent **random variable** \mathbf{y} — through $\boldsymbol{\varepsilon} \sim N(0, \boldsymbol{\Omega})$ — the observed value of **Moran's I** also **depends on the exogenous variables \mathbf{X}** through the projection matrix \mathbf{M} and the **exogenous spatial link matrix \mathbf{V}** .
- The spatial link matrix \mathbf{V} is **symmetrized** by the transformation $\frac{1}{2} \cdot (\mathbf{V} + \mathbf{V}^T)$.
While the observed value I^{obs} is invariant under this transformation, the evaluation of its distribution requires a symmetric structure.
- The **largest and smallest possible values** of Moran's I can be determined by the largest eigenvalue λ_1 and smallest eigenvalue λ_n of the matrix $\mathbf{M} \cdot \frac{1}{2} \cdot (\mathbf{V} + \mathbf{V}^T) \cdot \mathbf{M}$. spatial filtering


Test under the assumption of spatial independence

Exploratory visualization:

- The observed residuals ε_i can be plotted against the **average values of their neighboring values ε_i^{avg}** using the “Moran’s plot”.

For the row-sum standardized link matrix \mathbf{V} these averages are given by $\boldsymbol{\varepsilon}^{avg} = \mathbf{V} \cdot \boldsymbol{\varepsilon}$.

average value of observations around that single position



value of each observed position

Under the assumption of $\epsilon \sim N(0, \sigma^2 \cdot \mathbf{I})$:

- Assuming that the population disturbances are **independently identically distributed** with $\epsilon \sim N(0, \sigma^2 \cdot \mathbf{I})$ the moments of Moran's I can be evaluated directly through the matrix terms:

$$E(I|H_0) = \frac{\text{tr}[\mathbf{M} \cdot \mathbf{V}]}{n - K}$$

$$\text{Var}(I|H_0) = \frac{\text{tr}[\mathbf{M} \cdot \mathbf{V} \cdot \mathbf{M} \cdot \mathbf{V}^T] + \text{tr}[\mathbf{M} \cdot \mathbf{V} \cdot \mathbf{M} \cdot \mathbf{V}] + (\text{tr}[\mathbf{M} \cdot \mathbf{V}])^2}{(n - K) \cdot (n - K + 2)} - [E(I|H_0)]^2$$

- For **sufficiently large number** of spatial objects n and **"well-behaved"** link matrices \mathbf{V} one can use the normal approximation to evaluate the distribution of I^{obs} :

$$\frac{I^{obs} - E(I|H_0)}{\sqrt{Var(I|H_0)}} \sim N(0,1)$$

projection matrix are not right in here, but a lot of software using it

- This expression is frequently used in software implementations.
- However, those implementations frequently use incorrectly the projections matrix $\mathbf{M}_{(1)} = \mathbf{1} \cdot (\mathbf{1}^T \cdot \mathbf{1})^{-1} \cdot \mathbf{1}^T$
- To evaluate the exact significance values of an observed value of Moran's I^{obs} numerical integration or the saddle-point approximation need to be employed.


Under the assumption of $\boldsymbol{\varepsilon} \sim N(0, \sigma^2 \cdot \boldsymbol{\Omega}(\rho))$:

- Again, these conditional distribution functions $F(I|\mathbf{X}, \boldsymbol{\Omega}(\rho))$ of Moran's I can be evaluated by numerical integration.

Under the assumption of $\boldsymbol{\varepsilon} \sim i.i.d$ with unknown distribution:

- In case the distribution of the residuals is unknown randomization needs to be employed to evaluated the significance of the observed Moran's I^{obs} . Data generating Process
- Underlying idea:
 - The spatial association of the residuals is broken up by randomly assigning the residuals to different areas.
 - This random assignment generates a map pattern under the assumption of spatial independence.

- This random assignment is repeated r -times.
- For each random pattern the associated Moran's I^{random} is calculated.
- The distribution of these random Moran's I^{random} under the null hypothesis is recorded.
- If the observed Moran's I^{obs} falls into a tail of this distribution, then the observed pattern exhibits significant spatial autocorrelation.



feasible general least square

Modelling an Autogressive Spatial Process with FGLS

- Let \mathbf{V} be the a $n \times n$ coded spatial link matrix among the n spatial objects.
- The error structure of disturbances **Gaussian spatial processes** becomes

$$\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma^2 \cdot \boldsymbol{\Omega}(\rho))$$

with

$$\begin{aligned}\boldsymbol{\Omega}_{AR}(\rho) &= (\mathbf{I} - \rho \cdot \mathbf{V})^{-1} \cdot (\mathbf{I} - \rho \cdot \mathbf{V}^T)^{-1} \text{ for an autoregressive process,} \\ \boldsymbol{\Omega}_{MA}(\rho) &= (\mathbf{I} + \rho \cdot \mathbf{V}) \cdot (\mathbf{I} + \rho \cdot \mathbf{V}^T) \text{ for a moving average process, and} \\ \boldsymbol{\Omega}_{CAR}(\rho) &= (\mathbf{I} - \rho \cdot \mathbf{V})^{-1} \text{ for a conditional autoregressive process.}\end{aligned}$$

Notes: For the CAR process the **link matrix needs to be symmetric**.

In all processes, if $\rho = 0$ the covariance structure reduces to the identity matrix \mathbf{I} , which implies spatial independence.

transfer dependent disturbance to independent and follow Gaussian distribution

- For the **AR process** we get the simple transformation $\boldsymbol{\Omega}_{AR}(\rho)^{-\frac{1}{2}} = [\mathbf{I} - \rho \cdot \mathbf{V}]$ with $\boldsymbol{\eta} = \boldsymbol{\Omega}_{AR}(\rho)^{-\frac{1}{2}} \cdot \boldsymbol{\varepsilon}$ with $\boldsymbol{\eta} = N(\mathbf{0}, \sigma^2 \cdot \mathbf{I})$.
- Therefore, the generalized least squares model for the AR-process becomes

$$\boldsymbol{\Omega}_{AR}(\rho)^{-\frac{1}{2}} \cdot \mathbf{y} = \boldsymbol{\Omega}_{AR}(\rho)^{-\frac{1}{2}} \cdot \mathbf{X} \cdot \boldsymbol{\beta} + \boldsymbol{\eta}$$

$$[\mathbf{I} - \rho \cdot \mathbf{V}] \cdot \mathbf{y} = [\mathbf{I} - \rho \cdot \mathbf{V}] \cdot \mathbf{X} \cdot \boldsymbol{\beta} + \boldsymbol{\eta}$$

$$\mathbf{y} = \rho \cdot \mathbf{V} \cdot \mathbf{y} + \mathbf{X} \cdot \boldsymbol{\beta} + \mathbf{V} \cdot \mathbf{X} \cdot \underbrace{\boldsymbol{\beta}_{AR}}_{=-\rho \cdot \boldsymbol{\beta}} + \boldsymbol{\eta}$$

which can be feasibly estimated for the unknown parameter $\hat{\sigma}^2$, $\hat{\rho}$ and $\hat{\boldsymbol{\beta}}$ with a maximum likelihood estimator.

- Notes:

1. For this structural form of the autoregressive model the endogenous variable \mathbf{y} is on both sides of the equation.
2. The terms $\mathbf{V} \cdot \mathbf{y}$ and $\mathbf{V} \cdot \mathbf{X}$ are spatially lagged terms of the dependent and independent variables.
3. If $\boldsymbol{\beta}_{AR} = \mathbf{0}$ then the model becomes the “spatial lag” model with $\mathbf{y} = \rho \cdot \mathbf{V} \cdot \mathbf{y} + \mathbf{X} \cdot \boldsymbol{\beta} + \boldsymbol{\eta}$.