

## MATRIX

1. If  $A$  is a square matrix of order 2 and  $|A| = -2$ , then the value of  $|5A'|$  is:  
(A)  $-50$   
(B)  $-10$   
(C)  $10$   
(D)  $50$
2. The product of matrix  $P$  and  $Q$  is equal to a diagonal matrix. If the order of matrix  $Q$  is  $3 \times 2$ , then the order of matrix  $P$  is:  
(A)  $2 \times 2$   
(B)  $3 \times 3$   
(C)  $2 \times 3$   
(D)  $3 \times 2$
3. If the inverse of the matrix  $\begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$  is the matrix  $\begin{bmatrix} 1 & 3 & 3 \\ 1 & \lambda & 3 \\ 1 & 3 & 4 \end{bmatrix}$ , then the value of  $\lambda$  is:  
(A)  $-4$   
(B)  $1$   
(C)  $3$   
(D)  $4$
4. Find the matrix  $A^2$ , where  $A = [a_{ij}]$  is a  $2 \times 2$  matrix whose elements are given by  $a_{ij} = \text{maximum}(i, j) - \text{minimum}(i, j)$ :  
(A)  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$   
(B)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$   
(C)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(D)  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

5. If  $A$  is a square matrix of order 3 such that the value of  $|\text{adj}.A| = 8$ , then the value of  $|A^T|$  is:

- (A)  $\sqrt{2}$   
 (B)  $-\sqrt{2}$   
 (C) 8  
 (D)  $2\sqrt{2}$

6. If  $A$  and  $B$  are events such that  $P(A/B) = P(B/A) \neq 0$ , then:

- (A)  $A \subset B$ , but  $A \neq B$   
 (B)  $A = B$   
 (C)  $A \cap B = \phi$   
 (D)  $P(A) = P(B)$

### DERIVATIONS

1. Derivative of  $e^{\sin^2 x}$  with respect to  $\cos x$  is:

- (A)  $\sin x e^{\sin^2 x}$   
 (B)  $\cos x e^{\sin^2 x}$   
 (C)  $-2\cos x e^{\sin^2 x}$   
 (D)  $-2\sin^2 x \cos x e^{\sin^2 x}$

2. If  $\sin(xy) = 1$ , then  $\frac{dy}{dx}$  is equal to:

- (A)  $\frac{x}{y}$   
 (B)  $-\frac{x}{y}$   
 (C)  $\frac{y}{x}$   
 (D)  $-\frac{y}{x}$

3. The general solution of the differential equation

$\frac{dy}{dx} = e^{x+y}$  is:

- (A)  $e^x + e^{-y} = c$
- (B)  $e^{-x} + e^{-y} = c$
- (C)  $e^{x+y} = c$
- (D)  $2e^{x+y} = c$

### EQUATIONS

1. Given a curve  $y = 7x - x^3$  and  $x$  increases at the rate of 2 units per second. The rate at which the slope of the curve is changing, when  $x = 5$  is:
  - (A)  $-60$  units/sec
  - (B)  $60$  units/sec
  - (C)  $-70$  units/sec
  - (D)  $-140$  units/sec
2. The area of the region bounded by the curve  $y^2 = 4x$  and  $x = 1$  is:
  - (A)  $\frac{4}{3}$
  - (B)  $\frac{8}{3}$
  - (C)  $\frac{64}{3}$
  - (D)  $\frac{32}{3}$
3. The angle which the line  $\frac{x}{1} = \frac{y}{-1} = \frac{z}{0}$  makes with the positive direction of Y-axis is:
  - (A)  $\frac{5\pi}{6}$
  - (B)  $\frac{3\pi}{4}$
  - (C)  $\frac{5\pi}{4}$
  - (D)  $\frac{7\pi}{4}$

### FUNCTIONS

1. The function  $f(x) = \frac{x}{2} + \frac{2}{x}$  has a local minima at  $x$  is equal to:
  - (A) 2
  - (B) 1

- (C) 0  
(D) -2

2. A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined as  $f(x) = x^2 - 4x + 5$  is:

- (A) injective but not surjective.  
(B) surjective but not injective.  
(C) both injective and surjective.  
(D) neither injective nor surjective.

### INTEGRATIONS

1. The value of  $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cot \theta \operatorname{cosec}^2 \theta d\theta$  is:

- (A)  $\frac{1}{2}$   
(B)  $-\frac{1}{2}$   
(C) 0  
(D)  $-\frac{\pi}{8}$

2. The integral  $\int \frac{dx}{\sqrt{9-4x^2}}$  is equal to:

- (A)  $\frac{1}{6} \sin^{-1}\left(\frac{2x}{3}\right) + c$   
(B)  $\frac{1}{2} \sin^{-1}\left(\frac{2x}{3}\right) + c$   
(C)  $\sin^{-1}\left(\frac{2x}{3}\right) + c$   
(D)  $\frac{3}{2} \sin^{-1}\left(\frac{2x}{3}\right) + c$

### VECTORS

1. The Cartesian equation of the line passing through the point  $(1, -3, 2)$  and parallel to the line:

$$\mathbf{r} = (2 + \lambda)\hat{i} + \lambda\hat{j} + (2\lambda - 1)\hat{k} \text{ is}$$

- (A)  $\frac{x-1}{2} = \frac{y+3}{0} = \frac{z-2}{-1}$   
(B)  $\frac{x+1}{2} = \frac{y-3}{1} = \frac{z+2}{2}$

$$(C) \frac{x+1}{2} = \frac{y-3}{0} = \frac{z+2}{-1}$$

$$(D) \frac{x-1}{1} = \frac{y+3}{1} = \frac{z-2}{2}$$

2. The position vectors of points  $P$  and  $Q$  are  $\mathbf{p}$  and  $\mathbf{q}$  respectively. The point  $R$  divides the line segment  $PQ$  in the ratio 3:1 and  $S$  is the mid-point of line segment  $PR$ . The position vector of  $S$  is:

$$(A) \frac{\mathbf{p}+3\mathbf{q}}{4}$$

$$(B) \frac{\mathbf{p}+3\mathbf{q}}{8}$$

$$(C) \frac{5\mathbf{p}+3\mathbf{q}}{4}$$

$$(D) \frac{5\mathbf{p}+3\mathbf{q}}{8}$$

3. **Assertion (A) :** The vectors

$$\mathbf{a} = 6\hat{i} + 2\hat{j} - 8\hat{k}$$

$$\mathbf{b} = 10\hat{i} - 2\hat{j} - 6\hat{k}$$

$$\mathbf{c} = 4\hat{i} - 4\hat{j} + 2\hat{k}$$

represent the sides of a right angled triangle.

**Reason (R) :** Three non-zero vectors of which none of two are collinear forms a triangle if their resultant is zero vector or sum of any two vectors is equal to the third.