

# ASSIGNMENT-12

R.YAMINI

## 1 QUESTION No-2.28(OPTIMIZATION)

Two godowns A and B have grain capacity of 100 quintals and 50 quintals respectively. They supply to 3 ration shops, D, E and F whose requirements are 60, 50 and 40 quintals respectively. The cost of transportation per quintal from the godowns to the shops are given in the following table: How

Transportation cost per quintal (in rupees)		
From/To	A	B
D	6	4
E	3	2
F	2.50	3

TABLE 1.1: Transportation table

should the supplies be transported in order that the transportation cost is minimum? What is the minimum cost?

## 2 SOLUTION

Let A supply  $x$  quintals to D and  $y$  quintals to E. From the data given we have

$$x + y \leq 100 \quad (2.0.1)$$

$$x \leq 60 \quad (2.0.2)$$

$$y \leq 50 \quad (2.0.3)$$

$$x + y \geq 60 \implies -x - y \leq -60 \quad (2.0.4)$$

Now our aim is to minimize the transportation cost, we obtain the minimizing function as,

$$\min_{\mathbf{x}} Z = (2.5 \quad 1.5) \mathbf{x} + 410. \quad (2.0.5)$$

Subject to the constraints,

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & -1 \\ -1 & -1 \end{pmatrix} \mathbf{x} \leq \begin{pmatrix} 100 \\ 60 \\ 50 \\ -60 \end{pmatrix} \quad (2.0.6)$$

We solve the optimization problem by the method of Lagrange multipliers, thus the Lagrangian function is given as follows,

$$\begin{aligned} L(\mathbf{x}, \lambda) &= (2.5 \quad 1.5) \mathbf{x} + 410 + [(1 \quad 1) \mathbf{x} - 100] \lambda_1 \\ &+ [(1 \quad 0) \mathbf{x} - 60] \lambda_2 + [(0 \quad 1) \mathbf{x} - 50] \lambda_3 \\ &+ [(-1 \quad -1) \mathbf{x} + 60] \lambda_4 + [(-1 \quad 0) \mathbf{x}] \lambda_5 + [(0 \quad -1) \mathbf{x}] \lambda_6 \end{aligned} \quad (2.0.7)$$

Now, we have

$$\nabla L(\mathbf{x}, \lambda) = \begin{pmatrix} 2.5 + (1 \quad 1 \quad 0 \quad -1 \quad -1 \quad 0) \lambda \\ 1.5 + (1 \quad 0 \quad 1 \quad -1 \quad 0 \quad -1) \lambda \\ (1 \quad 1) \mathbf{x} - 100 \\ (1 \quad 0) \mathbf{x} - 60 \\ (0 \quad 1) \mathbf{x} - 50 \\ (-1 \quad -1) \mathbf{x} + 60 \\ (-1 \quad 0) \mathbf{x} \\ (0 \quad -1) \mathbf{x} \end{pmatrix} \quad (2.0.8)$$

where  $\lambda = \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \\ \lambda_4 \\ \lambda_5 \\ \lambda_6 \end{pmatrix}$ . Therefore, Lagrangian matrix is

given by

$$\begin{pmatrix} 0 & 0 & 1 & 1 & 0 & -1 & -1 & 0 \\ 0 & 0 & 1 & 0 & 1 & -1 & 0 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} -2.5 \\ -1.5 \\ 100 \\ 60 \\ 50 \\ -60 \\ 0 \\ 0 \end{pmatrix} \quad (2.0.9)$$

We see that  $\lambda_3, \lambda_4$  are the only active multiplier and hence we proceed by considering them,

$$\begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \lambda_{i=3,4} \end{pmatrix} = \begin{pmatrix} -2.5 \\ -1.5 \\ 50 \\ -60 \end{pmatrix} \quad (2.0.10)$$

Now we have,

$$\begin{pmatrix} \mathbf{x} \\ \lambda_{i=3,4} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{pmatrix}^{-1} \begin{pmatrix} -2.5 \\ -1.5 \\ 50 \\ -60 \end{pmatrix} \quad (2.0.11)$$

$$\Rightarrow \begin{pmatrix} \mathbf{x} \\ \lambda_{i=3,4} \end{pmatrix} = \begin{pmatrix} 0 & 0 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} -2.5 \\ -1.5 \\ 50 \\ -60 \end{pmatrix} \quad (2.0.12)$$

$$\Rightarrow \begin{pmatrix} \mathbf{x} \\ \lambda_{i=3,4} \end{pmatrix} = \begin{pmatrix} 10 \\ 50 \\ 1 \\ 2.5 \end{pmatrix} \quad (2.0.13)$$

$\therefore \lambda_3, \lambda_4 > 0$

Thus, the Optimal solution is given as follows

$$\mathbf{x} = \begin{pmatrix} 10 \\ 50 \end{pmatrix} \quad (2.0.14)$$

$$Z = (2.5 \quad 1.5) \mathbf{x} + 410 \quad (2.0.15)$$

$$= (2.5 \quad 1.5) \begin{pmatrix} 10 \\ 50 \end{pmatrix} \quad (2.0.16)$$

$$= 510 \quad (2.0.17)$$

Hence, in order to minimize the transportation cost we see that A must supply 10 quintals to D and 50 quintals to E, and the obtained minimum cost is ₹510.




Fig. 2.1: Graphical Solution