Reactivity of 12-tungstophosphoric acid and its inhibitor potency toward Na⁺/K⁺-ATPase: a combined ³¹P *NMR* study, *ab initio* calculations and crystallographic analysis

Nada Bošnjaković-Pavlović^{a*}, Danica Bajuk-Bogdanović^a, Joanna Zakrzewska^b,

Zeyin Yan^{c,d}, Ivanka Holclajtner-Antunović^a, Jean-Michel Gillet^{c,d}

and Anne Spasojević-de Biré^{c,d*}

^a Faculty of Physical Chemistry, University of Belgrade, P.O. Box 47, 11001 Belgrade, Serbia ^b Institute of General and Physical Chemistry, Belgrade, Serbia

^c Université Paris-Saclay, CentraleSupélec, Campus de Gif, Plateau du Moulon, 3 rue Joliot Curie, 91192 Gif-sur-Yvette, France

^d CNRS, UMR 8580, Laboratory "Structures Propriétés et Modélisation des Solides" (SPMS), Campus de Gif, Plateau du Moulon, 3 rue Joliot Curie, 91192 Gif-sur-Yvette, France

Supporting information

- S1. Structures containing Keggin anion (WPA³⁻) found in CSD database
- S2. Structures containing Keggin anion (WPA³⁻) found in COD database
- S3. Structures containing Keggin anion (WPA³⁻) found in ICSD database
- S4. Structures containing Keggin anion (WPA³⁻) found in PDB database

- S1. a) Y.Z. Han, C.L. Zhang, Q. Cai, Y. Cheng, S.X. Ai, Q. Deng, Syntheses, Crystal Structure, Characterization and Properties of Two Novel Superamolecular Phosphotungstate Pyridine Complexes, *Synth. React. Inorg. Met. Org. Chem.* 46 (**2016**) 351-364.
- b) D. Meziani, K. Abdmeziem, S. Bouacida, M. Trari, Photo-electrochemical and physical characterizations of a new single crystal POM-based material. Application in photocatalysis, *J. Mol. Struct.* 1125 (**2016**) 540-545.
- c) C.L. Teng, H.X. Xiao, Q. Cai, J.T. Tang, T.J. Cai, Q. Deng, Two multifunctional organic-inorganic hybrid complexes based on polyoxometalates, Bi EDTA and sodium linker: crystal structures, photochromic, and catalytic performances, *J. Coord. Chem.* 69 (14) (2016) 2148-2163.
- d) X. Wang, C.Y. Kong, J.J. Lai, M.L. Wei, Synthesis, Structure and Proton Conductivity of a Complex Based on Decorated Keggin-Type Cluster: {[Cu(dmbipy)(H₂O)₂Cl_{0.5}]₂[PW₁₂O₄₀]}.7H₂O (dmbipy = 4,4'-dimethyl-2,2'-bipyridine), *J. Clust. Sci.* 27 (**2016**) 645-656.
- e) D. Chen, Z. Deng, Q. Li, X. Wang, X. Liu, Synthesis, crystal structure and catalytic behavior of a novel polyoxometalate-based complex, *Inorg. Chim. Acta* 438 (**2015**) 208-211.
- f) H. Hu, W. Zhang, J. Gong, H. Dong, F. Zhao, H.Huang, Y.Liu, G. Zhang, Z. Kang, Can tetrazole-functionalized ligands realize the role-control of Keggin-type POMs in hybridframeworks?, *Cryst. Eng. Comm.* 16 (2014) 5642-5649.
- g) A.X. Tian, Y. Yang, N. Sun, J.C. Li, J. Ying, J.W. Zhang, X.L.Wang, Three 2-(4-thiazolyl)benzimidazole-based supramolecular assemblies oriented by Keggin and Wells–Dawson anions, *J. Coord. Chem.* 67 (9) (2014) 1550-1561.

- h) H. Zhang, K. Yu, S. Gao, C. Wang, C. Wang, H. Wang, B. Zhou, Assembly of two-fold interpenetrated silver supramolecular coordination polymer using Keggin phosphotungstate template, *Inorg. Chem. Commun.* 44 (2014) 91-95.
- i) D. Chen, A. Sahasrabudhe, P. Wang, A. Dasgupta, R. Yuan, S. Roy, Synthesis and properties of a novel quarternerized imidazolium $[\alpha\text{-PW}_{12}O_{40}]^{3}$ salt as a recoverable photopolymerization catalyst, *Dalton Trans.* 42(29) (2013) 10587-96.
- j) X. Li, L. Cheng, W. Fang, G. Yang, A Series of Open-Frameworks Constructed From Polyoxoanion Clusters and Copper-tetrazolate Complexes: Synthesis, Structure and Properties, Acta Chim. Sinica 71 (2013) 179-185.
- k) L. Liu, X.Z. Xu, X. Zhang, N.X. Zhang, S.G. Fan, Y.P. Huang, Synthesis, Crystal structure and Luminescent Property of a New 3D Supramolecular Compound $(C_6H_6NO_2)_3(C_6H_5NO_2)(PW_{12}O_{40})\cdot 2H_2O$, Chin. J. Struct. Chem. 32 (2013) 1813-1820.
- X. Wang, J. Peng, K. Alimaje, Z. Zhang, Z. Shi, Two Keggin polyoxoanion-based hybrids composed of CuI/tbz segments: Syntheses, structures and electrochemical properties, *Inorg. Chem. Commun.* 36 (2013) 141-145.
- m) M. Wei, X. Wang, J. Sun, X. Duan, A 3D POM–MOF composite based on Ni(II) ion and 2,2'-bipyridyl-3,3'-dicarboxylicacid: Crystal structure and proton conductivity, *J. Solid State Chem.* 202 (2013) 200-206.
- n) X. Zhao, X. Sun, Z. Han, C. Zhao, H. Yu, X. Zhai, Two new polyoxometalate-based hybrids consisting of Keggin-type cluster modified by {Ag₄} group, *J. Solid State Chem.* 207 (2013) 178-183.
- o) M.G. Liu, P.P. Zhang, J. Peng, H.X. Meng, X. Wang, M. Zhu, D.D. Wang, C.L. Meng, K. Alimaje, Organic-Inorganic Hybrids Constructed from Mixed-Valence Multinuclear,

- Copper Complexes and Templated by Keggin Polyoxometalates, *Cryst. Growth Des.* 12(3) (2012) 1273-1281.
- p) C.J. Zhang, H.J. Pang, Q. Tang, Y.G. Chen, A feasible route to approach 3D POM-based hybrids: utilizing substituted or reduced Keggin anions with high charge density, *Dalton Trans.* 41 (2012) 9365-9372.
- q) Y. Jiang, S. Liu, J. Miao, J. Zhang, L. Wu, Anisotropic ionic liquids built from nonmesogenic cation surfactants and Keggin-type polyoxoanions, *Chem. Commun.* 47 (2011) 10287-10289.
- r) C. Pichon, P. Mialane, J. Marrot, C. Binet, A. Vimont, A. Travert, J.-C. Lavalley, Combined XRD and infrared studies of pyridinium species in (PyH) 3[PW₁₂O₄₀] single crystals, *Phys. Chem. Chem. Phys.* 13 (**2011**) 322–327.
- s) L.N. Xiao, Y. Wang, C.L. Pan, J. N. Xu, T.G. Wang, H. Ding, Z.M. Gao, D.F. Zheng, X.B. Cui, J.Q. Xu, Three novel supramolecular hybrid compounds based on kegging polytungstates, *Cryst. Eng. Comm.* 13 (2011) 4878-4885.
- t) Y.P. Ren, X.J. Kong, X.Y. Hu, M. Sun, L.S. Long, R.B. Huang, L.S. Zheng, Influence of Steric Hindrance of Organic Ligand on the Structure of Keggin-Based Coordination Polymer, *Inorg Chem.* 45(10) (2006) 4016-4023.
- u) M. Schulz-Dobrick, M. Jansen, Supramolecular Intercluster Compounds Consisting of Gold Clusters and Keggin Anions, *Eur. J. Inorg. Chem.* (2006) 4498-4502.
- v) X. Lu, B. Liu, Sarula, J. Wang, C. Ye, Synthesis, crystal structure and NMR of $[Na(DB_{18}C_6)(CH_3CN)]_3[\alpha\text{-PW}_{12}O_{40}], \textit{Polyhedron 24 (2005) 2889-2893}$
- w) Z. Han, Y. Zhao, J. Peng, H. Ma, Q. Liu, E. Wang, N. Hu, Supramolecular assembly of organic bicapped Keggin Polyoxometalate, *J. Solid State Chem.* 177 (2004) 4325-4331.

- x) J. Wang, X. Duan, J. Niu, Synthesis and crystal structure of a 1D double-chain polyoxometalate based complex {[Ba(DMSO)₃(H₂O)₃] [C₅H₆N] PW₁₂O₄₀·DMSO·2H₂O}_n, J. Mol. Struct. 692 (2004) 17-21.
- y) W. Huang, T. Louis, P.A. Glenn, R.B. Yap, L.C. Francesconi. T. Polenova, ⁵¹V Magic Angle Spinning NMR Spectroscopy of Keggin Anions [PV_nW_{12-n}O₄₀]⁽³⁺ⁿ⁾⁻: Effect of Countercation and Vanadium Substitution on Fine Structure Constants, *J. Am. Chem. Soc.* 126 (2004) 11564-11573.
- z) C. Inman, J.M. Knaust, S. W. Keller, A polyoxometallate-templated coordination polymer: synthesisand crystal structure of [Cu₃(4,4'-bipy)₅(MeCN)₂]PW₁₂O₄₀·2C₆H₅CN, *Chem. Commun.* (2002) 156-157.
- α) Y. Xu, K.L. Zhang, Y. Zhang, X.Z. You, J.Q. Xu, Keggin unit supported transition metal complexes: hydrothermal synthesis and characterization of [Ni(2,2'-bipy)₃]_{1.5} [PW₁₂O₄₀Ni(2,2'-bipy)₂(H₂O)]·0.5H₂O and [Co(1,10'-phen)₃]_{1.5}[PMo₁₂O₄₀Co(1,10'-phen)₂(H₂O)]·0.5H₂O, *Chem. Commun.* (2000) 153-154.

S2. M.H. Alizadeh, I.M. Zonoz, 12-Tungstophosphoric acid-1,1'-methylenebis(imidazolidine-2,4-dione)-imidazolidine-2,4-dione-water, *Acta Crystallogr*. E66 (**2010**) m839.

- S3. a) F. Gschwind, M. Jansen, Crystal structure of hexakis (dimethylsulfoxide-κ-*O*) iron(III) dodecatungstophosphate, C₁₂H₃₆FeO₄₆PS₆W₁₂, *Z. Krist.-New Cryst. Struct.* 229 (**2014**) 5-6.
- b) S. Busbongthong, T. Ozeki, Structural Relationships among Methyl-, Dimethyl-, and Trimethylammonium Phosphododecatungstates, *Bull. Chem. Soc. Jpn.* 82 (2009) 1393-1397.
- c) Q.X. Han, C.Z. Wang, J.R. Ma, J.Y. Niu, J.P. Wang, Synthesis, Crystal Structure and Electrochemistry of Two Complexes $(YL_n)^{3+}[PW_{12}O_{40}]^{3-}$ (L=Me₂SO, Me₂NCHO), *Chem. Res. Chinese Univ.* 24 (**2008**) 255-259.
- d) A. Kremenović, D. Poleti, N.E. Ghermani, Lj. Karanović, U.B. Mioč, Z. Nedić, A. Spasojević-de Biré, Synthesis and crystal structure of hexaaquamagnesium hydrogen phosphododecatungstate tetrahydrate (Mg(H₂O)₆) (HPW₁₂O₄₀) * 4(H₂O), *Cryst. Res. Technol.* 42 (2007) 98-104.
- e) A. Kremenović, A. Spasojević-de Biré, F. Bourée, P. Colomban, R. Dimitrijević, M. Davidović, U.B. Mioč, Structural modifications of dodecatungstophosphoric acid hexahydrate induced by temperature in the 10–358 K range. In situ high-resolution neutron powder diffraction investigation, *Solid State Ion.* 150 (2002) 431-442.
- f) M.R. Spirlet, W.R. Busing, Dodecatungstophosphoric acid-21-water by neutron diffraction, *Acta Crystallogr.* B34 (1978) 907-910,
- g) G.M. Brown, M.R. Noe-Spirlet, W.R. Busing, H.A. Levy, Dodecatungstophosphoric acid hexahydrate, (H₅O₂)₃ (PW₁₂O₄₀), the true structure of Keggin's 'pentahydrate' from single crystal X-ray and neutron diffraction data, *Acta Crystallogr*. B33 (1977) 1038-1046.

- S4. a) S.C. Almo, J. Bonanno, J.M Sauder, S. Emtage, T.P. Dilorenzo, V. Malashkevich, S.R. Wasserman, S. Swaminathan, S. Eswaramoorthy, R Agarwal, D. Kumaran, M. Madegowda, S. Ragumani, Y. Patskovsky, J. Alvarado, U.A. Ramagopal, J. Faber-Barata, MR. Chance, A. Sali, A. Fiser, Z-Y. Zhang, D.S. Lawrence, S.K. Burley, Structural Genomics of Protein Phosphatases, *J. Struct. Funct. Genom.* 8 (2007) 121-140.
- b). G. Rudenko, L. Henry, K. Henderson, K. Ichtchenko, M.S. Brown, J.L. Goldstein, J. Deisenhofer, Structure of the LDL receptor extracellular domain at endosomal, *Science* 298 (2002) 2353-2358.
- c). Y. Ren, P. Schmiege, G. Blobel, Structural and biochemical analyses of the DEAD-box ATPase Sub2 in association with THO or Yra1, *eLife* 6 e20070 (**2017**) 1-17