

Assignment 3: Data Exploration

Yang Wang

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Exploration.

Directions

1. Change “Student Name” on line 3 (above) with your name.
2. Work through the steps, **creating code and output** that fulfill each instruction.
3. Be sure to **answer the questions** in this assignment document.
4. When you have completed the assignment, **Knit** the text and code into a single PDF file.
5. After Knitting, submit the completed exercise (PDF file) to the dropbox in Sakai. Add your last name into the file name (e.g., “Salk_A03_DataExploration.Rmd”) prior to submission.

The completed exercise is due on Tuesday, January 28 at 1:00 pm.

Set up your R session

1. Check your working directory, load necessary packages (tidyverse), and upload two datasets: the ECOTOX neonicotinoid dataset (ECOTOX_Neonicotinoids_Insects_raw.csv) and the Niwot Ridge NEON dataset for litter and woody debris (NEON_NIWO_Litter_massdata_2018-08_raw.csv). Name these datasets “Neonics” and “Litter”, respectively.

```
getwd()
```

```
## [1] "C:/Users/26059/OneDrive/Desktop/ENV 872 R/Yang_ENV872"
```

```
library(tidyverse)
Neonics <- read.csv("./Data/Raw/ECOTOX_Neonicotinoids_Insects_raw.csv")
Litter <- read.csv("./Data/Raw/NEON_NIWO_Litter_massdata_2018-08_raw.csv")
```

Learn about your system

2. The neonicotinoid dataset was collected from the Environmental Protection Agency’s ECOTOX Knowledgebase, a database for ecotoxicology research. Neonicotinoids are a class of insecticides used widely in agriculture. The dataset that has been pulled includes all studies published on insects. Why might we be interested in the ecotoxicology of neonicotinoids on insects? Feel free to do a brief internet search if you feel you need more background information.

Answer: Neonicotinoid pesticides are widely applied to agriculture land. The residue of neonicotinoid may pose risk to non-target beneficial insects. Insects are one of important parts of a healthy ecosystem. Therefore, we might be interested in the ecotoxicology of this chemical for ecosystem health and balance.

3. The Niwot Ridge litter and woody debris dataset was collected from the National Ecological Observatory Network, which collectively includes 81 aquatic and terrestrial sites across 20 ecoclimatic domains. 32 of these sites sample forest litter and woody debris, and we will focus on the Niwot Ridge long-term ecological research (LTER) station in Colorado. Why might we be interested in studying litter and woody debris that falls to the ground in forests? Feel free to do a brief internet search if you feel you need more background information.

Answer: Insects constantly contact with litter and woody debris. Knowing the concentration of chemical in litter and woody debris can help to identify the chemical exposure of insects.

4. How is litter and woody debris sampled as part of the NEON network? Read the NEON_Litterfall_UserGuide.pdf document to learn more. List three pieces of salient information about the sampling methods here:

Answer: *spatial sampling design* random for the tower plots, and target for the forest tower
airshed *temporal sampling design

Obtain basic summaries of your data (Neonics)

5. What are the dimensions of the dataset?

```
dim(Neonics)
```

```
## [1] 4623 30
```

6. Using the `summary` function, determine the most common effects that are studied. Why might these effects specifically be of interest?

```
summary(Neonics$Effect)
```

```
##      Accumulation      Avoidance      Behavior      Biochemistry
##           12           102           360           11
##      Cell(s)      Development      Enzyme(s)      Feeding behavior
##           9           136           62           255
##      Genetics      Growth      Histology      Hormone(s)
##          82           38           5           1
##      Immunological      Intoxication      Morphology      Mortality
##          16           12           22           1493
##      Physiology      Population      Reproduction
##           7           1803           197
```

Answer: Population and mortality are the most common effects that are studied. Because both of them can reflect the acute effect of the chemical, while other endpoints are reflecting chronic effect.

7. Using the `summary` function, determine the six most commonly studied species in the dataset (common name). What do these species have in common, and why might they be of interest over other insects? Feel free to do a brief internet search for more information if needed.

```
summary(Neonics$Species.Common.Name)
```

##	Honey Bee	Parasitic Wasp
##	667	285
##	Buff Tailed Bumblebee	Carniolan Honey Bee
##	183	152
##	Bumble Bee	Italian Honeybee
##	140	113
##	Japanese Beetle	Asian Lady Beetle
##	94	76
##	Euonymus Scale	Wireworm
##	75	69
##	European Dark Bee	Minute Pirate Bug
##	66	62
##	Asian Citrus Psyllid	Parastic Wasp
##	60	58
##	Colorado Potato Beetle	Parasitoid Wasp
##	57	51
##	Erythrina Gall Wasp	Beetle Order
##	49	47
##	Snout Beetle Family, Weevil	Sevenspotted Lady Beetle
##	47	46
##	True Bug Order	Buff-tailed Bumblebee
##	45	39
##	Aphid Family	Cabbage Looper
##	38	38
##	Sweetpotato Whitefly	Braconid Wasp
##	37	33
##	Cotton Aphid	Predatory Mite
##	33	33
##	Ladybird Beetle Family	Parasitoid
##	30	30
##	Scarab Beetle	Spring Tiphia
##	29	29
##	Thrip Order	Ground Beetle Family
##	29	27
##	Rove Beetle Family	Tobacco Aphid
##	27	27
##	Chalcid Wasp	Convergent Lady Beetle
##	25	25
##	Stingless Bee	Spider/Mite Class
##	25	24
##	Tobacco Flea Beetle	Citrus Leafminer
##	24	23
##	Ladybird Beetle	Mason Bee
##	23	22
##	Mosquito	Argentine Ant
##	22	21
##	Beetle	Flatheaded Appletree Borer
##	21	20
##	Horned Oak Gall Wasp	Leaf Beetle Family
##	20	20
##	Potato Leafhopper	Tooth-necked Fungus Beetle

##	20	20
##	Codling Moth	Black-spotted Lady Beetle
##	19	18
##	Calico Scale	Fairyfly Parasitoid
##	18	18
##	Lady Beetle	Minute Parasitic Wasps
##	18	18
##	Mirid Bug	Mulberry Pyralid
##	18	18
##	Silkworm	Vedalia Beetle
##	18	18
##	Araneoid Spider Order	Bee Order
##	17	17
##	Egg Parasitoid	Insect Class
##	17	17
##	Moth And Butterfly Order	Oystershell Scale Parasitoid
##	17	17
##	Hemlock Woolly Adelgid Lady Beetle	Hemlock Woolly Adelgid
##	16	16
##	Mite	Onion Thrip
##	16	16
##	Western Flower Thrips	Corn Earworm
##	15	14
##	Green Peach Aphid	House Fly
##	14	14
##	Ox Beetle	Red Scale Parasite
##	14	14
##	Spined Soldier Bug	Armoured Scale Family
##	14	13
##	Diamondback Moth	Eulophid Wasp
##	13	13
##	Monarch Butterfly	Predatory Bug
##	13	13
##	Yellow Fever Mosquito	Braconid Parasitoid
##	13	12
##	Common Thrip	Eastern Subterranean Termite
##	12	12
##	Jassid	Mite Order
##	12	12
##	Pea Aphid	Pond Wolf Spider
##	12	12
##	Spotless Ladybird Beetle	Glasshouse Potato Wasp
##	11	10
##	Lacewing	Southern House Mosquito
##	10	10
##	Two Spotted Lady Beetle	Ant Family
##	10	9
##	Apple Maggot	(Other)
##	9	670

Answer: The most common is honey bee, parasitic wasp, buff tailed bumblebee, carniolan honey bee, bumble bee and Italian honeybee. The six most commonly species are all pollinators. They got more attention than other species because they are important pollinators for agriculture crops.

8. Concentrations are always a numeric value. What is the class of `Conc.1..Author.` in the dataset, and why is it not numeric?

```
class(Neonics$Conc.1..Author.)
```

```
## [1] "factor"
```

```
str(Neonics$Conc.1..Author.)
```


```
## Factor w/ 1006 levels "~10","~30/","~40/",...: 639 510 813 622 442 637 500 642 814 784 ...
```

Answer: It is factor. Because it has non-numeric subjects and symbol like slash..

Explore your data graphically (Neonics)


9. Using `geom_freqpoly`, generate a plot of the number of studies conducted by publication year.

```
ggplot(Neonics) +  
  geom_freqpoly(aes(x = Publication.Year), binwidth = 1)
```



10. Reproduce the same graph but now add a color aesthetic so that different `Test.Location` are displayed as different colors.

```
ggplot(Neonics) +  
  geom_freqpoly(aes(x = Neonics$Publication.Year, color = Neonics$Test.Location ),binwidth = 1)
```




Interpret this graph. What are the most common test locations, and do they differ over time?

Answer:

11. Create a bar graph of Endpoint counts. What are the two most common end points, and how are they defined? Consult the ECOTOX_CodeAppendix for more information.

```
ggplot(Neonics, aes(x = Neonics$Endpoint)) +  
  geom_bar()
```



Answer: The most common end points are LOEL and NOEL. LOEL is the lowest observed adverse effect level. NOEL is the highest no observable effect level.

Explore your data (Litter)

- Determine the class of collectDate. Is it a date? If not, change to a date and confirm the new class of the variable. Using the `unique` function, determine which dates litter was sampled in August 2018.

```
class(Litter$collectDate)
```

```
## [1] "factor"
```

```
Litter$collectDate <- as.Date(Litter$collectDate,format="%Y-%m-%d")
class(Litter$collectDate)
```

```
## [1] "Date"
```

```
unique(Litter$collectDate)
```

```
## [1] "2018-08-02" "2018-08-30"
```

- Using the `unique` function, determine how many plots were sampled at Niwot Ridge. How is the information obtained from `unique` different from that obtained from `summary`?

```
unique(Litter$plotID)
```

```
## [1] NIWO_061 NIWO_064 NIWO_067 NIWO_040 NIWO_041 NIWO_063 NIWO_047  
## [8] NIWO_051 NIWO_058 NIWO_046 NIWO_062 NIWO_057  
## 12 Levels: NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 ... NIWO_067
```


```
summary(Litter$plotID)
```

```
## NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 NIWO_058 NIWO_061  
##      20      19      18      15      14       8      16      17  
## NIWO_062 NIWO_063 NIWO_064 NIWO_067  
##      14      14      16      17
```

Answer: 12 plots. The information obtained from unique function does not include the count of the same plot. However, It gives the number of the unique plots and has a rank of the plot code.


14. Create a bar graph of functionalGroup counts. This shows you what type of litter is collected at the Niwot Ridge sites. Notice that litter types are fairly equally distributed across the Niwot Ridge sites.

```
ggplot(Litter, aes(x = Litter$functionalGroup)) +  
  geom_bar()
```




15. Using `geom_boxplot` and `geom_violin`, create a boxplot and a violin plot of `dryMass` by `functionalGroup`.


```
ggplot(Litter) +  
  geom_boxplot(aes(x = Litter$functionalGroup, y = dryMass))
```



```
ggplot(Litter) +  
  geom_violin(aes(x = Litter$functionalGroup, y = dryMass),  
             scale="area")
```



Why is the boxplot a more effective visualization option than the violin plot in this case?

Answer: The dry mass between different functional groups vary too much.

What type(s) of litter tend to have the highest biomass at these sites?

Answer: Needles