

Wigner and MP laws

Contents

Wigner

Illustration of convergence of averaged eigenvalue distribution

Experiment

Plot

Theory

Experiment

Plot

Theory

Illustration of almost sure convergence of eigenvalue distribution

Experiment

Plot

Theory

Experiment

Plot

Theory

Experiment

Plot

Theory

MP law

Illustration of convergence of averaged eigenvalue distribution

Experiment

Plot

Theory

Illustration of almost sure convergence of eigenvalue distribution

Experiment

Plot

Theory

Wigner

Illustration of convergence of averaged eigenvalue distribution

```
close all;  
clear all;  
rng('default');
```

```

N = 5;
K = 1e4;
v = []; % eigenvalues
dx = 0.1; % bin size

```

Experiment

```

for i =1:K
    A =randn(N)/sqrt(N/2); % random NxN Gaussian matrix; Note: normalize by sqrt(N/2)
    and not sqrt(N) so AA will have unit variance.
    AA =(A+A')/2; % symmetrize
    v = [v; eig(AA)]; % eigenvalues
end


```

Plot

```

[count, x]= hist(v, -3:dx:3);
bar(x, count/(K*N*dx), 'y');
hold on ;

```




Theory

```

x = -2:0.01:2;
plot(x,sqrt(4-x.^2)/(2*pi), 'LineWidth', 2);
axis([-3 3 0 0.35]);
grid on;
title('N=5, K=10000');
xlabel('\lambda');

```

```
ylabel('p(\lambda)');  
%%%%%%%%  
%%%%%%%%  
figure;  
N = 20;  
K = 1e4;  
v = []; % eigenvalues  
dx = 0.1; % bin size
```



Experiment

```
for i =1:K
    A =randn(N)/sqrt(N/2); % random NxN Gaussian matrix; Note: normalize by sqrt(N/2)
    and not sqrt(N) so AA will have unit variance.
```

```

AA = (A+A')/2; % symmetrize
v = [v; eig(AA)]; % eigenvalues
end


```

Plot

```

[count, x]= hist(v, -3:dx:3);
bar(x, count/(K*N*dx), 'y');
hold on ;

```



Theory

```

x = -2:0.01:2;
plot(x, sqrt(4-x.^2)/(2*pi), 'LineWidth', 2);
axis([-3 3 0 0.35]);
grid on;
title('N=20, K=10000');
xlabel('\lambda');
ylabel('p(\lambda)');

```




Illustration of almost sure convergence of eigenvalue distribution


```
figure;
N = 10;
K = 1;
v = []; % eigenvalues
dx = 2/3; % bin size
```

Experiment

```
for i =1:K
    A =randn(N)/sqrt(N/2); % random NxN Gaussian matrix; Note: normalize by sqrt(N/2)
    and not sqrt(N) so AA will have unit variance.
    AA = (A+A')/2; % symmetrize
    v = [v; eig(AA)]; % eigenvalues
end
```


Plot

```
[count, x]= hist(v, -2:dx:2);
bar(x, count/(K*N*dx), 'y');
hold on ;
```



Theory

```
x = -2:0.01:2;
plot(x,sqrt(4-x.^2)/(2*pi), 'LineWidth', 2);
axis([-3 3 0 0.5]);
grid on;
title('N=10, K=1');
xlabel('\lambda');
ylabel('p(\lambda)');
%%%%%%%%%
%%%%%%%%%
figure;
N = 100;
K = 1;
v = []; % eigenvalues
dx = 0.4; % bin size
```

Experiment

```
for i =1:K
    A =randn(N)/sqrt(N/2); % random NxN Gaussian matrix; Note: normalize by sqrt(N/2)
    and not sqrt(N) so AA will have unit variance.
```

```

AA = (A+A')/2; % symmetrize
v = [v; eig(AA)]; % eigenvalues
end


```

Plot

```

[count, x]= hist(v, -3:dx:3);
bar(x, count/(K*N*dx), 'y');
hold on ;

```




Theory

```

x = -2:0.01:2;
plot(x,sqrt(4-x.^2)/(2*pi), 'LineWidth', 2);
axis([-3 3 0 0.35]);
grid on;
title('N=100, K=1');
xlabel('\lambda');
ylabel('p(\lambda)');
%%%%%%%%%
%%%%%%%%%
figure;
N = 1000;
K = 1;
v = []; % eigenvalues
dx = 1/6; % bin size

```



Experiment

```
for i =1:K
    A =randn(N)/sqrt(N/2); % random NxN Gaussian matrix; Note: normalize by sqrt(N/2)
    and not sqrt(N) so AA will have unit variance.
```

```

AA = (A+A')/2; % symmetrize
v = [v; eig(AA)]; % eigenvalues
end


```

Plot

```

[count, x]= hist(v, -2:dx:2);
bar(x, count/(K*N*dx), 'y');
hold on ;

```




Theory

```

x = -2:0.01:2;
plot(x,sqrt(4-x.^2)/(2*pi), 'LineWidth', 2);
axis([-3 3 0 0.35]);
grid on;
title('N=1000, K=1');
xlabel('\lambda');
ylabel('p(\lambda)');
%%%%%%%%
%%%%%%%%
figure;
N = 4000;
K = 1;
v = []; % eigenvalues
dx = 0.1; % bin size

```



Experiment

```

for i =1:K
    A =randn(N)/sqrt(N/2); % random NxN Gaussian matrix; Note: normalize by sqrt(N/2)
    and not sqrt(N) so AA will have unit variance.

```

```

AA = (A+A')/2; % symmetrize
v = [v; eig(AA)]; % eigenvalues
end


```

Plot

```

[count, x]= hist(v, -3:dx:3);
bar(x, count/(K*N*dx), 'y');
hold on ;

```




Theory

```

x = -2:0.01:2;
plot(x, sqrt(4-x.^2)/(2*pi), 'LineWidth', 2);
axis([-3 3 0 0.35]);
grid on;
title('N=4000, K=1');
xlabel('\lambda');
ylabel('p(\lambda)');

```



MP law

Illustration of convergence of averaged eigenvalue distribution


```
figure;
N = 100;
M = 200;
K = 3000;
v = []; % eigenvalues
dx = 0.25; % bin size
```

Experiment

```
for i =1:K
    X =randn(N,M); % random NxN Gaussian matrix;
    A =X*X'/N; % Wishart
    v = [v; eig(A)]; % eigenvalues
end
```

Plot

```
alpha = M/N;
%[count, x]= hist(v, (1-sqrt(alpha))^2:dx:(1+sqrt(alpha))^2);
[count, x]= hist(v, 0:dx:6);
bar(x, count/(K*N*dx), 'y');
hold on ;
```

Theory

```
x = (1-sqrt(alpha))^2:0.01:(1+sqrt(alpha))^2;
plot(x,sqrt(4*alpha-(x-1-alpha).^2)./(2*pi*x), 'LineWidth', 2);
axis([-0.6 0.5]);
grid on;
title('N=100, \alpha=2, K=3000');
xlabel('\lambda');
ylabel('p(\lambda)');
```




Illustration of almost sure convergence of eigenvalue distribution


```
figure;
N = 2000;
M = 4000;
K = 1;
v = []; % eigenvalues
dx = 0.1; % bin size
```

Experiment

```
for i =1:K
    X =randn(N,M); % random NxN Gaussian matrix;
    A =X*X'/N; % Wishart
    v = [v; eig(A)]; % eigenvalues
end
```


Plot

```
alpha = M/N;
%[count, x]= hist(v, (1-sqrt(alpha))^2:dx:(1+sqrt(alpha))^2);
[count, x]= hist(v, 0:dx:6);
bar(x, count/(K*N*dx), 'y');
hold on ;
```



Theory

```
x = (1-sqrt(alpha))^2:0.01:(1+sqrt(alpha))^2;
plot(x,sqrt(4*alpha-(x-1-alpha).^2)./(2*pi*x), 'LineWidth', 2);
axis([-0.6 6 0 0.5]);
grid on;
title('N=2000, \alpha=2, K=1');
xlabel('\lambda');
ylabel('p(\lambda)');
```



Published with MATLAB® R2018a