

Confidence intervals vs Bayesian statistics

Last updated: October 20, 2017

Recap: hypothesis testing

Hypothesis testing = statistical proof by contradiction

- ▶ There is an unknown population parameter θ . (eg: $\mu_{MA} - \mu_{BA}$)
- ▶ H_0 : θ = (some specific value, eg, 0)
- ▶ H_A : not H_0 .
- ▶ Observe evidence s

$$\text{p-value} = \mathbb{P}(\text{Evidence} | \text{Hypothesis}).$$

Recap: hypothesis testing

What are the population parameters θ in the following hypothesis tests?

- ▶ Linear regression: is β_0 significantly different from 0?
- ▶ t -test: is the mean equal to μ_0 ?
- ▶ z -test: is the proportion equal to p_0 ?
- ▶ Two-sample t -test: are the means of the two groups equal?
- ▶ F -test: are the means of all groups equal?

Recap: hypothesis testing


What are the population parameters θ in the following hypothesis tests?

- ▶ Linear regression: is β_0 significantly different from 0?
- ▶ t -test: is the mean equal to μ_0 ?
- ▶ z -test: is the proportion equal to p_0 ?
- ▶ Two-sample t -test: are the means of the two groups equal?
- ▶ F -test: are the means of all groups equal?

Another way to think of p -values: confidence intervals.

Confidence intervals

Definition. A 95% confidence interval for a population parameter θ is a RANDOM interval, constructed based on the data, such that there is a 95% chance that this interval contains θ .



See os3.pdf, section 4.2. **Give interpretations for the following**

- ▶ Confidence intervals for estimates of β_0 and β_1 in linear regression (eg: teacher: salary vs degree)
- ▶ Confidence interval in z-test (eg: UT Austin survey)

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.
- ▶ **A 'maybe' answer?** (eg: there is an 80% chance that $\mu_{MA} \neq \mu_{BA}$)

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.
- ▶ **A 'maybe' answer?** (eg: there is an 80% chance that $\mu_{MA} \neq \mu_{BA}$)
Nonsense. μ_{MA} and μ_{BA} are fixed, so either they are different, or they are not.

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.
- ▶ **A 'maybe' answer?** (eg: there is an 80% chance that $\mu_{MA} \neq \mu_{BA}$)
Nonsense. μ_{MA} and μ_{BA} are fixed, so either they are different, or they are not.
BUT, you CAN construct a RANDOM interval and say: there is a 95% chance that this interval contains $\mu_{MA} - \mu_{BA}$. This is the **confidence interval**.

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.
- ▶ **A 'maybe' answer?** (eg: there is an 80% chance that $\mu_{MA} \neq \mu_{BA}$)
Nonsense. μ_{MA} and μ_{BA} are fixed, so either they are different, or they are not.
BUT, you CAN construct a RANDOM interval and say: there is a 95% chance that this interval contains $\mu_{MA} - \mu_{BA}$. This is the **confidence interval**.
- ▶ **A whole range of nulls with different certainties?** (eg: $\mu_{MA} - \mu_{BA}$ is uniformly distributed between -500 and 2000)

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.
- ▶ **A 'maybe' answer?** (eg: there is an 80% chance that $\mu_{MA} \neq \mu_{BA}$)
Nonsense. μ_{MA} and μ_{BA} are fixed, so either they are different, or they are not.
BUT, you CAN construct a RANDOM interval and say: there is a 95% chance that this interval contains $\mu_{MA} - \mu_{BA}$. This is the **confidence interval**.
- ▶ **A whole range of nulls with different certainties?** (eg: $\mu_{MA} - \mu_{BA}$ is uniformly distributed between -500 and 2000)
Nonsense. $\mu_{MA} - \mu_{BA}$ is a fixed number so it cannot be 'uniformly distributed'

Recap: hypothesis testing

Example to keep in mind: teacher: salary vs degree (MA/BA).

What to do if we want...

- ▶ **A quantitative answer?** (eg: how much is $\mu_{MA} - \mu_{BA}$?).
Regression.
- ▶ **A 'maybe' answer?** (eg: there is an 80% chance that $\mu_{MA} \neq \mu_{BA}$)
Nonsense. μ_{MA} and μ_{BA} are fixed, so either they are different, or they are not.
BUT, you CAN construct a RANDOM interval and say: there is a 95% chance that this interval contains $\mu_{MA} - \mu_{BA}$. This is the **confidence interval**.
- ▶ **A whole range of nulls with different certainties?** (eg: $\mu_{MA} - \mu_{BA}$ is uniformly distributed between -500 and 2000)
Nonsense. $\mu_{MA} - \mu_{BA}$ is a fixed number so it cannot be 'uniformly distributed'

To say 'maybe', you need: **Bayesian statistics**

Bayesian = statistics for those who like to say "maybe"

Best explained in: https://ocw.mit.edu/courses/mathematics/18-05-introduction-to-probability-and-statistics-spring-2014/readings/MIT18_05S14_Reading20.pdf.

The Frequentist view:

$$p\text{-value} = \mathbb{P}(\textit{Evidence} | \textit{Hypothesis})$$

The Bayesian view

The Posterior	The Evidence	The Prior
	The probability of getting this evidence if this hypothesis were true	The probability of H being true, before gathering evidence
$P(H E)$	$P(H E)$	$P(H)$
The probability that the hypothesis (H) is true given the evidence (E)	$P(E)$	The marginal probability of the evidence (Prob of E over all possibilities)

Bayesian vs Frequentist example

Frequentist

- ▶ $\mu_1 - \mu_2$ is an unknown fixed number.
- ▶ Prior 'belief' H_0 : $\mu_1 - \mu_2 = 0$
- ▶ Observe data $S = s$.
Compute

$$p\text{-value} = \mathbb{P}(S > s | H_0).$$

- ▶ After calculations: new 'belief': $\mu_1 - \mu_2 = 0$ or $\mu_1 - \mu_2 \neq 0$.

Bayesian:

- ▶ $\mu_1 - \mu_2$ is random
- ▶ Prior belief: $\mu_1 - \mu_2 \sim P_0$
- ▶ Observe data $S = s$. Update the belief:

The Posterior	The Evidence	The Prior
	The probability of getting this evidence if this hypothesis were true	The probability of H being true, before gathering evidence
$P(H E) = \frac{P(H E)P(H)}{P(E)}$		
The probability that the hypothesis (H) is true given the evidence (E)	The marginal probability of the evidence (Prob of E over all possibilities)	

- ▶ After calculations: posterior belief: $\mu_1 - \mu_2 \sim P_1$.

The Bayesian philosophy

What we think about the world after seeing data =

What we thought about the world before seeing data \times

Chance we'd see our data under different assumptions about the world

$$Pr(world|data) = Pr(world) \times \frac{Pr(data|world)}{Pr(data)}.$$

$$Posterior = Prior \times Likelihood$$

Why Bayesian?

Why NOT Bayesian?

The Bayesian philosophy

What we think about the world after seeing data =

What we thought about the world before seeing data \times

Chance we'd see our data under different assumptions about the world

$$Pr(world|data) = Pr(world) \times \frac{Pr(data|world)}{Pr(data)}.$$

$$Posterior = Prior \times Likelihood$$

Why Bayesian?

- ▶ Intuitive logic
- ▶ Flexible: input = a distribution, output = a distribution (instead of a single estimate or yes/no)
- ▶ Data-driven: see enough data and the prior does not matter.

Why NOT Bayesian?

The Bayesian philosophy

What we think about the world after seeing data =

What we thought about the world before seeing data \times

Chance we'd see our data under different assumptions about the world

$$Pr(world|data) = Pr(world) \times \frac{Pr(data|world)}{Pr(data)}.$$

$$Posterior = Prior \times Likelihood$$

Why Bayesian?

- ▶ Intuitive logic
- ▶ Flexible: input = a distribution, output = a distribution (instead of a single estimate or yes/no)
- ▶ Data-driven: see enough data and the prior does not matter.

Why NOT Bayesian?

- ▶ Answers depend on prior, which can be different for different people
- ▶ Likelihood hard to compute: resort to simulations or certain family of priors. So actually, not THAT flexible.

How to be Bayesian?

There is a Bayesian equivalent of regression, t-test, z-test etc.
Doable in R. See R code example: proportion estimate.