

Image Formation and Cameras

Computer Vision I
CSE 252A
Lecture 3

CS252A, Fall 2018

Computer Vision I

Announcements

- Assignment 0 is due on Tuesday
- Read Chapters 1 & 2 of Forsyth & Ponce
- Lecture notes on web page
- Waitlist
- Two slide decks that might be useful:
 - [Linear Algebra Review](#)
 - [Probability and Random Variables Review](#)
- (Subset of?) Final exam can be used for CSE MS Comprehensive Exam [Pending]

CS252A, Fall 2018

Computer Vision I

The course

- Part 1: The physics of imaging
- Part 2: Early vision
- Part 3: Reconstruction
- Part 4: Recognition

Image Formation: Outline

- Factors in producing images
- Projection
- Perspective/Orthographic Projection
- Vanishing points
- Projective Geometry
- Rigid Transformation and $SO(3)$
- Lenses
- Sensors
- Quantization/Resolution
- Illumination
- Reflectance and Radiometry

Earliest Surviving Photograph



- First photograph on record, “la table service” by Nicéphore Niépce in 1822.
- Note: First photograph by Niépce was in 1816.

CS252A, Fall 2018

Computer Vision I

Compare to Paintings



Willem Kalf, Mid 1600's





Pedro Campos,

CS252A, Fall 2018

Computer Vision I

How Cameras Produce Images

- Basic process:
 - photons hit a detector
 - the detector becomes charged
 - the charge is read out as brightness
- Sensor types:
 - CCD (charge-coupled device)
 - high sensitivity
 - high power
 - cannot be individually addressed
 - blooming
 - CMOS
 - simple to fabricate (cheap)
 - lower sensitivity, lower power
 - can be individually addressed



CS252A, Fall 2018

Computer Vision I

Images are two-dimensional patterns of brightness values.




Figure from US Navy Manual of Basic Optics and Optical Instruments, prepared by Bureau of Naval Personnel. Reprinted by Dover Publications, Inc., 1969.

CS252A, Fall 2018

They are formed by the projection of 3D objects. Computer Vision I


Effect of Lighting: Monet



CS252A, Fall 2018

Computer Vision I

Change of Viewpoint: Monet




Haystack at Chailly at sunrise (1865)

CS252A, Fall 2018

Computer Vision I

Pinhole Camera: Perspective projection

- Abstract camera model - box with a small hole in it




CS252A, Fall 2018 Forsyth&Ponce

Computer Vision I

Camera Obscura

illum in tabula per radios Solis, quam in celo contin-
git: hoc est, si in celo superior pars deliquum patiatur, in
radiis apparet inferior deficere, ut ratio exigit optica.



Sic nos exacte Anno .1544. Louanii eclipsim Solis
obseruauimus, inuenimusq; deficere paulo plus q; dex-


"When images of illuminated objects ... penetrate through a small hole into a very dark room ... you will see [on the opposite wall] these objects in their proper form and color, reduced in size ... in a reversed position, owing to the intersection of the rays". --- Leonardo Da Vinci

http://www.acmi.net.au/AIC/CAMERA_OBSCURA.html (Russell Naughton)

CS252A, Fall 2018

Computer Vision I

Camera Obscura



- Used to observe eclipses (e.g., Bacon, 1214-1294)
- By artists (e.g., Vermeer).

CS252A, Fall 2018

Computer Vision I

Camera Obscura



Jetty at Margate England, 1898.




<http://brightbytes.com/cosite/collection2.html> (Jack and Beverly Wilgus)

CS252A, Fall 2018

Computer Vision I


Purely Geometric View of Perspective




The projection of the point \mathbf{P} on the image plane Π' is given by the point of intersection \mathbf{P}' of the ray defined by \mathbf{PO} with the plane Π' .

Geometric properties of projection

- 3-D points map to points
- 3-D lines map to lines
- Planes map to whole image or half-plane
- Polygons map to polygons
- Important point to note: Angles & distances not preserved, nor are inequalities of angles & distances.
- Nor are ratios of angles or distances
- Degenerate cases:
 - line through focal point project to point
 - plane through focal point projects to a line



Equation of Perspective Projection




Cartesian coordinates:

- We have, by similar triangles, that for $\mathbf{P}=(x, y, z)$, the intersection of OP with Π' is $(f' x/z, f' y/z, f')$
- Establishing an image plane coordinate system at C' aligned with i and j , we get $(x, y, z) \rightarrow (f' \frac{x}{z}, f' \frac{y}{z})$

CS252A, Fall 2018

Computer Vision I

Virtual Image Plane



- Virtual image plane in front of optical center.
- Image is ‘upright’

$$(x, y, z) \rightarrow (f'' \frac{x}{z}, f'' \frac{y}{z})$$


CS252A, Fall 2018

Computer Vision I


A Digression

Projective Geometry
and
Homogenous Coordinates

What is the intersection of
two lines in a plane?



Do two lines in the plane always intersect at a point?




No, Parallel lines don't meet at a point.

CS252A, Fall 2018

Computer Vision I

Can the perspective image of two parallel lines meet at a point?

YES



CS252A, Fall 2018

Computer Vision I

Projective geometry provides an elegant means for handling these different situations in a unified way and **homogenous coordinates** are a way to represent entities (points & lines) in projective spaces.

Projective Geometry

- Axioms of Projective Plane
 1. Every two distinct points define a line
 2. Every two distinct lines define a point (intersect at a point)
 3. There exists three points, A,B,C such that C does not lie on the line defined by A and B.
- Different than Euclidean (affine) geometry
- Projective plane is “bigger” than affine plane – includes “line at infinity”

$$\text{Projective Plane} = \text{Affine Plane} + \text{Line at Infinity}$$

Homogenous coordinates

A way to represent points in a projective space

- Use three numbers to represent a point on a projective plane

Why? The projective plane has to be bigger than the Cartesian plane so that it contains the point where parallel lines intersect.

How: Add an extra coordinate
e.g., $(x,y) \rightarrow (x,y,1)$

Impose equivalence relation
 $(x,y,z) \approx \lambda*(x,y,z)$
such that $\lambda \neq 0$
i.e., $(x,y,1) \approx (\lambda x, \lambda y, \lambda)$

- Points at infinity – zero for last coordinate
e.g., $(x,y,0)$

- Why do this?

- Possible to represent points “at infinity”
 - Where parallel lines intersect
 - Where parallel planes intersect
- Possible to write the action of a perspective camera as a matrix


Homogenous coordinates

A way to represent points in a projective space

Use three numbers to represent a point on a projective plane

Add an extra coordinate
e.g., $(x,y) \rightarrow (x,y,1)$

Impose equivalence relation
 $(x,y,z) \approx \lambda*(x,y,z)$
such that (λ not 0)
i.e., $(x,y,1) \approx (\lambda x, \lambda y, \lambda)$



Conversion


Euclidean \rightarrow Homogenous \rightarrow Euclidean

In 2-D

- Euclidean \rightarrow Homogenous:
 $(x, y) \rightarrow \lambda (x, y, 1)$
- Homogenous \rightarrow Euclidean:
 $(x, y, z) \rightarrow (x/z, y/z)$

In 3-D

- Euclidean \rightarrow Homogenous:
 $(x, y, z) \rightarrow \lambda (x, y, z, 1)$
- Homogenous \rightarrow Euclidean:
 $(x, y, z, w) \rightarrow (x/w, y/w, z/w)$




CS252A, Fall 2018

Computer Vision I

Points at infinity

Point at infinity – last coordinate is zero $(x, y, 0)$
 and equivalence relation
 $(x, y, 0) \approx \lambda^*(x, y, 0)$

No corresponding Euclidean point (you'd divide by zero).



$$\text{Projective Plane} = \text{Affine Plane} + \text{Line at Infinity}$$

CS252A, Fall 2018

Computer Vision I

Projective transformation

- 3×3 linear transformation of homogenous coordinates


$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{21} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

- Points map to points
- Lines map to lines
- Points in X can map to points at infinity in U . e.g.,
 $u_3=0$

CS252A, Fall 2018

Computer Vision I

The equation of projection



Cartesian coordinates:

$$(x, y, z) \rightarrow (f \frac{x}{z}, f \frac{y}{z})$$

$$\begin{pmatrix} U \\ V \\ W \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1/f & 0 \end{pmatrix} \begin{pmatrix} X \\ Y \\ Z \\ 1 \end{pmatrix}$$

CS252A, Fall 2018

Computer Vision I

End of the Digression

CS252A, Fall 2018

Computer Vision I

CS252A, Fall 2018

Computer Vision I

In a perspective image, parallel lines meet at a point, called the vanishing point




CS252A, Fall 2018

Doesn't need to be near the center of the image

Computer Vision I

Parallel lines meet in the image




- Vanishing point location: Intersection of 3-D line through O parallel to given line(s)
- A single line can have a vanishing point


CS252A, Fall 2018

Computer Vision I

Vanishing points




- A scene can have more than one vanishing point
- Different 3-D directions correspond different vanishing points



CS252A, Fall 2018

Computer Vision I

Vanishing Points



CS252A, Fall 2018

Computer Vision I


Vanishing Point

- In the **projective plane**, parallel lines meet at a point at infinity.
- The vanishing point is the perspective projection of that point at infinity, resulting from multiplication by the camera matrix.

What is a Camera?

- An mathematical expression that relates points in 3D to points in an image for different types of physical cameras or imaging situations

Simplified Camera Models




CS252A, Fall 2018

Computer Vision I

Affine Camera Model

Appropriate for
in neighborhood
of (x_0, y_0, z_0)



- Take perspective projection equation, and perform Taylor series expansion about some point (x_0, y_0, z_0)
- Drop terms that are higher order than linear
- Resulting expression is the Affine Camera Model

CS252A, Fall 2018

Computer Vision I

- Perspective

$$\begin{bmatrix} u \\ v \end{bmatrix} = \frac{f}{z} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Perform a Taylor series expansion about (x_0, y_0, z_0)

$$\begin{bmatrix} u \\ v \end{bmatrix} \approx \frac{f}{z_0} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} - \frac{f}{z_0^2} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} (z - z_0) + \frac{f}{z_0} \begin{bmatrix} 1 \\ 0 \end{bmatrix} (x - x_0) \\ + \frac{f}{z_0} \begin{bmatrix} 0 \\ 1 \end{bmatrix} (y - y_0) + \frac{f}{2} \frac{2}{z_0^3} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} (z - z_0)^2 + \dots$$

- Dropping higher order terms and regrouping.

$$\begin{bmatrix} u \\ v \end{bmatrix} \approx \frac{f}{z_0} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} + \begin{bmatrix} f/z_0 & 0 & -fx_0/z_0^2 \\ 0 & f/z_0 & -fy_0/z_0^2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{Ap} + \mathbf{b}$$

Affine camera model


$$\begin{bmatrix} u \\ v \end{bmatrix} \approx \frac{f}{z_0} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} + \begin{bmatrix} f/z_0 & 0 & -fx_0/z_0^2 \\ 0 & f/z_0 & -fy_0/z_0^2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \mathbf{Ap} + \mathbf{b}$$

Rewrite affine camera model
in terms of Homogenous Coordinates

$$\begin{bmatrix} u \\ v \\ w \end{bmatrix} \approx \begin{bmatrix} f/z_0 & 0 & -fx_0/z_0^2 & fx_0/z_0 \\ 0 & f/z_0 & -fy_0/z_0^2 & fy_0/z_0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

Scaled orthographic projection

Starting with Affine Camera Model, take Taylor series about $(x_0, y_0, z_0) = (0, 0, z_0)$ – a point on the optical axis



f/z_0 is the scale

CS252A, Fall 2018

Computer Vision I

The projection matrix for scaled orthographic projection

$$\begin{bmatrix} u \\ v \\ w \end{bmatrix} = \begin{bmatrix} f/z_0 & 0 & 0 & 0 \\ 0 & f/z_0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ t \end{bmatrix}$$

- Parallel lines project to parallel lines
- Ratios of distances are preserved under orthographic projection


CS252A, Fall 2018

Computer Vision I

Other camera models

- Generalized camera – maps points lying on rays and maps them to points on the image plane.

Omnicam (hemispherical)



CS252A, Fall 2018

Light Probe (spherical)




Computer Vision I

Some Alternative “Cameras”



CS252A, Fall 2018



Computer Vision I