

### Problem Set B

Aim: The purpose of this Problem Set is to reinforce your understanding of under which circumstances a unique solution exists.

1. On a convex polygonal domain  $\Omega$ , consider the differential equation

$$-\Delta u(\mathbf{x}) = f, \quad \mathbf{x} \in \Omega \quad (1)$$

$$\hat{n} \cdot \nabla u(\mathbf{x}) = g, \quad \mathbf{x} \in \partial\Omega \quad (2)$$

where  $f$  and  $g$  are sufficiently regular functions that are given.

- (a) Why is this problem **not** well-posed?
  - (b) If you would implement a standard finite element method for this problem (for example cG(1)<sup>1</sup>), would you be able to solve the linear system? What happens?
2. Consider the following Boundary Value Problem (BVP):

$$-(k(x)u'(x))' + u'(x) = f(x), \quad x \in \Omega \quad (3)$$

$$u = 0, \quad x \in \partial\Omega \quad (4)$$

with the coefficient,  $k(x)$ , being a continuous function such that  $k(x) \geq k_0 \geq 0$  for  $x \in \Omega = (0, 1)$ . Assume both  $k(x)$  and  $f(x)$  are given with  $f \in L^1(0, 1)$ .

- (a) Derive a variational formulation for the BVP: Find  $u \in V$  such that

$$a(u, v) = L(v), \quad \forall v \in V.$$

- i. Define the Hilbert Space  $V = \mathcal{H}_0^1(\Omega)$ .
  - ii. Define the bilinear form  $a : V \times V \rightarrow \mathbb{R}$  and linear form  $L : V \rightarrow \mathbb{R}$ .
- (b) Show that for  $\|w\|_E^2 = a(w, w)$  we have that

$$\|w\|_E^2 = \int_{\Omega} k(x)(w'(x))^2 dx$$

and that this represents a norm on  $E$ .

- (c) Prove that there exists a unique solution to the variational problem in 2a.
- (d) Formulate the cG(1) method for the given BVP on a subdivision of  $(0, 1)$  with mesh size  $h = h(x)$ .
- (e) Prove the following a priori error estimate:

$$\|u - u_h\|_E \leq C \|hu''\|_{L^2(0,1)}.$$

Here  $u$  is the exact solution to the given BVP and  $u_h$  is the approximation of  $u$  using the cG(1) method.

---

<sup>1</sup>FEM with a continuous piecewise linear approximation  $u_h$

- (f) Describe the steps in an adaptive algorithm for local mesh refinement based on a posteriori error estimation and given a tolerance TOL on  $\|u - u_h\|_E$ . Assume TOL is real. Why is adaptivity important?

3. Show that a spatial discretization of the equation

$$\begin{aligned} \dot{u} + \Delta u + u &= f, & \mathbf{x} \in \Omega \\ u &= 0, & \mathbf{x} \in \partial\Omega, t > 0 \\ u &= u_0, & \mathbf{x} \in \Omega, t = 0 \end{aligned}$$

leads to the algebraic system

$$M\ddot{\xi}(t) + A\xi(t) + M\xi(t) = b(t).$$

Identify the entries of the involved matrices and vectors.<sup>2</sup>

---

<sup>2</sup>Exercise 5.3 in M. G. Larson and F. Bengzon, *The Finite Element Method: Theory, Implementation, and Applications*