

On the Rankin Constants of Lattices


F. Oggier

WCC 2026

Cryptography


Suppose $\mathbf{v}_1, \dots, \mathbf{v}_m$ are linearly independent vectors in $\mathbb{R}^n$ ($m \leq n$). A **lattice** $L \subset \mathbb{R}^n$, generated by $\mathbf{v}_1, \dots, \mathbf{v}_m$, is the additive subgroup of $\mathbb{R}^n$ generated by $\mathbf{v}_1, \dots, \mathbf{v}_m$:

$$L = \{\mathbf{u}M, \mathbf{u} \in \mathbb{Z}^m\}, \quad M = \begin{bmatrix} \mathbf{v}_1 \\ \vdots \\ \mathbf{v}_m \end{bmatrix}, \quad G = MM^T.$$



$$M = \begin{bmatrix} 1 & 0 \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}, \quad G = \begin{bmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{bmatrix}.$$

The matrix M containing the basis vectors is called a **generator matrix**, the matrix $G = MM^T$ is called a **Gram matrix** (it contains the “geometry” of the lattice, that is the norms of the basis vectors on its diagonal, and their inner product or angle elsewhere).



$$M = \begin{bmatrix} 1 & 0 \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}, \quad G = \begin{bmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{bmatrix}.$$

✓ Lattices

Coding Theory

Cryptography

- A vector $\mathbf{x} = (x_1, \dots, x_k)$ of information symbols is encoded linearly into a codeword $\mathbf{c} = (c_1, \dots, c_n)$:

$$\mathbf{x} = (x_1, \dots, x_k) \mapsto \mathbf{c} = (c_1, \dots, c_n), n \geq k,$$

both have coefficients in a given alphabet, typically a finite field $\mathbb{F}_q$ or the finite ring $\mathbb{Z}_q = \{0, 1, \dots, q-1\}$ of integers modulo q (if $q = p$ is a prime, $\mathbb{Z}_p = \mathbb{F}_p$).

- The set of all codewords, denoted by C , is called a **linear code** of **length** n .

If the alphabet is $\mathbb{F}_q$, C forms a linear subspace of $\mathbb{F}_q^n$. It thus has a **dimension**, namely k , and we say that we consider an (n, k) linear code C over $\mathbb{F}_q$.

- Since an (n, k) linear code C is a subspace, it has a basis, containing k vectors, that we stack as rows of a $k \times n$ matrix G , called a **generator matrix** for C .
- There is a unique generator matrix of the form $G = [\mathbf{I}_k | A]$ where $\mathbf{I}_k$ is the identity matrix. The code is said to be in **systematic form**.

$$\underbrace{(x_1, \dots, x_k)}_{\text{information data}} \underbrace{\begin{bmatrix} & a_{11} & a_{1,n-k} \\ \mathbf{I}_k & \vdots & \vdots \\ & a_{k,1} & a_{k,n-k} \end{bmatrix}}_{\text{generator matrix } G} = \underbrace{(x_1, \dots, x_k, c_{k+1}, \dots, c_n)}_{\text{codeword}}$$

✓ Lattices

✓ Coding Theory

Cryptography

Let

$$\rho : \mathbb{Z} \rightarrow \mathbb{Z}_q = \{0, 1, \dots, q-1\}, \quad x \mapsto x \pmod{q},$$

be the map of reduction modulo q . Given $a \pmod{q}$, its pre-image $\rho^{-1}(a)$ is the set of integers that are mapped to a by ρ , that is $\rho^{-1}(a) = \{a + bq, b \in \mathbb{Z}\}$.




The pre-image $\rho^{-1}(a) \subset \mathbb{Z}$ of $a \pmod{5}$.

Consider

$$\rho: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}_q \times \mathbb{Z}_q, (x_1, x_2) \mapsto (x_1 \pmod{q}, x_2 \pmod{q}).$$


The pre-image $\rho^{-1}((a_1, a_2))$ is now the set of 2-dimensional vectors with integer entries that is mapped to (a_1, a_2) by ρ .




The pre-image $\rho^{-1}((a_1, a_2)) \subset \mathbb{Z}^2$ of $(a_1 \pmod{3}, a_2 \pmod{3})$.

$$\rho : \mathbb{Z}^n \rightarrow \mathbb{Z}_q^n, \mathbf{x} \mapsto \rho(\mathbf{x}).$$

One may take any arbitrary subset S of $\mathbb{Z}_q^n$ and compute $\rho^{-1}(S)$, but what if S has a structure? how is it carried over to $\rho^{-1}(S)$?



The pre-image $\rho^{-1}(S) \subset \mathbb{Z}^2$ of $S = \{(a_1 \pmod{3}, a_2 \pmod{3}), (a_1 + 1 \pmod{3}, a_2 + 1 \pmod{3})\}$.



The pre-image $\rho^{-1}(C) \subset \mathbb{Z}^2$ of the linear binary code $C = \{(0, 0), (1, 1)\}$.

A well known connection between linear codes in $\mathbb{Z}_q^n$ and lattices:
 Given a subset $S \subset \mathbb{Z}_q^n$ then $\rho^{-1}(S)$ is a lattice in $\mathbb{R}^n$ if and only if
 S is a linear code in $\mathbb{Z}_q^n$.

A linear code C in $\mathbb{Z}_q^n$ is
 an additive subgroup of
 $\mathbb{Z}_q^n$.

A lattice $L \subset \mathbb{R}^n$ is an
 additive subgroup of $\mathbb{R}^n$
 given by

$$L = \{uM, u \in \mathbb{Z}^m\}$$

for M a generator matrix,
 whose rows are $m \leq n$
 linearly independent vec-
 tors.

Construction A

Let C be a linear code in $\mathbb{Z}_q^n$, the integers modulo a positive integer $q \geq 2$, where q is either prime or composite. Let $\rho : \mathbb{Z}^n \rightarrow \mathbb{Z}_q^n$ be the componentwise reduction modulo q . Then the lattice $\Lambda_C = \rho^{-1}(C)$ is said to have been obtained via *Construction A*.

- Sometimes $\Lambda_C = \frac{1}{\sqrt{q}}\rho^{-1}(C)$ is called Construction A.
- There are other constructions, Construction B,C,D...
- The lattice Λ_C is also known as a q -ary lattice or modulo q lattice (especially in the context of lattice-based cryptography).

Let L be a lattice. Suppose d is the smallest distance between any two distinct points of L , the vectors in L with length d are the shortest vectors of L .

- The minimum norm of L , or the norm of L or the minimum of L , is the length of a shortest nonzero lattice vector of L :

$$N(L) = \min(L) = \inf_{\mathbf{x} \in L, \mathbf{x} \neq 0} N(\mathbf{x}),$$

where the norm of $\mathbf{x}$ is $N(\mathbf{x}) = \mathbf{x} \cdot \mathbf{x} = (\|\mathbf{x}\|_2)^2$.

- For $L = \mathbb{Z}^n$, $N(L) = 1$.
- Let L be a lattice of rank n , then

$$\gamma(L) = \frac{N(L)}{\sqrt[n]{\det(L)}}, \quad \det(L) = \det(G),$$

which is invariant by scaling.

- A sublattice $L' \leq L$ of rank l is a lattice included in L , generated by l basis vectors.
- Set

$$d_l(L) = \min_{\substack{L' \leq L, \\ \text{rank}(L')=l}} \det(L'), \quad 1 \leq l \leq n.$$

- We define

$$\begin{aligned} \gamma_{n,l}(L) &= d_l(L) \det(L)^{-l/n} \\ \gamma_{n,l} &= \sup_{\text{rank}(L)=n} \gamma_{n,l}(L). \end{aligned}$$

The constant $\gamma_{n,l}$ is called the **Rankin constant**, and when $l = 1$, it translates into the Hermite constant: $\gamma_{n,1} = \gamma_n$.

$$\gamma_{n,2} \geq \frac{3}{4^{2/n}}, \quad n \geq 3.$$

$$\begin{bmatrix} \mathbf{I}_k & A \end{bmatrix},$$

the generator matrix of C in systematic form, then a generator matrix of Λ_C is

$$\begin{bmatrix} \mathbf{I}_k & A \\ \mathbf{0} & q\mathbf{I}_{n-k} \end{bmatrix}.$$

For $n \geq 2$, $C = \{(a_1, \dots, a_{n-1}, \sum_{i=1}^{n-1} a_i), a_1, \dots, a_{n-1} \in \mathbb{F}_2\}$ has length n and dimension $n - 1$. A systematic generator is

$$\begin{bmatrix} \mathbf{I}_{n-1} & \mathbf{1} \end{bmatrix}.$$

A generator matrix for Λ_C is thus

$$\begin{bmatrix} \mathbf{I}_{n-1} & \mathbf{1} \\ \mathbf{0} & 2 \end{bmatrix}.$$

We have just constructed the lattice

$$D_n = \{(x_1, \dots, x_n), \sum_{i=1}^n x_i \text{ is even}\}.$$

Let $\mathcal{R}(r, m)$ denote the binary Reed-Mueller code of order r and length 2^m . A recursive formula for a generator matrix of $\mathcal{R}(r, m)$ is

$$\mathbf{B}_{r,m} = \begin{bmatrix} \mathbf{B}_{r,m-1} & \mathbf{B}_{r,m-1} \\ \mathbf{0} & \mathbf{B}_{r-1,m-1} \end{bmatrix},$$

with $\mathbf{B}_{0,i} = \mathbf{1}_{1,2^i}$ and $\mathbf{B}_{i,i} = I_{2^i}$ for $i \leq m$.

- The dimension k of this code is $k = \sum_{i=0}^r \binom{m}{i}$.
- We have $\mathcal{R}(i, m) \subseteq \mathcal{R}(j, m)$ for $i \leq j$.
- Given the lattice $\Lambda_{\mathcal{R}(j, m)}$, a natural sublattice is generated by $\mathbf{B}_{i, m}$, a generator matrix of $\mathcal{R}(i, m)$, which we will denote by $\Lambda(\mathbf{B}_{i, m})$.

The Case $r = 1$

1. For $l = 1$,

$$\gamma_{n,1}(\Lambda_{\mathcal{R}(1,m)}) = \begin{cases} \frac{2^{3/2}}{2} = \sqrt{2} & m = 2 \\ 2^{2(m+1)/2^m} & m \geq 3, \end{cases}$$

which is optimal for $m = 2, 3$ (that is in dimension 4 and 8).

2. For $l = 2$, there exists a sublattice $\Lambda(\mathbf{B})$ such that

$$\gamma(\Lambda_{\mathcal{R}(1,m)}, \Lambda(\mathbf{B})) = \begin{cases} 3 & m = 3 \ (n = 8) \\ \frac{1}{2\sqrt{2}} & m = 4 \ (n = 16) \end{cases}$$

and $\gamma_{8,2}(\Lambda_{\mathcal{R}(1,3)}) = 3$, which is optimal in dimension 8.

3. For $l \geq 3$, there exists a sublattice $\Lambda(\mathbf{B})$ such that

$$\gamma(\Lambda_{\mathcal{R}(1,m)}, \Lambda(\mathbf{B})) = \frac{4 \cdot 2^{(m-2)l}}{\left(\frac{2^{2m}}{2^{m+1}}\right)^{2l/n}}$$

for $m \leq \frac{2l-2}{l} + 2$. This quantity simplifies to 4 when $m = 3$ and $l = 3, 4$.

About $\gamma'_{n,l}(\Lambda_C)$

- We recall that $\gamma'_{n,I}(L) = \sqrt{d_I(L)d_I(L^*)}$ where L^* is the dual lattice of L .
- For speaking of duality, it is often preferred to work with $\Lambda_C = \frac{1}{\sqrt{q}}\rho^{-1}(C)$ (we will see why next).
- The results hold for $\mathbb{F}_q$ (though we consider $q = p$).

$$C^\perp = \{\mathbf{v} \in \mathbb{F}_q^n, \mathbf{c} \cdot \mathbf{v} = 0 \text{ for all } \mathbf{c} \in C\} \quad \Bigg| \quad L^* = \{\mathbf{y} \in \mathbb{R}^n, \mathbf{y} \cdot \mathbf{x} \in \mathbb{Z} \text{ for all } \mathbf{x} \in L\}$$

More on $\gamma_{5,2}$

We have $\gamma_{n,2} \geq \sqrt{3}$ for $n \geq 5$ so

$$1.7321 \leq \gamma_{5,2} \leq 2.$$

- We use $\gamma'_{n,l} \leq \gamma_{n,l}$ for the lower bounds.

Natural correspondence between codes and lattices:

dual code

self-dual code

weight enumerator

?

dual lattice

unimodular lattice

theta series

Rankin constants

- Known lattices:
 - Give known optimal constructions in small dimensions but still valid bounds for large dimensions.
- Bounds on $\gamma_{n,2}$ and $\gamma'_{n,2}$.
 - Due to the ability to provide a lower bound, which give new bounds for $n = 5, 7$.
- Bounds on families of $\gamma_{n,l}(\Lambda_C)$.

Open Questions

- $\gamma_{5,2}$, $\gamma'_{5,2}$ and $\gamma_{7,2}$, $\gamma'_{7,2}$?
- Other families of codes?

