

## RESEARCH

# Chemotherapy weakly contributes to predicted neoantigen expression in ovarian cancer

Timothy O'Donnell<sup>1\*</sup>, Elizabeth L. Christie<sup>2</sup>, Arun Ahuja<sup>1</sup>, Jacqueline Buros<sup>1</sup>, B. Arman Aksoy<sup>1</sup>, David D. L. Bowtell<sup>2</sup>, Alexandra Snyder<sup>3†</sup> and Jeff Hammerbacher<sup>1†</sup>

\*Correspondence:

tim@hammerlab.org

<sup>1</sup>Icahn School of Medicine at Mount Sinai, New York, USA

Full list of author information is available at the end of the article

†Co-senior author

## Abstract

**Background:** Patients with highly mutated tumors, such as melanoma or smoking-related lung cancer, have higher rates of response to immune checkpoint blockade therapy, perhaps due to increased neoantigen expression. Many chemotherapies including platinum compounds are known to be mutagenic, but the impact of standard treatment protocols on mutational burden and resulting neoantigen expression in most human cancers is unknown.

**Methods:** We sought to quantify the effect of chemotherapy treatment on computationally predicted neoantigen expression for 12 high grade serous ovarian carcinoma (HGSC) patients with pre- and post-chemotherapy samples collected in the Australian Ovarian Cancer Study. We additionally analyzed 16 patients from the cohort with post-treatment samples only, including five primary surgical samples exposed to neoadjuvant chemotherapy. Our approach integrates tumor whole genome and RNA sequencing with class I MHC binding prediction and mutational signatures of chemotherapy exposure extracted from two preclinical studies.

**Results:** The mutational signatures for cisplatin and cyclophosphamide identified in a preclinical model had significant but inexact associations with the relevant exposure in the clinical samples. In an analysis stratified by tissue type (solid tumor or ascites), relapse samples collected after chemotherapy harbored a median of 90% more expressed neoantigens than untreated primary samples, a figure that combines the effects of chemotherapy and other mutagenic processes operative during relapse. Neoadjuvant-treated primary samples showed no detectable increase over untreated samples. The contribution from chemotherapy-associated signatures was small, accounting for a mean of 5% (range 0–16) of the expressed neoantigen burden in relapse samples. In both treated and untreated samples, most neoantigens were attributed to COSMIC *Signature (3)*, associated with BRCA disruption, *Signature (1)*, associated with a slow mutagenic process active in healthy tissue, and *Signature (8)*, of unknown etiology.

**Conclusion:** Relapsed HGSC tumors harbor nearly double the predicted expressed neoantigen burden of primary samples, but mutations associated with chemotherapy signatures account for only a small part of this increase. The mutagenic processes responsible for most neoantigens are similar between primary and relapse samples. Our analyses are based on mutations detectable from whole genome sequencing of bulk samples and do not account for neoantigens present in small populations of cells.

**Keywords:** neoantigen; mutational signature; chemotherapy

## Background

Many chemotherapies including platinum compounds [1], cyclophosphamide [2], and etoposide [3] exert their effect through DNA damage, and recent studies have found evidence for chemotherapy-induced mutations in post-treatment acute myeloid leukaemia [4], glioma [5], and esophageal adenocarcinoma [6]. Successful development of immune checkpoint-mediated therapy [7] has focused attention on the importance of T cell responses to somatic mutations in coding genes that generate neoantigens [8]. Studies based on bulk-sequencing of tumor samples followed by computational peptide-class I MHC affinity prediction [9] have suggested that tumors with more mutations and predicted mutant MHC I peptide ligands are more likely to respond to checkpoint blockade immunotherapy [10, 11]. Ovarian cancers fall into an intermediate group of solid tumors in terms of mutational load present in pre-treatment surgical samples [12]. However, the effect of standard chemotherapy regimes on tumor mutation burden and resulting neoantigen expression in ovarian cancer is poorly understood.

Investigators associated with the Australian Ovarian Cancer Study (AOCS) performed whole genome and RNA sequencing of 79 pre-treatment and 35 post-treatment cancer samples from 92 HGSC patients, including 12 patients with both pre- and post-treatment samples [13]. The samples were obtained from solid tissue resections, autopsies, and ascites drained to relieve abdominal distension. Treatment regimes varied but primary treatment always included platinum-based chemotherapy. In their analysis, Patch *et al.* reported that post-treatment samples harbored more somatic mutations than pre-treatment samples and exhibited evidence of chemotherapy-associated mutations. Here we extend these results by quantifying the mutations and predicted neoantigens attributable to chemotherapy-associated mutational signatures. We find that, while neoantigen expression increases after treatment and relapse, only a small part of the increase is due to mutations associated with chemotherapy signatures.

## Content

Text and results for this section, as per the individual journal's instructions for authors.

## Section title

Text for this section ...

### Sub-heading for section

Text for this sub-heading ...

#### *Sub-sub heading for section*

Text for this sub-sub-heading ...

*Sub-sub-sub heading for section* Text for this sub-sub-sub-heading ... In this section we examine the growth rate of the mean of  $Z_0$ ,  $Z_1$  and  $Z_2$ . In addition, we examine a common modeling assumption and note the importance of considering

the tails of the extinction time  $T_x$  in studies of escape dynamics. We will first consider the expected resistant population at  $vT_x$  for some  $v > 0$ , (and temporarily assume  $\alpha = 0$ )

$$E[Z_1(vT_x)] = E\left[\mu T_x \int_0^{v \wedge 1} Z_0(uT_x) \exp(\lambda_1 T_x(v-u)) du\right].$$

If we assume that sensitive cells follow a deterministic decay  $Z_0(t) = xe^{\lambda_0 t}$  and approximate their extinction time as  $T_x \approx -\frac{1}{\lambda_0} \log x$ , then we can heuristically estimate the expected value as

$$\begin{aligned} E[Z_1(vT_x)] &= \frac{\mu}{r} \log x \int_0^{v \wedge 1} x^{1-u} x^{(\lambda_1/r)(v-u)} du \\ &= \frac{\mu}{r} x^{1-\lambda_1/\lambda_0 v} \log x \int_0^{v \wedge 1} x^{-u(1+\lambda_1/r)} du \\ &= \frac{\mu}{\lambda_1 - \lambda_0} x^{1+\lambda_1/rv} \left(1 - \exp\left[-(v \wedge 1) \left(1 + \frac{\lambda_1}{r}\right) \log x\right]\right). \quad (1) \end{aligned}$$

Thus we observe that this expected value is finite for all  $v > 0$  (also see [14, ?, ?, ?, ?]).

#### Competing interests

The authors declare that they have no competing interests.

#### Author's contributions

Text for this section ...

#### Acknowledgements

Text for this section ...

#### Author details

<sup>1</sup>Icahn School of Medicine at Mount Sinai, New York, USA. <sup>2</sup>Peter MacCallum Cancer Centre, East Melbourne, Victoria 3002 Australia. <sup>3</sup>Department of Medicine, Memorial Sloan-Kettering Cancer Center, Weill Cornell Medical College, New York, USA.

#### References

- Hannan, M.A., Al-Dakan, A.A., Hussain, S.S., Amer, M.H.: Mutagenicity of cisplatin and carboplatin used alone and in combination with four other anticancer drugs. *Toxicology* **55**(1-2), 183–191 (1989). doi:10.1016/0300-483x(89)90185-6
- Anderson, D., Bishop, J.B., Garner, R.C., Ostrosky-Wegman, P., Selby, P.B.: Cyclophosphamide: Review of its mutagenicity for an assessment of potential germ cell risks. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **330**(1-2), 115–181 (1995). doi:10.1016/0027-5107(95)00039-1
- Nakanomyo, H., Hiraoka, M., Shiraya, M.: Mutagenicity tests of etoposide and teniposide. *J. Toxicol. Sci.* **11**(Supplement1), 301–310 (1986). doi:10.2131/jts.11.supplement1\_301
- Ding, L., Ley, T.J., Larson, D.E., Miller, C.A., Koboldt, D.C., Welch, J.S., Ritchey, J.K., Young, M.A., Lamprecht, T., McLellan, M.D., McMichael, J.F., Wallis, J.W., Lu, C., Shen, D., Harris, C.C., Dooling, D.J., Fulton, R.S., Fulton, L.L., Chen, K., Schmidt, H., Kalicki-Veizer, J., Magrini, V.J., Cook, L., McGrath, S.D., Vickery, T.L., Wendl, M.C., Heath, S., Watson, M.A., Link, D.C., Tomasson, M.H., Shannon, W.D., Payton, J.E., Kulkarni, S., Westervelt, P., Walter, M.J., Graubert, T.A., Mardis, E.R., Wilson, R.K., DiPersio, J.F.: Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* **481**(7382), 506–510 (2012). doi:10.1038/nature10738
- Johnson, B.E., Mazar, T., Hong, C., Barnes, M., Aihara, K., McLean, C.Y., Fouse, S.D., Yamamoto, S., Ueda, H., Tatsuno, K., Asthana, S., Jalbert, L.E., Nelson, S.J., Bollen, A.W., Gustafson, W.C., Charron, E., Weiss, W.A., Smirnov, I.V., Song, J.S., Olshen, A.B., Cha, S., Zhao, Y., Moore, R.A., Mungall, A.J., Jones, S.J.M., Hirst, M., Marra, M.A., Saito, N., Aburatani, H., Mukasa, A., Berger, M.S., Chang, S.M., Taylor, B.S., Costello, J.F.: Mutational Analysis Reveals the Origin and Therapy-Driven Evolution of Recurrent Glioma. *Science* **343**(6167), 189–193 (2013). doi:10.1126/science.1239947
- Murugaesu, N., Wilson, G.A., Birkbak, N.J., Watkins, T.B.K., McGranahan, N., Kumar, S., Abbassi-Ghadi, N., Salm, M., Mitter, R., Horswell, S., Rowan, A., Phillimore, B., Biggs, J., Begum, S., Matthews, N., Hochhauser, D., Hanna, G.B., Swanton, C.: Tracking the Genomic Evolution of Esophageal Adenocarcinoma through Neoadjuvant Chemotherapy. *Cancer Discovery* **5**(8), 821–831 (2015). doi:10.1158/2159-8290.cd-15-0412

7. Chen, D.S., Mellman, I.: Oncology Meets Immunology: The Cancer-Immunity Cycle. *Immunity* **39**(1), 1–10 (2013). doi:10.1016/j.immuni.2013.07.012

8. Schumacher, T.N., Schreiber, R.D.: Neoantigens in cancer immunotherapy. *Science* **348**(6230), 69–74 (2015). doi:10.1126/science.aaa4971

9. Lundegaard, C., Lund, O., Kesmir, C., Brunak, S., Nielsen, M.: Modeling the adaptive immune system: predictions and simulations. *Bioinformatics* **23**(24), 3265–3275 (2007). doi:10.1093/bioinformatics/btm471

10. Allen, E.M.V., Miao, D., Schilling, B., Shukla, S.A., Blank, C., Zimmer, L., Sucker, A., Hillen, U., Foppen, M.H.G., Goldinger, S.M., Utikal, J., Hassel, J.C., Weide, B., Kaehler, K.C., Loquai, C., Mohr, P., Gutzmer, R., Dummer, R., Gabriel, S., Wu, C.J., Schadendorf, D., Garraway, L.A.: Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science* **350**(6257), 207–211 (2015). doi:10.1126/science.aad0095

11. Rizvi, N.A., Hellmann, M.D., Snyder, A., Kvistborg, P., Makarov, V., Havel, J.J., Lee, W., Yuan, J., Wong, P., Ho, T.S., Miller, M.L., Rekhtman, N., Moreira, A.L., Ibrahim, F., Bruggeman, C., Gasmir, B., Zappasodi, R., Maeda, Y., Sander, C., Garon, E.B., Merghoub, T., Wolchok, J.D., Schumacher, T.N., Chan, T.A.: Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* **348**(6230), 124–128 (2015). doi:10.1126/science.aaa1348

12. Lawrence, M.S., Stojanov, P., Polak, P., Kryukov, G.V., Cibulskis, K., Sivachenko, A., Carter, S.L., Stewart, C., Mermel, C.H., Roberts, S.A., Kiezun, A., Hammerman, P.S., McKenna, A., Drier, Y., Zou, L., Ramos, A.H., Pugh, T.J., Stransky, N., Helman, E., Kim, J., Sougnez, C., Ambrogio, L., Nickerson, E., Shefler, E., Cortés, M.L., Auclair, D., Saksena, G., Voet, D., Noble, M., DiCara, D., Lin, P., Lichtenstein, L., Heiman, D.I., Fennell, T., Imielinski, M., Hernandez, B., Hodis, E., Baca, S., Dulak, A.M., Lohr, J., Landau, D.-A., Wu, C.J., Melendez-Zajgla, J., Hidalgo-Miranda, A., Koren, A., McCarroll, S.A., Mora, J., Lee, R.S., Crompton, B., Onofrio, R., Parkin, M., Winckler, W., Ardlie, K., Gabriel, S.B., Roberts, C.W.M., Biegel, J.A., Stegmaier, K., Bass, A.J., Garraway, L.A., Meyerson, M., Golub, T.R., Gordenin, D.A., Sunyaev, S., Lander, E.S., Getz, G.: Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* **499**(7457), 214–218 (2013). doi:10.1038/nature12213

13. Patch, A.-M., Christie, E.L., Etemadmoghadam, D., Garsed, D.W., George, J., Fereday, S., Nones, K., Cowin, P., Alsop, K., Bailey, P.J., Kassahn, K.S., Newell, F., Quinn, M.C.J., Kazakoff, S., Quek, K., Wilhelm-Benartzi, C., Curry, E., Leong, H.S., Hamilton, A., Mileskhin, L., Au-Yeung, G., Kennedy, C., Hung, J., Chiew, Y.-E., Harnett, P., Friedlander, M., Quinn, M., Pyman, J., Cordner, S., O'Brien, P., Leditschke, J., Young, G., Strachan, K., Waring, P., Azar, W., Mitchell, C., Traficante, N., Hendley, J., Thorne, H., Shackleton, M., Miller, D.K., Arnau, G.M., Tothill, R.W., Holloway, T.P., Semple, T., Harliwong, I., Nourse, C., Nourbakhsh, E., Manning, S., Idrisoglu, S., Bruxner, T.J.C., Christ, A.N., Poudel, B., Holmes, O., Anderson, M., Leonard, C., Lonie, A., Hall, N., Wood, S., Taylor, D.F., Xu, Q., Fink, J.L., Waddell, N., Drapkin, R., Stronach, E., Gabra, H., Brown, R., Jewell, A., Nagaraj, S.H., Markham, E., Wilson, P.J., Ellul, J., McNally, O., Doyle, M.A., Vedururu, R., Stewart, C., Lengyel, E., Pearson, J.V., Waddell, N., deFazio, A., Grimmond, S.M., Bowtell, D.D.L.: Whole-genome characterization of chemoresistant ovarian cancer. *Nature* **521**(7553), 489–494 (2015). doi:10.1038/nature14410

14. Koonin, E.V., Altschul, S.F., Bork, P.: BRCA1 protein products: functional motifs. *Nat Genet* **13**, 266–267 (1996)

Figures

**Figure 1** Sample figure title. A short description of the figure content should go here.

**Figure 2** Sample figure title. Figure legend text.

Tables

**Table 1** Sample table title. This is where the description of the table should go.

	B1	B2	B3
A1	0.1	0.2	0.3
A2	...	..	.
A3	..	.	.

Additional Files

Additional file 1 — Sample additional file title  
Additional file descriptions text (including details of how to view the file, if it is in a non-standard format or the file extension). This might refer to a multi-page table or a figure.

Additional file 2 — Sample additional file title  
Additional file descriptions text.